

wwPDB EM Validation Summary Report (i)

Mar 31, 2021 - 09:16 am BST

EMDB ID : EMD-10244

Title : Structure of the native C. jejuni flagellum filament

Authors: Al-Otaibi, N.S.; Bergeron, J.R.C.

Deposited on : 2019-08-22

Resolution : 8.70 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMMapValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.0.dev75

Validation Pipeline (wwPDB-VP) : 2.18

1 Experimental information (i)

| Property | Value | Source | |
|----------------------------------|---|-----------|--|
| EM reconstruction method | HELICAL | Depositor | |
| Imposed symmetry | HELICAL, twist=65.4°, rise=4.5 Å, axial | Depositor | |
| | sym=C1 | | |
| Number of segments used | 254041 | Depositor | |
| Resolution determination method | FSC 0.143 CUT-OFF | Depositor | |
| CTF correction method | Not provided | | |
| Microscope | FEI TECNAI ARCTICA | Depositor | |
| Voltage (kV) | 200 | Depositor | |
| Electron dose $(e^-/\text{Å}^2)$ | 20.0 | Depositor | |
| Minimum defocus (nm) | Not provided | | |
| Maximum defocus (nm) | Not provided | | |
| Magnification | Not provided | | |
| Image detector | FEI FALCON III (4k x 4k) | Depositor | |
| Maximum map value | 0.150 | Depositor | |
| Minimum map value | -0.074 | Depositor | |
| Average map value | 0.001 | Depositor | |
| Map value standard deviation | 0.011 | Depositor | |
| Recommended contour level | 0.05 | Depositor | |
| Map size (Å) | 609.0, 609.0, 609.0 | wwPDB | |
| Map dimensions | 300, 300, 300 | wwPDB | |
| Map angles (°) | 90.0, 90.0, 90.0 | wwPDB | |
| Pixel spacing (Å) | 2.03, 2.03, 2.03 | Depositor | |




2 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-10244. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

2.1 Orthogonal projections (i)


2.1.1 Primary map





The images above show the map projected in three orthogonal directions.

2.2 Central slices (i)

2.2.1 Primary map








ndox. 100





The images above show central slices of the map in three orthogonal directions.

2.3 Largest variance slices (i)

2.3.1 Primary map







X Index: 169

Y Index: 141


Z Index: 92


The images above show the largest variance slices of the map in three orthogonal directions.

2.4 Orthogonal surface views (i)

2.4.1 Primary map







Y

The images above show the 3D surface view of the map at the recommended contour level 0.05. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



2.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



3 Map analysis (i)

This section contains the results of statistical analysis of the map.


3.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



3.2 Volume estimate (i)




The volume at the recommended contour level is $3136~\mathrm{nm^3}$; this corresponds to an approximate mass of $2833~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



3.3 Rotationally averaged power spectrum (i)




^{*}Reported resolution corresponds to spatial frequency of 0.115 $\rm \AA^{-1}$



4 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

4.1 FSC (i)



*Reported resolution corresponds to spatial frequency of 0.115 $\rm \AA^{-1}$



4.2 Resolution estimates (i)

| Resolution estimate (Å) | \mid Estimation criterion (FSC cut-off) \mid | | |
|---------------------------|--|-------|----------|
| resolution estimate (A) | 0.143 | 0.5 | Half-bit |
| Reported by author | 8.70 | - | - |
| Author-provided FSC curve | 8.56 | 10.06 | 8.64 |
| Calculated* | - | _ | - |

^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

