

# Full wwPDB EM Validation Report (i)

May 17, 2021 – 10:11 am BST

EMDB ID : EMD-11015

Title: CryoEM Structure of CtBP2 Confirms Tetrameric Architecture

Authors : Jecrois, A.M. Deposited on : 2020-05-08

Resolution : 5.00 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at  $\frac{\text{https://www.wwpdb.org/validation/2017/EMMapValidationReportHelp}}{\text{with specific help available everywhere you see the } \vec{\textbf{i}} \text{ symbol.}$ 

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.0.dev75Validation Pipeline (wwPDB-VP) : 2.18

# 1 Experimental information (i)

| Property                         | Value               | Source    |
|----------------------------------|---------------------|-----------|
| EM reconstruction method         | SINGLE PARTICLE     | Depositor |
| Imposed symmetry                 | POINT, D2           | Depositor |
| Number of particles used         | 46426               | Depositor |
| Resolution determination method  | FSC 0.143 CUT-OFF   | Depositor |
| CTF correction method            | Not provided        |           |
| Microscope                       | FEI TITAN KRIOS     | Depositor |
| Voltage (kV)                     | 300                 | Depositor |
| Electron dose $(e^-/\text{Å}^2)$ | 40.0                | Depositor |
| Minimum defocus (nm)             | Not provided        |           |
| Maximum defocus (nm)             | Not provided        |           |
| Magnification                    | 105000.0            | Depositor |
| Image detector                   | GATAN K3 (6k x 4k)  | Depositor |
| Maximum map value                | 0.038               | Depositor |
| Minimum map value                | -0.016              | Depositor |
| Average map value                | 0.000               | Depositor |
| Map value standard deviation     | 0.003               | Depositor |
| Recommended contour level        | 0.0156              | Depositor |
| Map size (Å)                     | 261.0, 261.0, 261.0 | wwPDB     |
| Map dimensions                   | 300, 300, 300       | wwPDB     |
| Map angles (°)                   | 90.0, 90.0, 90.0    | wwPDB     |
| Pixel spacing (Å)                | 0.87, 0.87, 0.87    | Depositor |




## 2 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-11015. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 2.1 Orthogonal projections (i)

#### 2.1.1 Primary map



#### 2.1.2 Raw map





The images above show the map projected in three orthogonal directions.




## 2.2 Central slices (i)

#### 2.2.1 Primary map









X Index: 150


Y Index: 150

Z Index: 150

#### 2.2.2 Raw map









The images above show central slices of the map in three orthogonal directions.

## 2.3 Largest variance slices (i)

#### 2.3.1 Primary map



#### 2.3.2 Raw map




The images above show the largest variance slices of the map in three orthogonal directions.




### 2.4 Orthogonal surface views (i)

#### 2.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0156. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

#### 2.4.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.

### 2.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



## 3 Map analysis (i)

This section contains the results of statistical analysis of the map.


### 3.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



### 3.2 Volume estimate (i)




The volume at the recommended contour level is  $177 \text{ nm}^3$ ; this corresponds to an approximate mass of 160 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 3.3 Rotationally averaged power spectrum (i)



<sup>\*</sup>Reported resolution corresponds to spatial frequency of 0.200  $\rm \mathring{A}^{-1}$ 



# 4 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.

