

### Nov 16, 2021 – 03:01 pm GMT

EM	DB ID	:	EMD-11402				
Title			VAP-A engaged in in vitro membrane contact sites with the OSBP construct				
			N-PH-FFAT at 19.6 A resolution				
Authors		:	de la Mora, E.; diCicco, A.; Dezi, M.; Levy, D.				
Deposited on							
Resolution		:	19.60  Å(reported)				
	This is a Full wwPDB EM Validation Report for a publicly released PDB entry.						

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMMapValidationReportHelp with specific help available everywhere you see the (i) symbol.

EMDB validation analysis : 0.0.0.dev97 Validation Pipeline (wwPDB-VP) : 2.23.2

The following versions of software and data (see references (1)) were used in the production of this report:

# 1 Experimental information (i)

Property	Value	Source
EM reconstruction method	SUBTOMOGRAM AVERAGING	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of subtomograms used	6948	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	Not provided	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	3.5	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	14.678	Depositor
Minimum map value	-15.540	Depositor
Average map value	0.024	Depositor
Map value standard deviation	1.107	Depositor
Recommended contour level	4.12	Depositor
Map size (Å)	353.6, 353.6, 353.6	wwPDB
Map dimensions	52, 52, 52	wwPDB
Map angles ( $^{\circ}$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	6.8, 6.8, 6.8	Depositor




## 2 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-11402. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

## 2.1 Orthogonal projections (i)


### 2.1.1 Primary map




The images above show the map projected in three orthogonal directions.

### 2.2 Central slices (i)


### 2.2.1 Primary map



X Index: 26



Y Index: 26




Z Index: 26



The images above show central slices of the map in three orthogonal directions.


#### Largest variance slices (i) 2.3

#### 2.3.1Primary map



X Index: 27

Y Index: 23



Z Index: 29

The images above show the largest variance slices of the map in three orthogonal directions.

#### Orthogonal surface views (i) $\mathbf{2.4}$

#### 2.4.1Primary map



The images above show the 3D surface view of the map at the recommended contour level 4.12. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



## 2.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



## 3 Map analysis (i)

This section contains the results of statistical analysis of the map.


## 3.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



### 3.2 Volume estimate (i)




The volume at the recommended contour level is  $118 \text{ nm}^3$ ; this corresponds to an approximate mass of 107 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



### 3.3 Rotationally averaged power spectrum (i)




\*Reported resolution corresponds to spatial frequency of 0.051  ${\rm \AA^{-1}}$ 



## 4 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 4.1 FSC (i)



\*Reported resolution corresponds to spatial frequency of 0.051  $\text{\AA}^{-1}$ 



## 4.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
Resolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	19.60	-	-	
Author-provided FSC curve	19.31	26.74	20.28	
Unmasked-calculated*	-	-	-	

\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

