

# Full wwPDB EM Validation Report (i)

Apr 10, 2022 – 09:33 am BST

EMDB ID : EMD-14074

Title: Cryo-EM structure of human full-length extrasynaptic alpha4beta3delta

GABA(A)R in complex with THIP (gaboxadol), histamine and nanobody

Nb25

Authors: Sente, A.; Desai, R.; Naydenova, K.; Malinauskas, T.; Jounaidi, Y.; Miehling,

J.; Zhou, X.; Masiulis, S.; Hardwick, S.W.; Chirgadze, D.Y.; Miller, K.W.;

Aricescu, A.R.

Deposited on : 2021-12-20

Resolution : 2.90 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/EMMapValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.0.dev97

Validation Pipeline (wwPDB-VP) : 2.27

# 1 Experimental information (i)

Property	Value	Source	
EM reconstruction method	SINGLE PARTICLE	Depositor	
Imposed symmetry	Not Provided		
Number of particles used	61854	Depositor	
Resolution determination method	FSC 0.143 CUT-OFF	Depositor	
CTF correction method	Not provided		
Microscope	FEI TITAN KRIOS	Depositor	
Voltage (kV)	300	Depositor	
Electron dose $(e^-/\text{Å}^2)$	40.0	Depositor	
Minimum defocus (nm)	0.7000000000000001	Depositor	
Maximum defocus (nm)	2.1	Depositor	
Magnification	130000.0	Depositor	
Image detector	GATAN K3 (6k x 4k)	Depositor	
Maximum map value	0.107	Depositor	
Minimum map value	-0.040	Depositor	
Average map value	0.000	Depositor	
Map value standard deviation	0.003	Depositor	
Recommended contour level	0.008	Depositor	
Map size (Å)	259.84, 259.84, 259.84	wwPDB	
Map dimensions	290, 290, 290	wwPDB	
Map angles (°)	90.0, 90.0, 90.0	wwPDB	
Pixel spacing (Å)	0.89599997, 0.89599997, 0.89599997	Depositor	




# 2 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-14074. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 2.1 Orthogonal projections (i)

### 2.1.1 Primary map




The images above show the map projected in three orthogonal directions.

### 2.2 Central slices (i)


### 2.2.1 Primary map



X Index: 145



Y Index: 145




Z Index: 145





The images above show central slices of the map in three orthogonal directions.

#### Largest variance slices (i) 2.3

#### 2.3.1 Primary map







X Index: 155

Y Index: 163

Z Index: 121

The images above show the largest variance slices of the map in three orthogonal directions.

#### Orthogonal surface views (i) 2.4

#### 2.4.1 Primary map







 $\mathbf{Z}$ 

The images above show the 3D surface view of the map at the recommended contour level 0.008. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



# 2.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



# 3 Map analysis (i)

This section contains the results of statistical analysis of the map.


### 3.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



### 3.2 Volume estimate (i)




The volume at the recommended contour level is  $278~\mathrm{nm}^3$ ; this corresponds to an approximate mass of  $251~\mathrm{kDa}$ .

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



## 3.3 Rotationally averaged power spectrum (i)




<sup>\*</sup>Reported resolution corresponds to spatial frequency of 0.345  $\rm \mathring{A}^{-1}$ 



# 4 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 4.1 FSC (i)



\*Reported resolution corresponds to spatial frequency of 0.345  $\rm \AA^{-1}$ 



# 4.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
rtesolution estimate (A)	0.143	0.5	Half-bit
Reported by author	2.90	-	-
Author-provided FSC curve	2.89	3.36	2.93
Unmasked-calculated*	-	-	-

<sup>\*</sup>Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

