

Mar 31, 2021 - 09:58 am BST

		EMD-1593 Electron crystallographic reconstruction of human copper transporter (hCTR1).
Deposited on	:	DeFeo, C.J.; Aller, S.G.; Siluvai, G.S.; Blackburn, N.J.; Unger, V.M.
This is	a l	Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/EMMapValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

1 Experimental information (i)

Property	Value	Source
EM reconstruction method	CRYSTALLOGRAPHY	Depositor
Imposed symmetry	2D CRYSTAL, $a=89$ Å, $b=89$ Å, $c=400$ Å,	Depositor
	$\gamma = 120^{\circ}$, space group=Not provided	
Number of images used	Not provided	
Resolution determination method	OTHER	Depositor
CTF correction method	Each image	Depositor
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	200	Depositor
Electron dose $(e^-/\text{\AA}^2)$	10	Depositor
Minimum defocus (nm)	0.4	Depositor
Maximum defocus (nm)	1.5	Depositor
Magnification	50000.0	Depositor
Image detector	KODAK SO-163 FILM	Depositor
Maximum map value	250.000	Depositor
Minimum map value	-211.531	Depositor
Average map value	-2.757	Depositor
Map value standard deviation	31.055	Depositor
Recommended contour level	75.0	Depositor
Map size (Å)	88.99969, 88.99969, 400.0	wwPDB
Map dimensions	123, 123, 401	wwPDB
Map angles (°)	90.0, 90.0, 120.0	wwPDB
Pixel spacing (Å)	1.01136, 1.01136, 1.0	Depositor




2 Map visualisation (i)


This section contains visualisations of the EMDB entry EMD-1593. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

2.1 Orthogonal projections (i)

2.1.1 Primary map





 \mathbf{Z}


The images above show the map projected in three orthogonal directions.




2.2 Central slices (i)

2.2.1 Primary map







Z Index: 200

X Index: 61


The images above show central slices of the map in three orthogonal directions.

2.3 Largest variance slices (i)


2.3.1 Primary map



X Index: 61



Y Index: 61




Z Index: 172

The images above show the largest variance slices of the map in three orthogonal directions.



2.4 Orthogonal surface views (i)

2.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 75.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

2.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



3 Map analysis (i)

This section contains the results of statistical analysis of the map.


3.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



3.2 Volume estimate (i)



The volume at the recommended contour level is 138 nm^3 ; this corresponds to an approximate mass of 125 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

3.3 Rotationally averaged power spectrum (i)

This section was not generated. The rotationally averaged power spectrum is only generated for cubic maps.



4 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.

