

# wwPDB EM Validation Summary Report (i)

Dec 21, 2020 – 05:52 PM EST

EMDB ID : EMD-1724

Title : Ribosome dynamics and tRNA movement as visualized by time-resolved elec-

tron cryomicroscopy

Authors: Fischer, N.; Konevega, A.L.; Wintermeyer, W.; Rodnina, M.V.; Stark, H.

Deposited on : 2010-04-30

Resolution : 9.00 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

The following versions of software and data (see references (i)) were used in the production of this report:

EMDB validation analysis : 0.0.0.dev61Validation Pipeline (wwPDB-VP) : 2.16

# 1 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	Not Provided	
Number of particles used	26429	Depositor
Resolution determination method	FSC 0.5 CUT-OFF	Depositor
CTF correction method	Local	Depositor
Microscope	FEI/PHILIPS CM200FEG	Depositor
Voltage (kV)	160	Depositor
Electron dose $(e^-/\text{Å}^2)$	20	Depositor
Minimum defocus (nm)	0.5	Depositor
Maximum defocus (nm)	2.0	Depositor
Magnification	162740.0	Depositor
Image detector	GENERIC TVIPS (4k x 4k)	Depositor
Maximum map value	223.953	Depositor
Minimum map value	-122.023	Depositor
Average map value	-0.756	Depositor
Map value standard deviation	22.478	Depositor
Recommended contour level	40.0	Depositor
Map size (Å)	359.04, 359.04, 359.04	wwPDB
Map dimensions	192, 192, 192	wwPDB
Map angles (°)	90, 90, 90	wwPDB
Pixel spacing (Å)	1.87, 1.87, 1.87	Depositor




## 2 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-1724. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 2.1 Orthogonal projections (i)


#### 2.1.1 Primary map



The images above show the map projected in three orthogonal directions.

#### 2.2 Central slices (i)

#### 2.2.1 Primary map






The images above show central slices of the map in three orthogonal directions.

### 2.3 Largest variance slices (i)


#### 2.3.1 Primary map



The images above show the largest variance slices of the map in three orthogonal directions.

## 2.4 Orthogonal surface views (i)

#### 2.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 40.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



# 2.5 Mask visualisation (i)


This section was not generated. No masks/segmentation were deposited.



# 3 Map analysis (i)

This section contains the results of statistical analysis of the map.


### 3.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



### 3.2 Volume estimate (i)




The volume at the recommended contour level is  $1986~\mathrm{nm^3}$ ; this corresponds to an approximate mass of  $1794~\mathrm{kDa}$ .

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



## 3.3 Rotationally averaged power spectrum (i)



<sup>\*</sup>Reported resolution corresponds to spatial frequency of 0.111  $\rm \AA^{-1}$ 



# 4 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.

