



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 2, 2024 – 06:48 pm GMT

PDB ID : 5A0M
Title : THE CRYSTAL STRUCTURE OF I-SCEI IN COMPLEX WITH ITS TARGET DNA IN THE PRESENCE OF MN
Authors : Prieto, J.; Redondo, P.; Merino, N.; Villate, M.; Blanco, F.J.; Molina, R.
Deposited on : 2015-04-21
Resolution : 2.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2172 (2.90-2.90)
Ramachandran outliers	138981	2115 (2.90-2.90)
Sidechain outliers	138945	2117 (2.90-2.90)
RSRZ outliers	127900	1906 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	235	
2	B	235	
3	C	14	
3	E	14	
4	D	16	
4	H	16	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	I	9	 67% 33%
5	L	9	 44% 44% 11%
6	J	11	 55% 45%
6	K	11	 36% 55% 9%

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 5761 atoms, of which 13 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INTRON-ENCODED ENDONUCLEASE I-SCEI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	224	1862	1213	308	330	11	0	0	0

- Molecule 2 is a protein called INTRON-ENCODED ENDONUCLEASE I-SCEI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	227	1874	1220	309	334	11	0	0	0

- Molecule 3 is a DNA chain called 5'-D(*CP*AP*CP*GP*CP*TP*AP*GP*GP*GP*AP*T P*AP*AP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	13	268	128	55	73	12	0	0	0
3	E	14	287	137	58	79	13	0	0	0

- Molecule 4 is a DNA chain called 5'-D(*GP*GP*TP*AP*TP*TP*AP*CP*CP*CP*TP*G P*TP*TP *AP*TP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	D	15	304	147	48	94	15	0	0	0
4	H	15	301	147	48	92	14	0	0	0

- Molecule 5 is a DNA chain called 5'-D(*CP*CP*CP*TP*AP*GP*CP*GP*TP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
5	I	9	182	86	31	56	9	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
5	L	9	Total	C	H	N	O	P	0	0	0
			194	86	12	31	56	9			

- Molecule 6 is a DNA chain called 5'-D(*CP*AP*GP*GP*GP*TP*AP*AP*TP*AP*CP)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
6	J	11	Total	C	N	O	P	0	0	0	
			229	108	45	65	11				
6	K	11	Total	C	H	N	O	P	0	0	0
			230	108	1	45	65	11			

- Molecule 7 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	3	Total	Mn	0	0
			3	3		
7	B	3	Total	Mn	0	0
			3	3		
7	D	1	Total	Mn	0	0
			1	1		
7	H	1	Total	Mn	0	0
			1	1		


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	8	Total	O	0	0
			8	8		
8	B	5	Total	O	0	0
			5	5		
8	C	1	Total	O	0	0
			1	1		
8	D	2	Total	O	0	0
			2	2		
8	E	2	Total	O	0	0
			2	2		
8	H	1	Total	O	0	0
			1	1		
8	I	3	Total	O	0	0
			3	3		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: INTRON-ENCODED ENDONUCLEASE I-SCEI




- Molecule 2: INTRON-ENCODED ENDONUCLEASE I-SCEI




- Molecule 3: 5'-D(*CP*AP*CP*GP*CP*TP*AP*GP*GP*GP*AP*TP*AP*AP)-3'




- Molecule 3: 5'-D(*CP*AP*CP*GP*CP*TP*AP*GP*GP*GP*AP*TP*AP*AP)-3'






- Molecule 4: 5'-D(*GP*GP*TP*AP*TP*TP*AP*CP*CP*CP*TP*GP*TP*TP *AP*TP)-3'

Chain D: 6% 75% 12% 6%



- Molecule 4: 5'-D(*GP*GP*TP*AP*TP*TP*AP*CP*CP*CP*TP*GP*TP*TP *AP*TP)-3'

Chain H: 19% 62% 12% 6%



- Molecule 5: 5'-D(*CP*CP*CP*TP*AP*GP*CP*GP*TP)-3'

Chain I: 67% 33%



- Molecule 5: 5'-D(*CP*CP*CP*TP*AP*GP*CP*GP*TP)-3'

Chain L: 44% 44% 11%



- Molecule 6: 5'-D(*CP*AP*GP*GP*GP*TP*AP*AP*TP*AP*CP)-3'

Chain J: 55% 45%



- Molecule 6: 5'-D(*CP*AP*GP*GP*GP*TP*AP*AP*TP*AP*CP)-3'

Chain K: 36% 55% 9%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	80.11Å 80.56Å 130.87Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.90 – 2.90 42.90 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (42.90-2.90) 100.0 (42.90-2.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.56 (at 2.90Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.186 , 0.249 0.199 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	61.6	Xtrriage
Anisotropy	0.954	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 36.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.057 for k,h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5761	wwPDB-VP
Average B, all atoms (Å ²)	69.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/1909	0.57	0/2580
2	B	0.50	0/1921	0.60	0/2599
3	C	1.14	1/302 (0.3%)	1.05	0/465
3	E	1.91	10/323 (3.1%)	1.57	5/497 (1.0%)
4	D	0.99	0/338	1.29	3/519 (0.6%)
4	H	1.27	3/335 (0.9%)	1.37	2/515 (0.4%)
5	I	1.35	1/202 (0.5%)	1.21	0/307
5	L	1.10	1/202 (0.5%)	1.10	0/307
6	J	1.21	1/257 (0.4%)	1.07	0/393
6	K	1.18	1/257 (0.4%)	0.99	1/393 (0.3%)
All	All	0.88	18/6046 (0.3%)	0.90	11/8575 (0.1%)

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	I	17	DC	OP3-P	-11.43	1.47	1.61
3	E	2	DA	P-O5'	10.86	1.70	1.59
6	K	15	DC	OP3-P	-10.68	1.48	1.61
6	J	15	DC	OP3-P	-10.63	1.48	1.61
3	E	1	DC	C2-O2	10.31	1.33	1.24

The worst 5 of 11 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	E	2	DA	O5'-P-OP2	-18.32	88.72	110.70
3	E	1	DC	O4'-C1'-N1	10.37	115.26	108.00
4	D	16	DT	O4'-C4'-C3'	-9.06	100.57	106.00
3	E	2	DA	OP1-P-OP2	8.49	132.33	119.60
3	E	2	DA	O4'-C1'-N9	6.61	112.63	108.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1862	0	1876	60	0
2	B	1874	0	1877	73	0
3	C	268	0	147	9	0
3	E	287	0	158	22	1
4	D	304	0	173	24	0
4	H	301	0	174	19	1
5	I	182	0	102	1	0
5	L	182	12	102	12	0
6	J	229	0	124	3	0
6	K	229	1	124	11	0
7	A	3	0	0	0	0
7	B	3	0	0	0	0
7	D	1	0	0	0	0
7	H	1	0	0	0	0
8	A	8	0	0	2	0
8	B	5	0	0	0	0
8	C	1	0	0	0	0
8	D	2	0	0	0	0
8	E	2	0	0	0	0
8	H	1	0	0	0	0
8	I	3	0	0	0	0
All	All	5748	13	4857	201	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 201 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:2:DA:H8	3:E:2:DA:H5''	1.11	1.13
3:E:2:DA:H5''	3:E:2:DA:C8	1.85	1.11
3:C:5:DC:H2''	3:C:6:DT:H5''	1.37	1.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:L:23:DC:H2''	5:L:24:DG:H5'	1.41	0.98
2:B:480:ARG:NH1	2:B:487:CYS:O	2.00	0.94

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:1:DC:N4	4:H:2:DG:N7[1_545]	1.86	0.34

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	222/235 (94%)	201 (90%)	20 (9%)	1 (0%)	29	61
2	B	225/235 (96%)	211 (94%)	13 (6%)	1 (0%)	34	66
All	All	447/470 (95%)	412 (92%)	33 (7%)	2 (0%)	34	66

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	463	ASN
1	A	525	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	207/218 (95%)	192 (93%)	15 (7%)	14	39
2	B	207/218 (95%)	195 (94%)	12 (6%)	20	50
All	All	414/436 (95%)	387 (94%)	27 (6%)	17	45

5 of 27 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	504	SER
2	B	350	ARG
2	B	503	MET
2	B	349	SER
2	B	373	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	364	ASN
2	B	377	GLN
2	B	402	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 8 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/235 (95%)	0.42	13 (5%) 23 19	52, 76, 108, 130	0
2	B	227/235 (96%)	0.07	1 (0%) 92 93	44, 64, 94, 132	0
3	C	13/14 (92%)	-0.02	0 100 100	51, 72, 95, 96	0
3	E	14/14 (100%)	-0.15	1 (7%) 16 12	41, 61, 80, 90	0
4	D	15/16 (93%)	-0.56	0 100 100	57, 63, 83, 92	0
4	H	15/16 (93%)	-0.53	0 100 100	45, 56, 72, 74	0
5	I	9/9 (100%)	-0.60	0 100 100	43, 63, 76, 82	0
5	L	9/9 (100%)	-0.30	0 100 100	58, 79, 98, 116	0
6	J	11/11 (100%)	-0.18	0 100 100	44, 62, 73, 74	0
6	K	11/11 (100%)	-0.32	0 100 100	49, 72, 80, 91	0
All	All	548/570 (96%)	0.14	15 (2%) 54 50	41, 69, 100, 132	0

The worst 5 of 15 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	456	THR	5.8
1	A	352	GLU	5.1
1	A	454	ASN	4.8
1	A	508	PHE	3.9
1	A	312	LEU	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	MN	A	1528	1/1	0.96	0.20	56,56,56,56	0
7	MN	A	1529	1/1	0.96	0.15	56,56,56,56	0
7	MN	B	1530	1/1	0.97	0.18	49,49,49,49	0
7	MN	B	1531	1/1	0.97	0.20	52,52,52,52	0
7	MN	H	1017	1/1	0.97	0.24	61,61,61,61	0
7	MN	B	1529	1/1	0.98	0.20	57,57,57,57	0
7	MN	D	1017	1/1	0.98	0.18	64,64,64,64	0
7	MN	A	1527	1/1	0.98	0.19	63,63,63,63	0

6.5 Other polymers [i](#)

There are no such residues in this entry.