



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 21, 2022 – 04:59 pm GMT

PDB ID : 7A0X  
Title : Structure of dimeric sodium proton antiporter NhaA, at pH 6.0, crystallized with chimeric Fab antibodies  
Authors : Fippel, A.; Lentes, C.J.; Mir, S.H.; Wirth, C.; Hunte, C.  
Deposited on : 2020-08-11  
Resolution : 2.37 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.26  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.26


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5509 (2.40-2.36)
Clashscore	141614	6082 (2.40-2.36)
Ramachandran outliers	138981	5973 (2.40-2.36)
Sidechain outliers	138945	5975 (2.40-2.36)
RSRZ outliers	127900	5397 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	
1	B	406	
2	C	252	
2	E	252	
3	D	210	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	210	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	PG4	E	301	-	-	-	X

## 2 Entry composition [i](#)

There are 11 unique types of molecules in this entry. The entry contains 12318 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Na(+)/H(+) antiporter NhaA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	370	Total	C	N	O	S	55	2	0
			2783	1855	449	465	14			
1	B	373	Total	C	N	O	S	58	4	0
			2826	1890	457	465	14			

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	389	SER	-	expression tag	UNP Q8ZRZ3
A	390	GLU	-	expression tag	UNP Q8ZRZ3
A	391	ASN	-	expression tag	UNP Q8ZRZ3
A	392	LEU	-	expression tag	UNP Q8ZRZ3
A	393	TYR	-	expression tag	UNP Q8ZRZ3
A	394	PHE	-	expression tag	UNP Q8ZRZ3
A	395	GLN	-	expression tag	UNP Q8ZRZ3
A	396	GLY	-	expression tag	UNP Q8ZRZ3
A	397	GLY	-	expression tag	UNP Q8ZRZ3
A	398	ARG	-	expression tag	UNP Q8ZRZ3
A	399	GLY	-	expression tag	UNP Q8ZRZ3
A	400	SER	-	expression tag	UNP Q8ZRZ3
A	401	HIS	-	expression tag	UNP Q8ZRZ3
A	402	HIS	-	expression tag	UNP Q8ZRZ3
A	403	HIS	-	expression tag	UNP Q8ZRZ3
A	404	HIS	-	expression tag	UNP Q8ZRZ3
A	405	HIS	-	expression tag	UNP Q8ZRZ3
A	406	HIS	-	expression tag	UNP Q8ZRZ3
B	389	SER	-	expression tag	UNP Q8ZRZ3
B	390	GLU	-	expression tag	UNP Q8ZRZ3
B	391	ASN	-	expression tag	UNP Q8ZRZ3
B	392	LEU	-	expression tag	UNP Q8ZRZ3
B	393	TYR	-	expression tag	UNP Q8ZRZ3
B	394	PHE	-	expression tag	UNP Q8ZRZ3
B	395	GLN	-	expression tag	UNP Q8ZRZ3

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	396	GLY	-	expression tag	UNP Q8ZRZ3
B	397	GLY	-	expression tag	UNP Q8ZRZ3
B	398	ARG	-	expression tag	UNP Q8ZRZ3
B	399	GLY	-	expression tag	UNP Q8ZRZ3
B	400	SER	-	expression tag	UNP Q8ZRZ3
B	401	HIS	-	expression tag	UNP Q8ZRZ3
B	402	HIS	-	expression tag	UNP Q8ZRZ3
B	403	HIS	-	expression tag	UNP Q8ZRZ3
B	404	HIS	-	expression tag	UNP Q8ZRZ3
B	405	HIS	-	expression tag	UNP Q8ZRZ3
B	406	HIS	-	expression tag	UNP Q8ZRZ3


- Molecule 2 is a protein called chimeric antibody Fab-F6, heavy chain, chimeric antibody Fab-F6, heavy chain, chimeric antibody Fab-F6, heavy chain, chimeric antibody Fab-F6, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	C	210	Total 1556	C 977	N 268	O 306	S 5	42	2	0
2	E	223	Total 1631	C 1019	N 280	O 327	S 5	5	1	0

- Molecule 3 is a protein called chimeric antibody Fab-F6, light chain, chimeric antibody Fab-F6, light chain, chimeric antibody Fab-F6, light chain, chimeric antibody Fab-F6, light chain.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	D	207	Total 1555	C 972	N 263	O 316	S 4	20	0	0
3	F	209	Total 1568	C 979	N 265	O 320	S 4	4	0	0

- Molecule 4 is CARDIOLIPIN (three-letter code: CDL) (formula: C<sub>81</sub>H<sub>156</sub>O<sub>17</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
4	A	1	54	35	17	2	0	0

- Molecule 5 is 1,2-DIACYL-GLYCEROL-3-SN-PHOSPHATE (three-letter code: 3PH) (formula:  $C_{39}H_{77}O_8P$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
5	A	1	27	18	8	1	0	0
5	B	1	27	18	8	1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
5	B	1	22	13	8	1	0	0

- Molecule 6 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
6	A	1	5	4	1	0	0
6	B	1	5	4	1	0	0

- Molecule 7 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: C<sub>8</sub>H<sub>18</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			13	8	5		
7	B	1	Total	C	O	0	0
			13	8	5		
7	E	1	Total	C	O	0	0
			13	8	5		
7	E	1	Total	C	O	0	0
			13	8	5		

- Molecule 8 is 1,2-Distearoyl-sn-glycerophosphoethanolamine (three-letter code: 3PE) (formula:  $C_{41}H_{82}NO_8P$ ).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
8	B	1	Total	C	N	O	P	0	0
			26	16	1	8	1		

- Molecule 9 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	C	2	Total	Cl	0	0
			2	2		
9	F	1	Total	Cl	0	0
			1	1		

- Molecule 10 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	F	1	Total	Ca	0	0
			1	1		


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	8	Total	O	0	0
			8	8		
11	B	3	Total	O	0	0
			3	3		
11	C	29	Total	O	0	0
			29	29		
11	D	34	Total	O	0	0
			34	34		
11	E	51	Total	O	0	0
			51	51		
11	F	52	Total	O	0	0
			52	52		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Na(+)/H(+) antiporter NhaA




- Molecule 1: Na(+)/H(+) antiporter NhaA




- Molecule 2: chimeric antibody Fab-F6, heavy chain, chimeric antibody Fab-F6, heavy chain, chimeric antibody Fab-F6, heavy chain, chimeric antibody Fab-F6, heavy chain







- Molecule 2: chimeric antibody Fab-F6, heavy chain, chimeric antibody Fab-F6, heavy chain, chimeric antibody Fab-F6, heavy chain, chimeric antibody Fab-F6, heavy chain



- Molecule 3: chimeric antibody Fab-F6, light chain, chimeric antibody Fab-F6, light chain, chimeric antibody Fab-F6, light chain, chimeric antibody Fab-F6, light chain



- Molecule 3: chimeric antibody Fab-F6, light chain, chimeric antibody Fab-F6, light chain, chimeric antibody Fab-F6, light chain, chimeric antibody Fab-F6, light chain



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	113.01Å 91.71Å 140.00Å 90.00° 109.80° 90.00°	Depositor
Resolution (Å)	24.90 – 2.37 24.90 – 2.37	Depositor EDS
% Data completeness (in resolution range)	63.1 (24.90-2.37) 63.2 (24.90-2.37)	Depositor EDS
$R_{merge}$	0.16	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.70 (at 2.36Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, $R_{free}$	0.205 , 0.241 0.204 , 0.239	Depositor DCC
$R_{free}$ test set	2222 reflections (3.20%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	63.8	Xtrriage
Anisotropy	0.018	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	12318	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	77.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PG4, 3PE, PO4, 3PH, CA, CDL, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.24	0/2848	0.41	0/3884
1	B	0.24	0/2903	0.41	0/3960
2	C	0.24	0/1593	0.46	0/2167
2	E	0.24	0/1670	0.46	0/2272
3	D	0.24	0/1590	0.44	0/2165
3	F	0.24	0/1603	0.45	0/2182
All	All	0.24	0/12207	0.43	0/16630

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2783	0	2953	29	0
1	B	2826	0	3014	40	0
2	C	1556	0	1525	20	0
2	E	1631	0	1595	15	0
3	D	1555	0	1511	11	0
3	F	1568	0	1520	9	0
4	A	54	0	52	4	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	27	0	27	0	0
5	B	49	0	44	2	0
6	A	5	0	0	0	0
6	B	5	0	0	0	0
7	A	13	0	18	0	0
7	B	13	0	18	0	0
7	E	26	0	36	2	0
8	B	26	0	26	1	0
9	C	2	0	0	1	0
9	F	1	0	0	0	0
10	F	1	0	0	0	0
11	A	8	0	0	0	0
11	B	3	0	0	0	0
11	C	29	0	0	0	0
11	D	34	0	0	0	0
11	E	51	0	0	1	0
11	F	52	0	0	0	0
All	All	12318	0	12339	122	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 122 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:10:ALA:HB2	3:D:16:VAL:HG23	1.72	0.72
2:C:52:TYR:OH	2:C:102:ARG:NH1	2.26	0.68
3:D:182:TYR:O	3:D:207:ARG:NH2	2.27	0.68
3:D:33:LYS:NZ	3:D:76:GLU:O	2.25	0.67
1:A:88:LEU:HA	1:A:94:ALA:HB2	1.79	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	370/406 (91%)	345 (93%)	25 (7%)	0	100	100
1	B	375/406 (92%)	348 (93%)	27 (7%)	0	100	100
2	C	208/252 (82%)	200 (96%)	8 (4%)	0	100	100
2	E	222/252 (88%)	214 (96%)	8 (4%)	0	100	100
3	D	205/210 (98%)	194 (95%)	11 (5%)	0	100	100
3	F	207/210 (99%)	198 (96%)	9 (4%)	0	100	100
All	All	1587/1736 (91%)	1499 (94%)	88 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	286/314 (91%)	279 (98%)	7 (2%)	49	66
1	B	290/314 (92%)	282 (97%)	8 (3%)	43	61
2	C	168/199 (84%)	166 (99%)	2 (1%)	71	84
2	E	177/199 (89%)	173 (98%)	4 (2%)	50	68
3	D	176/178 (99%)	173 (98%)	3 (2%)	60	76
3	F	177/178 (99%)	176 (99%)	1 (1%)	86	93
All	All	1274/1382 (92%)	1249 (98%)	25 (2%)	55	72

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	333	ILE
3	D	16	VAL
3	F	162	GLN
2	C	220	VAL
3	D	73	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 4 are monoatomic - leaving 11 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	PO4	A	503	-	4,4,4	0.91	0	6,6,6	0.43	0
5	3PH	B	502	-	21,21,47	0.29	0	25,26,52	0.72	1 (4%)
8	3PE	B	503	-	25,25,50	0.41	0	28,30,55	0.39	0
7	PG4	B	505	-	12,12,12	0.45	0	11,11,11	0.29	0
5	3PH	B	501	-	26,26,47	0.28	0	30,31,52	0.44	0
7	PG4	A	504	-	12,12,12	0.45	0	11,11,11	0.29	0
7	PG4	E	302	-	12,12,12	0.43	0	11,11,11	0.43	0
7	PG4	E	301	-	12,12,12	0.45	0	11,11,11	0.36	0
6	PO4	B	504	-	4,4,4	0.92	0	6,6,6	0.42	0
4	CDL	A	501	-	53,53,99	1.26	4 (7%)	59,65,111	1.22	6 (10%)
5	3PH	A	502	-	26,26,47	0.26	0	30,31,52	0.46	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	3PH	B	502	-	-	4/23/23/49	-
8	3PE	B	503	-	-	8/29/29/54	-
7	PG4	B	505	-	-	1/10/10/10	-
5	3PH	B	501	-	-	2/28/28/49	-
7	PG4	A	504	-	-	1/10/10/10	-
7	PG4	E	302	-	-	2/10/10/10	-
7	PG4	E	301	-	-	2/10/10/10	-
4	CDL	A	501	-	-	25/64/64/110	-
5	3PH	A	502	-	-	6/28/28/49	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	501	CDL	OA8-CA7	4.38	1.46	1.33
4	A	501	CDL	OB8-CB7	4.24	1.45	1.33
4	A	501	CDL	OA6-CA5	4.17	1.46	1.34
4	A	501	CDL	OB6-CB5	4.12	1.45	1.34

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	501	CDL	OB6-CB5-C51	4.15	120.45	111.50
4	A	501	CDL	OA6-CA5-C11	3.53	119.12	111.50
4	A	501	CDL	OA8-CA7-C31	2.79	120.66	111.91
4	A	501	CDL	CA4-OA6-CA5	-2.59	111.41	117.79
4	A	501	CDL	OB8-CB7-C71	2.47	119.67	111.91

There are no chirality outliers.

5 of 51 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	501	CDL	CB2-C1-CA2-OA2
4	A	501	CDL	C1-CA2-OA2-PA1
4	A	501	CDL	CA2-OA2-PA1-OA4
4	A	501	CDL	CA3-OA5-PA1-OA3
4	A	501	CDL	CB2-OB2-PB2-OB4



There are no ring outliers.


6 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	502	3PH	1	0
8	B	503	3PE	1	0
5	B	501	3PH	1	0
7	E	302	PG4	2	0
7	E	301	PG4	1	0
4	A	501	CDL	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	370/406 (91%)	0.54	46 (12%) 4 4	42, 77, 131, 204	13 (3%)
1	B	373/406 (91%)	0.63	59 (15%) 2 2	47, 87, 144, 184	14 (3%)
2	C	207/252 (82%)	0.97	42 (20%) 1 1	32, 63, 136, 156	6 (2%)
2	E	223/252 (88%)	0.49	28 (12%) 3 4	33, 54, 93, 157	2 (0%)
3	D	207/210 (98%)	0.94	44 (21%) 0 1	36, 68, 145, 165	6 (2%)
3	F	209/210 (99%)	0.11	17 (8%) 12 13	34, 52, 79, 107	1 (0%)
All	All	1589/1736 (91%)	0.61	236 (14%) 2 2	32, 71, 136, 204	42 (2%)

The worst 5 of 236 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	325	TYR	10.0
1	A	324	THR	8.8
2	E	141	SER	7.6
1	A	244	GLY	7.5
1	A	245	ARG	7.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	3PH	A	502	27/48	0.71	0.34	67,97,137,147	0
7	PG4	E	302	13/13	0.74	0.26	88,99,112,112	0
8	3PE	B	503	26/51	0.75	0.35	60,112,142,143	0
7	PG4	E	301	13/13	0.77	0.42	67,78,90,93	0
5	3PH	B	501	27/48	0.81	0.27	75,95,146,157	0
7	PG4	B	505	13/13	0.85	0.26	56,70,83,87	0
6	PO4	B	504	5/5	0.88	0.13	85,95,105,128	0
4	CDL	A	501	54/100	0.90	0.20	68,94,155,156	0
7	PG4	A	504	13/13	0.91	0.12	44,54,80,82	0
5	3PH	B	502	22/48	0.92	0.15	76,93,106,121	0
6	PO4	A	503	5/5	0.94	0.10	85,91,100,116	0
9	CL	C	302	1/1	0.94	0.18	76,76,76,76	0
10	CA	F	302	1/1	0.97	0.18	97,97,97,97	0
9	CL	F	301	1/1	0.99	0.25	59,59,59,59	0
9	CL	C	301	1/1	0.99	0.12	59,59,59,59	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around 3PH A 502:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around 3PE B 503:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)









**Electron density around 3PH B 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around CDL A 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.