



Full wwPDB X-ray Structure Validation Report ⓘ

Dec 12, 2023 – 09:45 pm GMT

PDB ID : 4A2H
Title : Crystal Structure of Laccase from *Corioloopsis gallica* pH 7.0
Authors : De La Mora, E.; Valderrama, B.; Horjales, E.; Rudino-Pinera, E.
Deposited on : 2011-09-27
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	496	 2% 92% 7%
2	B	2	 100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	BMA	A	1555	-	-	-	X

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 4135 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LACCASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	496	3727	2368	631	719	9	0	0	0

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	39	ASP	TYR	engineered mutation	UNP Q1W6B1
A	151	ASN	GLN	engineered mutation	UNP Q1W6B1
A	178	LYS	ARG	engineered mutation	UNP Q1W6B1
A	182	PRO	ALA	engineered mutation	UNP Q1W6B1
A	183	VAL	ILE	engineered mutation	UNP Q1W6B1
A	229	TYR	HIS	engineered mutation	UNP Q1W6B1
A	256	LEU	ILE	engineered mutation	UNP Q1W6B1
A	287	THR	ASN	engineered mutation	UNP Q1W6B1
A	288	GLN	THR	engineered mutation	UNP Q1W6B1
A	294	THR	VAL	engineered mutation	UNP Q1W6B1
A	314	THR	ALA	engineered mutation	UNP Q1W6B1
A	329	LYS	GLU	engineered mutation	UNP Q1W6B1
A	356	ASN	ARG	engineered mutation	UNP Q1W6B1
A	358	THR	SER	engineered mutation	UNP Q1W6B1
A	423	VAL	ALA	engineered mutation	UNP Q1W6B1

- Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	2	28	16	2	10	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	A	1	14	8	1	5	0	0
3	A	1	14	8	1	5	0	0

- Molecule 4 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cu		
4	A	4	4	4	0	0

- Molecule 5 is beta-D-mannopyranose (three-letter code: BMA) (formula: $C_6H_{12}O_6$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 12 6 6	0	0
5	A	1	Total C O 12 6 6	0	0
5	A	1	Total C O 12 6 6	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	312	Total O 312 312	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LACCASE



- Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	56.47Å 86.21Å 152.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.26 – 2.30 34.26 – 2.30	Depositor EDS
% Data completeness (in resolution range)	95.3 (34.26-2.30) 95.3 (34.26-2.30)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.94 (at 2.29Å)	Xtrriage
Refinement program	REFMAC 5.5.0110	Depositor
R, R_{free}	0.189 , 0.225 0.187 , 0.220	Depositor DCC
R_{free} test set	1642 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	31.0	Xtrriage
Anisotropy	0.160	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 25.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4135	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.05% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, CU, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.87	3/3838 (0.1%)	0.67	3/5277 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	343	ASP	CG-OD1	42.43	2.23	1.25
1	A	343	ASP	CG-OD2	8.51	1.45	1.25
1	A	343	ASP	CB-CG	-7.10	1.36	1.51

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	343	ASP	OD1-CG-OD2	-14.24	96.25	123.30
1	A	343	ASP	CB-CG-OD1	11.87	128.98	118.30
1	A	343	ASP	CB-CG-OD2	11.41	128.57	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	343	ASP	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3727	0	3541	27	0
2	B	28	0	25	0	0
3	A	28	0	26	2	0
4	A	4	0	0	0	0
5	A	36	0	36	2	0
6	A	312	0	0	1	0
All	All	4135	0	3628	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:329:LYS:H	1:A:329:LYS:HE2	1.32	0.93
1:A:343:ASP:OD1	1:A:343:ASP:CG	2.22	0.78
1:A:276:ASN:HD21	1:A:300:ARG:HE	1.32	0.74
1:A:242:GLU:HB3	1:A:265:SER:HB2	1.71	0.72
5:A:1553:BMA:H3	5:A:1554:BMA:H4	1.73	0.70
1:A:284:ASN:H	1:A:284:ASN:HD22	1.41	0.69
1:A:302:ASP:C	1:A:302:ASP:N	2.47	0.69
1:A:276:ASN:ND2	1:A:300:ARG:HE	1.93	0.66
1:A:33:ASN:HD22	1:A:74:THR:H	1.48	0.61
1:A:422:HIS:HE1	1:A:487:GLU:OE1	1.82	0.61
1:A:33:ASN:ND2	1:A:74:THR:H	1.99	0.60
1:A:22:ALA:HB2	1:A:163:ASP:OD2	2.03	0.58
1:A:226:ASP:HB3	1:A:227:PRO:HD3	1.88	0.55
5:A:1553:BMA:H1	5:A:1554:BMA:H61	1.90	0.54
1:A:135:THR:HA	1:A:474:ASP:OD2	2.06	0.54
1:A:329:LYS:HE2	1:A:329:LYS:N	2.13	0.54
1:A:227:PRO:HA	1:A:284:ASN:HD21	1.75	0.51
1:A:168:THR:HG22	1:A:221:VAL:HG13	1.92	0.50
1:A:422:HIS:CE1	1:A:487:GLU:OE1	2.64	0.50
1:A:396:ALA:HB2	1:A:488:ASP:CG	2.35	0.47
3:A:1503:NAG:O4	3:A:1504:NAG:C1	2.63	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:1503:NAG:O4	3:A:1504:NAG:C2	2.64	0.45
1:A:439:ASN:HD22	1:A:439:ASN:N	2.14	0.45
1:A:67:LEU:HD13	1:A:86:TRP:CZ3	2.53	0.44
1:A:260:ALA:O	1:A:261:ALA:HB3	2.20	0.42
1:A:436:ASN:HD21	1:A:439:ASN:ND2	2.17	0.42
1:A:36:VAL:HG23	1:A:38:PRO:HD3	2.01	0.42
1:A:244:ASP:OD2	1:A:443:ARG:HB2	2.20	0.41
1:A:226:ASP:HB3	1:A:227:PRO:CD	2.50	0.41
1:A:302:ASP:N	6:A:2166:HOH:O	2.53	0.41
1:A:57:ILE:HB	1:A:145:ILE:HG12	2.03	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	493/496 (99%)	478 (97%)	13 (3%)	2 (0%)	34 42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	226	ASP
1	A	79	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	400/401 (100%)	390 (98%)	10 (2%)	47 65

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	141	LEU
1	A	237	SER
1	A	263	ARG
1	A	282	LEU
1	A	284	ASN
1	A	294	THR
1	A	329	LYS
1	A	439	ASN
1	A	468	PHE
1	A	481	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	33	ASN
1	A	44	GLN
1	A	91	GLN
1	A	92	HIS
1	A	104	ASN
1	A	269	ASN
1	A	276	ASN
1	A	284	ASN
1	A	318	ASN
1	A	380	GLN
1	A	422	HIS
1	A	439	ASN
1	A	500	GLN
1	A	517	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates i

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	B	1	1,2	14,14,15	0.62	0	17,19,21	1.14	2 (11%)
2	NAG	B	2	2	14,14,15	0.71	0	17,19,21	0.97	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
2	B	1	NAG	O5-C5-C6	2.22	110.68	107.20
2	B	2	NAG	O5-C5-C6	2.15	110.58	107.20
2	B	1	NAG	O4-C4-C5	-2.02	104.29	109.30

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	2	NAG	C4-C5-C6-O6
2	B	2	NAG	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 4 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	BMA	A	1555	-	12,12,12	0.57	0	17,17,17	0.67	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	A	1504	-	14,14,15	0.58	0	17,19,21	1.75	6 (35%)
5	BMA	A	1553	-	12,12,12	0.51	0	17,17,17	0.78	0
5	BMA	A	1554	-	12,12,12	0.55	0	17,17,17	0.58	0
3	NAG	A	1503	1	14,14,15	0.51	0	17,19,21	0.87	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BMA	A	1555	-	-	0/2/22/22	0/1/1/1
3	NAG	A	1504	-	-	3/6/23/26	0/1/1/1
5	BMA	A	1553	-	-	2/2/22/22	0/1/1/1
5	BMA	A	1554	-	-	2/2/22/22	0/1/1/1
3	NAG	A	1503	1	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1504	NAG	C3-C4-C5	3.62	116.69	110.24
3	A	1504	NAG	C1-O5-C5	2.87	116.08	112.19
3	A	1504	NAG	O5-C1-C2	-2.47	107.39	111.29
3	A	1504	NAG	C8-C7-N2	2.45	120.25	116.10
3	A	1503	NAG	O5-C5-C6	2.40	110.96	107.20
3	A	1504	NAG	C2-N2-C7	2.23	126.07	122.90
3	A	1504	NAG	O5-C5-C6	2.06	110.44	107.20

There are no chirality outliers.

All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	1553	BMA	O5-C5-C6-O6
5	A	1553	BMA	C4-C5-C6-O6
5	A	1554	BMA	C4-C5-C6-O6
3	A	1504	NAG	C8-C7-N2-C2
3	A	1504	NAG	O7-C7-N2-C2
5	A	1554	BMA	O5-C5-C6-O6
3	A	1504	NAG	O5-C5-C6-O6

There are no ring outliers.

4 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1504	NAG	2	0
5	A	1553	BMA	2	0
5	A	1554	BMA	2	0
3	A	1503	NAG	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	496/496 (100%)	-0.19	11 (2%) 62 69	16, 24, 36, 44	1 (0%)

All (11) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	354	GLY	3.4
1	A	352	PHE	3.1
1	A	223	LEU	2.6
1	A	353	ALA	2.5
1	A	182	PRO	2.5
1	A	221	VAL	2.4
1	A	445	VAL	2.3
1	A	356	ASN	2.2
1	A	409	ALA	2.1
1	A	317	THR	2.1
1	A	220	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	NAG	B	2	14/15	0.92	0.14	27,31,35,35	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	NAG	B	1	14/15	0.97	0.09	25,27,30,30	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	BMA	A	1555	12/12	0.42	0.44	111,112,112,112	0
5	BMA	A	1554	12/12	0.49	0.31	101,102,103,103	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	BMA	A	1553	12/12	0.55	0.31	123,123,124,124	0
3	NAG	A	1504	14/15	0.72	0.25	60,65,68,68	0
3	NAG	A	1503	14/15	0.95	0.11	29,31,33,34	0
4	CU	A	1518	1/1	0.99	0.10	11,11,11,11	1
4	CU	A	1519	1/1	0.99	0.04	26,26,26,26	1
4	CU	A	1521	1/1	0.99	0.08	96,96,96,96	0
4	CU	A	1520	1/1	1.00	0.07	36,36,36,36	1

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.