



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 10, 2020 – 12:16 PM GMT

PDB ID : 7A5X
Title : Two copies of the catalytic domain of NanA sialidase from *Streptococcus pneumoniae* juxtaposed in the P212121 space group, in complex with DANA derivatized with a PEG linker on the glycerol group.
Authors : Bridot, C.; Bouckaert, J.
Deposited on : 2020-08-24
Resolution : 1.94 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.14.6
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.14.6


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.94 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	498	
1	B	498	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	R7H	A	801	X	-	-	-
2	R7H	B	801	X	-	-	-

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 15685 atoms, of which 7353 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Sialidase A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	472	7375	2345	3633	660	725	12	0	2	0
1	B	472	7434	2361	3668	668	725	12	0	5	0

There are 48 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	279	MET	-	initiating methionine	UNP P62575
A	280	GLY	-	expression tag	UNP P62575
A	281	HIS	-	expression tag	UNP P62575
A	282	HIS	-	expression tag	UNP P62575
A	283	HIS	-	expression tag	UNP P62575
A	284	HIS	-	expression tag	UNP P62575
A	285	HIS	-	expression tag	UNP P62575
A	286	HIS	-	expression tag	UNP P62575
A	287	HIS	-	expression tag	UNP P62575
A	288	HIS	-	expression tag	UNP P62575
A	289	HIS	-	expression tag	UNP P62575
A	290	HIS	-	expression tag	UNP P62575
A	291	SER	-	expression tag	UNP P62575
A	292	SER	-	expression tag	UNP P62575
A	293	GLY	-	expression tag	UNP P62575
A	294	HIS	-	expression tag	UNP P62575
A	295	ILE	-	expression tag	UNP P62575
A	296	ASP	-	expression tag	UNP P62575
A	297	ASP	-	expression tag	UNP P62575
A	298	ASP	-	expression tag	UNP P62575
A	299	ASP	-	expression tag	UNP P62575
A	300	LYS	-	expression tag	UNP P62575
A	301	HIS	-	expression tag	UNP P62575
A	302	MET	-	expression tag	UNP P62575
B	279	MET	-	initiating methionine	UNP P62575

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	280	GLY	-	expression tag	UNP P62575
B	281	HIS	-	expression tag	UNP P62575
B	282	HIS	-	expression tag	UNP P62575
B	283	HIS	-	expression tag	UNP P62575
B	284	HIS	-	expression tag	UNP P62575
B	285	HIS	-	expression tag	UNP P62575
B	286	HIS	-	expression tag	UNP P62575
B	287	HIS	-	expression tag	UNP P62575
B	288	HIS	-	expression tag	UNP P62575
B	289	HIS	-	expression tag	UNP P62575
B	290	HIS	-	expression tag	UNP P62575
B	291	SER	-	expression tag	UNP P62575
B	292	SER	-	expression tag	UNP P62575
B	293	GLY	-	expression tag	UNP P62575
B	294	HIS	-	expression tag	UNP P62575
B	295	ILE	-	expression tag	UNP P62575
B	296	ASP	-	expression tag	UNP P62575
B	297	ASP	-	expression tag	UNP P62575
B	298	ASP	-	expression tag	UNP P62575
B	299	ASP	-	expression tag	UNP P62575
B	300	LYS	-	expression tag	UNP P62575
B	301	HIS	-	expression tag	UNP P62575
B	302	MET	-	expression tag	UNP P62575

- Molecule 2 is (4 {S},5 {R},6 {R})-5-acetamido-6-[(1 {R},2 {S})-3-[[1-(2-ethoxyethyl)-1,2,3-triazol-4-yl]methylsulfanyl]-1,2-bis(oxidanyl)propyl]-4-oxidanyl-oxane-2-carboxylic acid (three-letter code: R7H) (formula: C₁₈H₃₀N₄O₈S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	A	1	Total	C	H	N	O	S	0	0
			57	18	26	4	8	1		
2	B	1	Total	C	H	N	O	S	0	0
			57	18	26	4	8	1		

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	O S	0	0
			5	4 1		
3	B	1	Total	O S	0	0
			5	4 1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	3	Total Cl	0	0
			3 3		
4	A	1	Total Cl	0	0
			1 1		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Ca	0	0
			1 1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	253	Total	O	0	0
			253	253		
6	B	493	Total	O	0	0
			493	493		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.20Å 96.74Å 226.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.60 – 1.94 56.60 – 1.94	Depositor EDS
% Data completeness (in resolution range)	99.3 (56.60-1.94) 99.3 (56.60-1.94)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.93 (at 1.94Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.189 , 0.238 0.189 , 0.238	Depositor DCC
R_{free} test set	4008 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	35.6	Xtrriage
Anisotropy	0.180	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 46.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	15685	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: R7H, CA, SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.64	0/3829	0.74	1/5169 (0.0%)
1	B	0.82	1/3859 (0.0%)	0.82	0/5208
All	All	0.73	1/7688 (0.0%)	0.78	1/10377 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	586	SER	CB-OG	5.20	1.49	1.42

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	419	ASP	CB-CG-OD1	6.09	123.78	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	761	ALA	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3742	3633	3666	64	0
1	B	3766	3668	3704	34	0
2	A	31	26	0	1	0
2	B	31	26	0	3	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	1	0	0	0	0
4	B	3	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	253	0	0	12	1
6	B	493	0	0	14	2
All	All	8332	7353	7370	102	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 102 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:680:TYR:OH	6:A:901:HOH:O	1.67	1.00
1:A:546[B]:ASP:OD2	6:A:902:HOH:O	1.82	0.98
1:B:528:ASP:OD2	1:B:532:THR:HG22	1.69	0.90
1:B:546[A]:ASP:OD2	6:B:901:HOH:O	1.90	0.89
1:A:711:VAL:HG21	1:A:769:PHE:HZ	1.46	0.81

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:B:909:HOH:O	6:B:1112:HOH:O[3_555]	1.99	0.21
6:A:1130:HOH:O	6:B:1340:HOH:O[4_455]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	472/498 (95%)	447 (95%)	23 (5%)	2 (0%)	34	24
1	B	475/498 (95%)	452 (95%)	22 (5%)	1 (0%)	47	39
All	All	947/996 (95%)	899 (95%)	45 (5%)	3 (0%)	41	32

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	402	ASP
1	A	680	TYR
1	B	680	TYR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	404/426 (95%)	396 (98%)	8 (2%)	55	42
1	B	407/426 (96%)	403 (99%)	4 (1%)	76	71
All	All	811/852 (95%)	799 (98%)	12 (2%)	65	56

5 of 12 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	630	ASN
1	A	723	THR
1	B	627	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	612	GLN
1	B	562	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 6 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	SO4	B	802	-	4,4,4	0.54	0	6,6,6	0.40	0
2	R7H	A	801	-	28,32,32	3.14	7 (25%)	28,43,43	7.61	10 (35%)
2	R7H	B	801	-	28,32,32	2.85	9 (32%)	28,43,43	7.00	11 (39%)
3	SO4	A	802	-	4,4,4	0.18	0	6,6,6	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	R7H	B	801	-	1/1/9/9	6/23/43/43	0/2/2/2
2	R7H	A	801	-	1/1/9/9	8/23/43/43	0/2/2/2

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	801	R7H	C3-C2	-12.38	1.33	1.52
2	B	801	R7H	C3-C2	-8.90	1.38	1.52
2	A	801	R7H	C4-C5	6.83	1.59	1.53
2	B	801	R7H	C4-C5	6.70	1.59	1.53
2	B	801	R7H	CAD-NAZ	5.55	1.41	1.35

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	801	R7H	CAV-CAM-S9	29.60	170.87	113.78
2	B	801	R7H	CAV-CAM-S9	27.73	167.27	113.78
2	A	801	R7H	C9-S9-CAM	23.30	150.71	101.25
2	B	801	R7H	C9-S9-CAM	18.87	141.29	101.25
2	B	801	R7H	O6-C2-C3	7.36	122.78	109.87

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	801	R7H	C2
2	B	801	R7H	C2

5 of 14 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	801	R7H	OAA-CAG-CAJ-NAZ
2	A	801	R7H	CAG-CAJ-NAZ-CAD
2	A	801	R7H	C7-C8-C9-S9
2	A	801	R7H	O8-C8-C9-S9
2	B	801	R7H	C7-C8-C9-S9


There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	801	R7H	1	0
2	B	801	R7H	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	472/498 (94%)	1.19	93 (19%) 1 1	26, 51, 90, 176	0
1	B	472/498 (94%)	0.30	6 (1%) 77 81	20, 33, 52, 84	0
All	All	944/996 (94%)	0.74	99 (10%) 6 9	20, 40, 80, 176	0

The worst 5 of 99 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	306	ALA	11.7
1	A	308	LEU	9.1
1	A	375	TRP	8.8
1	A	766	PHE	8.2
1	A	776	LYS	8.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.


6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	CA	B	806	1/1	0.69	0.13	92,92,92,92	0
3	SO4	B	802	5/5	0.76	0.34	58,66,85,91	0
2	R7H	A	801	31/31	0.78	0.23	44,63,97,99	22
3	SO4	A	802	5/5	0.80	0.30	94,100,104,110	0
5	CA	A	804	1/1	0.83	0.29	79,79,79,79	0
2	R7H	B	801	31/31	0.89	0.19	25,47,118,120	22
4	CL	B	805	1/1	0.93	0.09	62,62,62,62	0
4	CL	B	804	1/1	0.94	0.20	63,63,63,63	0
4	CL	B	803	1/1	0.95	0.14	34,34,34,34	0
4	CL	A	803	1/1	0.99	0.14	37,37,37,37	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.