



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 22, 2020 – 12:36 PM BST

PDB ID : 4ADD  
Title : Structural and functional study of succinyl-ornithine transaminase from E. coli  
Authors : Newman, J.; Peat, T.S.  
Deposited on : 2011-12-23  
Resolution : 2.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	 2% 72% 23%
1	B	406	 73% 21%
1	C	406	 75% 20%
1	D	406	 2% 72% 22%

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 12731 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SUCCINYLORNITHINE TRANSAMINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	400	Total 3074	C 1945	N 547	O 571	S 11	0	5	0
1	B	400	Total 3076	C 1947	N 546	O 572	S 11	0	5	0
1	C	400	Total 3064	C 1940	N 540	O 573	S 11	0	4	0
1	D	400	Total 3073	C 1947	N 544	O 571	S 11	0	6	0

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	B	1	Total 15	C 8	N 1	O 5	P 1	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	D	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is N 2 -(3-CARBOXYPROPANOYL)-L-ORNITHINE (three-letter code: SUO) (formula: C<sub>9</sub>H<sub>16</sub>N<sub>2</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			16	9	2	5		
3	B	1	Total	C	N	O	0	0
			16	9	2	5		
3	C	1	Total	C	N	O	0	0
			16	9	2	5		
3	D	1	Total	C	N	O	0	0
			16	9	2	5		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	118	Total	O	0	0
			118	118		
4	B	79	Total	O	0	0
			79	79		
4	C	73	Total	O	0	0
			73	73		

Continued on next page...



*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	D	50	Total	O	0	0
			50	50		





● Molecule 1: SUCCINYLORNITHINE TRANSAMINASE



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	183.94Å 118.26Å 109.24Å 90.00° 96.82° 90.00°	Depositor
Resolution (Å)	108.46 – 2.45 19.76 – 2.45	Depositor EDS
% Data completeness (in resolution range)	99.8 (108.46-2.45) 100.0 (19.76-2.45)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.21 (at 2.44Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, $R_{free}$	0.170 , 0.225 0.172 , 0.227	Depositor DCC
$R_{free}$ test set	4254 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.2	Xtrriage
Anisotropy	0.012	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 35.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	12731	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.36% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SUO, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.01	2/3153 (0.1%)	1.09	10/4282 (0.2%)
1	B	1.00	4/3158 (0.1%)	1.07	10/4288 (0.2%)
1	C	0.92	3/3145 (0.1%)	0.95	5/4271 (0.1%)
1	D	0.87	2/3162 (0.1%)	0.95	6/4295 (0.1%)
All	All	0.95	11/12618 (0.1%)	1.02	31/17136 (0.2%)

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	34	TRP	CD2-CE2	6.69	1.49	1.41
1	B	13	TRP	CD2-CE2	6.49	1.49	1.41
1	D	72	TRP	CD2-CE2	5.74	1.48	1.41
1	C	72	TRP	CD2-CE2	5.71	1.48	1.41
1	D	34	TRP	CD2-CE2	5.68	1.48	1.41

The worst 5 of 31 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	212	LEU	CA-CB-CG	9.95	138.19	115.30
1	A	32	ARG	NE-CZ-NH1	9.61	125.10	120.30
1	A	299	ILE	CG1-CB-CG2	-7.95	93.92	111.40
1	D	14	MET	CG-SD-CE	-7.79	87.74	100.20
1	A	32	ARG	NE-CZ-NH2	-7.76	116.42	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3074	0	3002	92	0
1	B	3076	0	3009	70	0
1	C	3064	0	2996	76	0
1	D	3073	0	3004	77	0
2	A	15	0	7	0	0
2	B	15	0	6	0	0
2	C	15	0	6	1	0
2	D	15	0	7	0	0
3	A	16	0	13	1	0
3	B	16	0	12	2	0
3	C	16	0	12	2	0
3	D	16	0	12	4	0
4	A	118	0	0	5	0
4	B	79	0	0	3	0
4	C	73	0	0	4	0
4	D	50	0	0	5	0
All	All	12731	0	12086	291	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 291 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:325:ARG:HG3	1:C:325:ARG:HH11	0.98	1.11
1:D:271[B]:ARG:CG	1:D:271[B]:ARG:HH11	1.65	1.10
1:D:271[B]:ARG:NH1	1:D:271[B]:ARG:HG2	1.65	1.05
1:D:353:GLN:H	1:D:353:GLN:HE21	1.05	1.03
1:D:271[B]:ARG:HG2	1:D:271[B]:ARG:HH11	0.82	0.98

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	403/406 (99%)	369 (92%)	32 (8%)	2 (0%)	29	34
1	B	403/406 (99%)	380 (94%)	21 (5%)	2 (0%)	29	34
1	C	402/406 (99%)	368 (92%)	30 (8%)	4 (1%)	15	16
1	D	404/406 (100%)	369 (91%)	32 (8%)	3 (1%)	22	25
All	All	1612/1624 (99%)	1486 (92%)	115 (7%)	11 (1%)	22	25

5 of 11 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	251	ALA
1	D	164	ASP
1	D	251	ALA
1	C	251	ALA
1	C	252	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/312 (100%)	285 (92%)	26 (8%)	11	12
1	B	312/312 (100%)	283 (91%)	29 (9%)	9	9
1	C	311/312 (100%)	287 (92%)	24 (8%)	13	15
1	D	312/312 (100%)	282 (90%)	30 (10%)	8	8
All	All	1246/1248 (100%)	1137 (91%)	109 (9%)	10	11

5 of 109 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	325	ARG
1	C	99	VAL
1	D	287	LEU
1	B	328	LEU
1	C	14	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	3	GLN
1	C	121	HIS
1	D	323	ASN
1	C	103	ASN
1	C	139	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SUO	A	411	2	9,15,15	0.70	0	10,18,18	2.29	3 (30%)
3	SUO	C	411	2	9,15,15	0.40	0	10,18,18	1.47	1 (10%)
3	SUO	B	411	2	9,15,15	1.21	2 (22%)	10,18,18	1.98	3 (30%)
3	SUO	D	411	2	9,15,15	1.30	2 (22%)	10,18,18	1.66	2 (20%)
2	PLP	D	410	3	15,15,16	2.58	5 (33%)	20,22,23	1.46	3 (15%)
2	PLP	B	410	3	15,15,16	2.85	3 (20%)	20,22,23	1.67	5 (25%)
2	PLP	C	410	3	15,15,16	3.15	4 (26%)	20,22,23	1.66	4 (20%)
2	PLP	A	410	3	15,15,16	2.58	5 (33%)	20,22,23	1.49	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SUO	A	411	2	-	1/11/17/17	-
3	SUO	C	411	2	-	1/11/17/17	-
3	SUO	B	411	2	-	2/11/17/17	-
3	SUO	D	411	2	-	2/11/17/17	-
2	PLP	D	410	3	-	3/6/6/8	0/1/1/1
2	PLP	B	410	3	-	2/6/6/8	0/1/1/1
2	PLP	C	410	3	-	0/6/6/8	0/1/1/1
2	PLP	A	410	3	-	1/6/6/8	0/1/1/1

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	410	PLP	C3-C2	8.55	1.49	1.40
2	B	410	PLP	C5-C4	7.45	1.48	1.40
2	D	410	PLP	C3-C2	7.16	1.48	1.40
2	C	410	PLP	C5-C4	7.10	1.48	1.40
2	A	410	PLP	C3-C2	6.86	1.47	1.40

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	411	SUO	CB-CA-N	-5.19	102.64	110.19
2	B	410	PLP	O2P-P-O4P	-4.48	94.82	106.73
2	C	410	PLP	O2P-P-O4P	-4.40	95.02	106.73

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	411	SUO	CW-CX-CY	3.93	119.26	112.67
3	D	411	SUO	CA-N-CV	3.89	129.51	123.33

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	411	SUO	C-CA-CB-CG
3	D	411	SUO	CV-CW-CX-CY
2	D	410	PLP	C5A-O4P-P-O1P
2	D	410	PLP	C5A-O4P-P-O2P
2	B	410	PLP	C5A-O4P-P-O1P

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	411	SUO	1	0
3	C	411	SUO	2	0
3	B	411	SUO	2	0
3	D	411	SUO	4	0
2	C	410	PLP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	400/406 (98%)	-0.57	3 (0%) 86 86	9, 22, 37, 77	15 (3%)
1	B	400/406 (98%)	-0.59	2 (0%) 91 92	12, 22, 37, 55	17 (4%)
1	C	400/406 (98%)	-0.44	2 (0%) 91 92	16, 27, 46, 76	19 (4%)
1	D	400/406 (98%)	-0.31	7 (1%) 68 65	16, 33, 55, 76	19 (4%)
All	All	1600/1624 (98%)	-0.48	14 (0%) 84 85	9, 26, 47, 77	70 (4%)

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	402	SER	4.3
1	A	402	SER	3.7
1	D	4	PRO	3.5
1	D	324[A]	HIS	3.4
1	D	3	GLN	2.9

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SUO	B	411	16/16	0.92	0.13	22,33,39,41	0
3	SUO	D	411	16/16	0.92	0.14	26,46,53,56	0
3	SUO	C	411	16/16	0.93	0.13	33,50,54,60	0
3	SUO	A	411	16/16	0.95	0.11	25,32,41,45	0
2	PLP	D	410	15/16	0.98	0.10	15,17,21,23	0
2	PLP	B	410	15/16	0.99	0.09	11,12,13,13	0
2	PLP	C	410	15/16	0.99	0.07	20,25,27,28	0
2	PLP	A	410	15/16	0.99	0.08	15,17,18,21	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.