



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 23, 2023 – 04:17 AM EDT

PDB ID : 3AFQ
Title : Crystal structure of the single-stranded DNA binding protein from *Mycobacterium leprae* (Form II)
Authors : Kaushal, P.S.; Singh, P.; Sharma, A.; Muniyappa, K.; Vijayan, M.
Deposited on : 2010-03-10
Resolution : 2.80 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	168	
1	B	168	
1	C	168	
1	D	168	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3319 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Single-stranded DNA-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	105	Total 774	C 477	N 140	O 156	S 1	0	0	0
1	B	113	Total 831	C 517	N 147	O 166	S 1	0	0	0
1	C	105	Total 768	C 475	N 141	O 151	S 1	0	0	0
1	D	107	Total 792	C 495	N 140	O 156	S 1	0	0	0


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	48	Total 48	O 48	0	0
2	B	35	Total 35	O 35	0	0
2	C	33	Total 33	O 33	0	0
2	D	38	Total 38	O 38	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Single-stranded DNA-binding protein



- Molecule 1: Single-stranded DNA-binding protein



- Molecule 1: Single-stranded DNA-binding protein



VAL
GLY
ASP
GLU
GLU
PRO
PRO
PHE

• Molecule 1: Single-stranded DNA-binding protein

Chain D: 

MET A2 G3 D4 T5 T6 I7 T8 I9 V10 G11 N12 A15 R20 F21 T22 S23 S24 V28 V29 N30 V33 T36 I39 Y40 D41 ARG GLN SER GLY GLU K47 K48 D49 G50 E51 A52 L53 C57 N58 I59 W60 R61 A64 L71 T72 R73 G74 A75 V79

R82 R83 R84 Q85 R86 S87 F88 E89 THR ARG ARG GLU GLY GLY LYS T97 V98 V99 E102 V103 D104 E105 I106 G107 F108 A113 T114 A115 A120 SER ARG SER GLY GLY GLY GLY GLY PHE GLY SER GLY SER ARG ARG ALA PRO ALA MET SER GLY VAL ASP

ASP PRO TRP GLY SER ALA PRO THR SER GLY SER PHE GLY VAL GLY ASP GLU PRO PRO PHE

4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	102.49Å 102.49Å 120.76Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.80 29.82 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (30.00-2.80) 99.2 (29.82-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.01 (at 2.80Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.211 , 0.235 0.246 , 0.270	Depositor DCC
R_{free} test set	916 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	68.1	Xtrriage
Anisotropy	0.002	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 62.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.000 for -h,-k,l	Xtrriage
Reported twinning fraction	0.610 for H, K, L 0.390 for -h,-k,l	Depositor
Outliers	0 of 18500 reflections	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	3319	wwPDB-VP
Average B, all atoms (Å ²)	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.86 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.2313e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.84	0/780	0.80	0/1061
1	B	0.89	2/839 (0.2%)	0.76	0/1139
1	C	0.76	0/774	0.77	0/1052
1	D	0.87	1/800 (0.1%)	0.72	0/1088
All	All	0.84	3/3193 (0.1%)	0.76	0/4340

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	84	LYS	CE-NZ	5.86	1.63	1.49
1	D	57	CYS	CB-SG	5.74	1.92	1.82
1	B	86	ARG	CZ-NH2	5.68	1.40	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	774	0	757	32	0
1	B	831	0	806	26	0
1	C	768	0	755	39	0
1	D	792	0	775	51	0
2	A	48	0	0	2	0
2	B	35	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	33	0	0	0	0
2	D	38	0	0	1	0
All	All	3319	0	3093	118	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 118 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:PHE:H	1:B:85:GLN:HE22	1.04	0.97
1:D:88:PHE:HB2	1:D:97:THR:N	1.78	0.96
1:C:5:THR:HB	1:D:9:ILE:HG22	1.50	0.93
1:D:39:ILE:HG22	1:D:40:TYR:H	1.34	0.91
1:C:79:VAL:HG22	1:C:106:ILE:HG13	1.56	0.88

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/168 (59%)	89 (90%)	8 (8%)	2 (2%)	7	24
1	B	107/168 (64%)	98 (92%)	7 (6%)	2 (2%)	8	26
1	C	99/168 (59%)	93 (94%)	5 (5%)	1 (1%)	15	44
1	D	101/168 (60%)	92 (91%)	8 (8%)	1 (1%)	15	44
All	All	406/672 (60%)	372 (92%)	28 (7%)	6 (2%)	10	33

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	51	GLU
1	B	3	GLY
1	C	3	GLY
1	D	3	GLY
1	A	119	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	80/128 (62%)	75 (94%)	5 (6%)	18	46
1	B	84/128 (66%)	80 (95%)	4 (5%)	25	58
1	C	78/128 (61%)	67 (86%)	11 (14%)	3	10
1	D	81/128 (63%)	72 (89%)	9 (11%)	6	19
All	All	323/512 (63%)	294 (91%)	29 (9%)	9	28

5 of 29 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	49	ASP
1	D	88	PHE
1	C	96	ARG
1	D	73	ARG
1	C	73	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	66	ASN
1	D	30	ASN
1	D	66	ASN
1	B	66	ASN
1	A	66	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	105/168 (62%)	0.37	5 (4%) 30 21	43, 71, 92, 104	0
1	B	113/168 (67%)	0.49	9 (7%) 12 6	43, 72, 100, 104	0
1	C	105/168 (62%)	0.37	7 (6%) 17 10	43, 73, 95, 111	0
1	D	107/168 (63%)	0.46	9 (8%) 11 5	43, 72, 97, 120	0
All	All	430/672 (63%)	0.43	30 (6%) 16 9	43, 72, 98, 120	0

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	47	TRP	7.5
1	B	47	TRP	6.4
1	D	40	TYR	5.7
1	C	49	ASP	4.9
1	B	46	GLU	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.