



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 24, 2023 – 07:12 AM EDT

PDB ID : 3AK5  
Title : Hemoglobin protease (Hbp) passenger missing domain-2  
Authors : Nishimura, K.; Park, S.-Y.; Tame, J.R.H.  
Deposited on : 2010-07-07  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	975	 0.2% 78% 19% ..
1	B	975	 0.2% 78% 19% ..
1	C	975	 2% 77% 18% ..
1	D	975	 0.2% 76% 18% ..

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 30307 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemoglobin-binding protease hbp.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	970	7278	4511	1268	1485	14	7	0	0
1	B	970	7278	4511	1268	1485	14	4	0	0
1	C	952	7149	4435	1241	1459	14	4	1	0
1	D	948	7113	4411	1236	1452	14	9	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	607	GLY	-	SEE REMARK 999	UNP O88093
B	607	GLY	-	SEE REMARK 999	UNP O88093
C	607	GLY	-	SEE REMARK 999	UNP O88093
D	607	GLY	-	SEE REMARK 999	UNP O88093

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total 1	Ca 1	0	0
2	B	1	Total 1	Ca 1	0	0
2	C	1	Total 1	Ca 1	0	0
2	D	1	Total 1	Ca 1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	426	Total 426	O 426	0	0
3	B	430	Total 430	O 430	0	0
3	C	321	Total 321	O 321	0	0
3	D	308	Total 308	O 308	0	0


### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Hemoglobin-binding protease hbp





- Molecule 1: Hemoglobin-binding protease hbp







• Molecule 1: Hemoglobin-binding protease hbp



• Molecule 1: Hemoglobin-binding protease hbp





## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	292.80Å 53.94Å 335.95Å 90.00° 107.32° 90.00°	Depositor
Resolution (Å)	48.56 – 2.20 48.56 – 2.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (48.56-2.20) 93.1 (48.56-2.20)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.33 (at 2.20Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, $R_{free}$	0.226 , 0.280 0.224 , 0.279	Depositor DCC
$R_{free}$ test set	11985 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.4	Xtrriage
Anisotropy	0.084	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 44.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	30307	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 94.28 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 3.1752e-09. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.07	3/7407 (0.0%)	0.99	17/10069 (0.2%)
1	B	1.15	6/7407 (0.1%)	1.04	20/10069 (0.2%)
1	C	1.09	5/7278 (0.1%)	0.99	9/9895 (0.1%)
1	D	1.26	7/7236 (0.1%)	1.12	11/9835 (0.1%)
All	All	1.14	21/29328 (0.1%)	1.04	57/39868 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	2
1	C	0	2
1	D	0	2
All	All	0	7

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	89	GLU	CD-OE2	-51.39	0.69	1.25
1	B	851	GLU	CG-CD	-31.69	1.04	1.51
1	B	955	LYS	CE-NZ	29.51	2.22	1.49
1	D	89	GLU	CD-OE1	25.27	1.53	1.25
1	A	619	LYS	CE-NZ	-23.79	0.89	1.49

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	523	ASP	CB-CG-OD2	-43.35	79.29	118.30
1	D	89	GLU	OE1-CD-OE2	31.72	161.37	123.30
1	C	523	ASP	CB-CG-OD1	-26.34	94.59	118.30
1	D	89	GLU	CG-CD-OE1	-20.08	78.14	118.30
1	B	851	GLU	CG-CD-OE1	19.40	157.10	118.30

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	968	SER	Peptide
1	B	1069	LEU	Peptide
1	B	963	PHE	Peptide
1	C	523	ASP	Sidechain
1	C	968	SER	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	7278	0	6993	115	0
1	B	7278	0	6993	109	0
1	C	7149	0	6858	141	0
1	D	7113	0	6823	136	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	426	0	0	15	0
3	B	430	0	0	15	0
3	C	321	0	0	9	0
3	D	308	0	0	7	0
All	All	30307	0	27667	497	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 497 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:364[B]:ASN:OD1	1:D:364:ASN:ND2	1.76	1.16
1:B:632:VAL:HG12	1:B:634:GLY:H	1.27	0.99
1:A:401:ILE:HD12	1:A:421:LEU:HD11	1.40	0.99
1:C:842:THR:HG22	3:C:1285:HOH:O	1.62	0.97
1:D:172:THR:HG22	1:D:174:GLN:H	1.31	0.94

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	966/975 (99%)	898 (93%)	55 (6%)	13 (1%)	12	9
1	B	966/975 (99%)	900 (93%)	61 (6%)	5 (0%)	29	31
1	C	945/975 (97%)	876 (93%)	63 (7%)	6 (1%)	25	26
1	D	938/975 (96%)	870 (93%)	60 (6%)	8 (1%)	17	16
All	All	3815/3900 (98%)	3544 (93%)	239 (6%)	32 (1%)	19	19

5 of 32 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	338	GLY
1	A	1022	ASP
1	C	85	ASN
1	D	85	ASN
1	A	85	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	785/787 (100%)	735 (94%)	50 (6%)	17	20
1	B	785/787 (100%)	721 (92%)	64 (8%)	11	11
1	C	772/787 (98%)	724 (94%)	48 (6%)	18	21
1	D	767/787 (98%)	719 (94%)	48 (6%)	18	20
All	All	3109/3148 (99%)	2899 (93%)	210 (7%)	16	17

5 of 210 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	75	ARG
1	C	796	THR
1	D	912	GLN
1	C	123	VAL
1	C	312	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 51 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	467	ASN
1	C	946	ASN
1	D	957	ASN
1	C	621	ASN
1	C	808	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	970/975 (99%)	-0.25	7 (0%) 87 86	13, 31, 53, 66	4 (0%)
1	B	970/975 (99%)	-0.24	5 (0%) 91 90	13, 30, 53, 70	2 (0%)
1	C	952/975 (97%)	-0.23	16 (1%) 70 68	14, 32, 54, 69	2 (0%)
1	D	948/975 (97%)	-0.23	14 (1%) 73 72	15, 32, 54, 65	4 (0%)
All	All	3840/3900 (98%)	-0.24	42 (1%) 80 79	13, 31, 54, 70	12 (0%)

The worst 5 of 42 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	991	LEU	6.0
1	D	1049	PHE	5.3
1	B	633	THR	5.0
1	C	968	SER	4.9
1	D	909	LEU	4.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	CA	A	1200	1/1	0.99	0.14	22,22,22,22	0
2	CA	B	1200	1/1	0.99	0.10	19,19,19,19	0
2	CA	D	1200	1/1	0.99	0.10	22,22,22,22	0
2	CA	C	1200	1/1	1.00	0.10	19,19,19,19	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.