



wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 26, 2024 – 06:07 AM EDT

PDB ID : 7AM7
Title : Crystal structure of Peptilgase mutant - M222P/L217H/A225N/F189W/N218D
Authors : Rozeboom, H.J.; Janssen, D.J.
Deposited on : 2020-10-08
Resolution : 2.61 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3797 (2.64-2.60)
Clashscore	141614	4168 (2.64-2.60)
Ramachandran outliers	138981	4093 (2.64-2.60)
Sidechain outliers	138945	4093 (2.64-2.60)
RSRZ outliers	127900	3731 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	 3% 84% 13% ..
1	C	272	 2% 89% 8% .
2	B	272	 % 91% 8% .
3	P	13	 8% 31% 8% 8% 54%

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 5975 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Subtilisin BPN'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	267	1888	1173	328	381	6	0	0	0
1	C	265	1915	1193	330	386	6	0	3	0

There are 68 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	LYS	GLN	engineered mutation	UNP P00782
A	3	CYS	SER	engineered mutation	UNP P00782
A	5	SER	PRO	engineered mutation	UNP P00782
A	43	ASN	LYS	engineered mutation	UNP P00782
A	50	PHE	MET	engineered mutation	UNP P00782
A	?	-	ALA	deletion	UNP P00782
A	?	-	ALA	deletion	UNP P00782
A	?	-	LEU	deletion	UNP P00782
A	?	-	ASN	deletion	UNP P00782
A	?	-	ASN	deletion	UNP P00782
A	?	-	SER	deletion	UNP P00782
A	?	-	ILE	deletion	UNP P00782
A	?	-	GLY	deletion	UNP P00782
A	?	-	VAL	deletion	UNP P00782
A	74	ALA	GLY	engineered mutation	UNP P00782
A	156	SER	GLU	engineered mutation	UNP P00782
A	166	SER	GLY	engineered mutation	UNP P00782
A	169	ALA	GLY	engineered mutation	UNP P00782
A	188	PRO	SER	engineered mutation	UNP P00782
A	189	TRP	PHE	engineered mutation	UNP P00782
A	206	CYS	GLN	engineered mutation	UNP P00782
A	217	HIS	TYR	engineered mutation	UNP P00782
A	218	ASP	ASN	engineered mutation	UNP P00782
A	221	CSO	SER	engineered mutation	UNP P00782
A	222	PRO	MET	engineered mutation	UNP P00782

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	225	ASN	PRO	engineered mutation	UNP P00782
A	254	ALA	THR	engineered mutation	UNP P00782
A	271	GLU	GLN	engineered mutation	UNP P00782
A	276	HIS	-	expression tag	UNP P00782
A	277	HIS	-	expression tag	UNP P00782
A	278	HIS	-	expression tag	UNP P00782
A	279	HIS	-	expression tag	UNP P00782
A	280	HIS	-	expression tag	UNP P00782
A	281	HIS	-	expression tag	UNP P00782
C	2	LYS	GLN	engineered mutation	UNP P00782
C	3	CYS	SER	engineered mutation	UNP P00782
C	5	SER	PRO	engineered mutation	UNP P00782
C	43	ASN	LYS	engineered mutation	UNP P00782
C	50	PHE	MET	engineered mutation	UNP P00782
C	?	-	ALA	deletion	UNP P00782
C	?	-	ALA	deletion	UNP P00782
C	?	-	LEU	deletion	UNP P00782
C	?	-	ASN	deletion	UNP P00782
C	?	-	ASN	deletion	UNP P00782
C	?	-	SER	deletion	UNP P00782
C	?	-	ILE	deletion	UNP P00782
C	?	-	GLY	deletion	UNP P00782
C	?	-	VAL	deletion	UNP P00782
C	74	ALA	GLY	engineered mutation	UNP P00782
C	156	SER	GLU	engineered mutation	UNP P00782
C	166	SER	GLY	engineered mutation	UNP P00782
C	169	ALA	GLY	engineered mutation	UNP P00782
C	188	PRO	SER	engineered mutation	UNP P00782
C	189	TRP	PHE	engineered mutation	UNP P00782
C	206	CYS	GLN	engineered mutation	UNP P00782
C	217	HIS	TYR	engineered mutation	UNP P00782
C	218	ASP	ASN	engineered mutation	UNP P00782
C	221	CSO	SER	engineered mutation	UNP P00782
C	222	PRO	MET	engineered mutation	UNP P00782
C	225	ASN	PRO	engineered mutation	UNP P00782
C	254	ALA	THR	engineered mutation	UNP P00782
C	271	GLU	GLN	engineered mutation	UNP P00782
C	276	HIS	-	expression tag	UNP P00782
C	277	HIS	-	expression tag	UNP P00782
C	278	HIS	-	expression tag	UNP P00782
C	279	HIS	-	expression tag	UNP P00782
C	280	HIS	-	expression tag	UNP P00782

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	281	HIS	-	expression tag	UNP P00782

- Molecule 2 is a protein called Subtilisin BPN'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	268	1897	1179	331	381	6	0	0	0

There are 34 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	2	LYS	GLN	engineered mutation	UNP P00782
B	3	CYS	SER	engineered mutation	UNP P00782
B	5	SER	PRO	engineered mutation	UNP P00782
B	43	ASN	LYS	engineered mutation	UNP P00782
B	50	PHE	MET	engineered mutation	UNP P00782
B	?	-	ALA	deletion	UNP P00782
B	?	-	ALA	deletion	UNP P00782
B	?	-	LEU	deletion	UNP P00782
B	?	-	ASN	deletion	UNP P00782
B	?	-	ASN	deletion	UNP P00782
B	?	-	SER	deletion	UNP P00782
B	?	-	ILE	deletion	UNP P00782
B	?	-	GLY	deletion	UNP P00782
B	?	-	VAL	deletion	UNP P00782
B	74	ALA	GLY	engineered mutation	UNP P00782
B	156	SER	GLU	engineered mutation	UNP P00782
B	166	SER	GLY	engineered mutation	UNP P00782
B	169	ALA	GLY	engineered mutation	UNP P00782
B	188	PRO	SER	engineered mutation	UNP P00782
B	189	TRP	PHE	engineered mutation	UNP P00782
B	206	CYS	GLN	engineered mutation	UNP P00782
B	217	HIS	TYR	engineered mutation	UNP P00782
B	218	ASP	ASN	engineered mutation	UNP P00782
B	221	CYS	SER	engineered mutation	UNP P00782
B	222	PRO	MET	engineered mutation	UNP P00782
B	225	ASN	PRO	engineered mutation	UNP P00782
B	254	ALA	THR	engineered mutation	UNP P00782
B	271	GLU	GLN	engineered mutation	UNP P00782
B	276	HIS	-	expression tag	UNP P00782
B	277	HIS	-	expression tag	UNP P00782
B	278	HIS	-	expression tag	UNP P00782

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	279	HIS	-	expression tag	UNP P00782
B	280	HIS	-	expression tag	UNP P00782
B	281	HIS	-	expression tag	UNP P00782

- Molecule 3 is a protein called Eglin C fragment.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
3	P	6	Total	C	N	O	0	0	0
			43	27	6	10			

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		
4	B	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	C	O	0	0
			6	3	3		
4	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	B	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		
5	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	C	1	Total	C	O	0	0
			10	6	4		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	43	Total	O	0	0
			43	43		
7	B	49	Total	O	0	0
			49	49		
7	C	51	Total	O	0	0
			51	51		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Subtilisin BPN'




- Molecule 1: Subtilisin BPN'



- Molecule 2: Subtilisin BPN'



- Molecule 3: Eglin C fragment



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	105.80Å 105.80Å 191.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.59 – 2.61 48.56 – 2.61	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.59-2.61) 99.8 (48.56-2.61)	Depositor EDS
R_{merge}	0.26	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.196 , 0.237 0.203 , 0.241	Depositor DCC
R_{free} test set	1661 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	34.2	Xtrriage
Anisotropy	0.424	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 27.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5975	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CSO, SO4, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/1921	0.82	0/2625
1	C	0.74	0/1946	0.85	0/2654
2	B	0.73	0/1939	0.85	0/2652
3	P	1.00	0/43	1.66	1/59 (1.7%)
All	All	0.73	0/5849	0.85	1/7990 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
3	P	78	ASP	CA-C-O	6.32	133.38	120.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	213[B]	LYS	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1888	0	1833	21	0
1	C	1915	0	1856	26	0
2	B	1897	0	1840	15	0
3	P	43	0	42	5	0
4	A	12	0	16	1	0
4	B	30	0	40	1	0
4	C	12	0	16	0	0
5	A	5	0	0	0	0
5	B	10	0	0	0	0
5	C	10	0	0	0	0
6	C	10	0	14	0	0
7	A	43	0	0	0	0
7	B	49	0	0	1	0
7	C	51	0	0	1	0
All	All	5975	0	5657	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 53 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:221:CYS:SG	1:C:213[B]:LYS:HB2	2.24	0.77
1:C:155:ASN:HD22	3:P:78:ASP:HB2	1.50	0.74
2:B:221:CYS:SG	1:C:213[B]:LYS:CB	2.75	0.73
1:C:196:LEU:HD11	1:C:199:MET:HE1	1.71	0.72
1:A:125:SER:HB3	1:A:221:CSO:HB2	1.75	0.68

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/272 (97%)	257 (97%)	5 (2%)	2 (1%)	19	36
1	C	262/272 (96%)	255 (97%)	6 (2%)	1 (0%)	34	55
2	B	266/272 (98%)	258 (97%)	7 (3%)	1 (0%)	34	55
3	P	4/13 (31%)	4 (100%)	0	0	100	100
All	All	796/829 (96%)	774 (97%)	18 (2%)	4 (0%)	29	50

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	276	HIS
1	A	2	LYS
1	C	63	SER
1	A	36	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/204 (98%)	190 (96%)	9 (4%)	27	50
1	C	202/204 (99%)	197 (98%)	5 (2%)	47	71
2	B	201/205 (98%)	197 (98%)	4 (2%)	55	77
3	P	6/12 (50%)	4 (67%)	2 (33%)	0	0
All	All	608/625 (97%)	588 (97%)	20 (3%)	40	62

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	105	SER
1	C	217	HIS
3	P	78	ASP
3	P	73	SER
1	A	183	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	61	ASN
1	C	155	ASN
1	C	275	GLN
1	A	225	ASN
1	A	252	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSO	A	221	1	3,6,7	0.87	0	0,6,8	-	-
1	CSO	C	221	1	3,6,7	0.81	0	0,6,8	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSO	A	221	1	-	1/1/5/7	-
1	CSO	C	221	1	-	1/1/5/7	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	221	CSO	N-CA-CB-SG
1	C	221	CSO	N-CA-CB-SG

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	221	CSO	2	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	C	302	-	5,5,5	0.07	0	5,5,5	0.22	0
4	GOL	A	301	-	5,5,5	0.11	0	5,5,5	0.32	0
4	GOL	A	302	-	5,5,5	0.15	0	5,5,5	0.33	0
4	GOL	C	301	-	5,5,5	0.11	0	5,5,5	0.32	0
5	SO4	B	306	-	4,4,4	0.35	0	6,6,6	0.10	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GOL	B	304	-	5,5,5	0.12	0	5,5,5	0.32	0
4	GOL	B	305	-	5,5,5	0.21	0	5,5,5	0.38	0
5	SO4	C	304	-	4,4,4	0.35	0	6,6,6	0.06	0
6	PGE	C	303	-	9,9,9	0.24	0	8,8,8	0.11	0
5	SO4	B	307	-	4,4,4	0.36	0	6,6,6	0.05	0
4	GOL	B	301	-	5,5,5	0.19	0	5,5,5	0.39	0
5	SO4	A	303	-	4,4,4	0.34	0	6,6,6	0.07	0
4	GOL	B	303	-	5,5,5	0.20	0	5,5,5	0.60	0
5	SO4	C	305	-	4,4,4	0.36	0	6,6,6	0.11	0
4	GOL	B	302	-	5,5,5	0.10	0	5,5,5	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	C	302	-	-	0/4/4/4	-
4	GOL	A	301	-	-	2/4/4/4	-
4	GOL	A	302	-	-	2/4/4/4	-
4	GOL	C	301	-	-	2/4/4/4	-
4	GOL	B	304	-	-	0/4/4/4	-
4	GOL	B	305	-	-	2/4/4/4	-
6	PGE	C	303	-	-	3/7/7/7	-
4	GOL	B	301	-	-	0/4/4/4	-
4	GOL	B	303	-	-	1/4/4/4	-
4	GOL	B	302	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	302	GOL	O1-C1-C2-O2
4	A	302	GOL	O1-C1-C2-C3
4	B	305	GOL	C1-C2-C3-O3
4	A	301	GOL	O1-C1-C2-O2
6	C	303	PGE	O2-C3-C4-O3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	302	GOL	1	0
4	B	304	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	266/272 (97%)	-0.17	8 (3%) 50 44	27, 38, 57, 96	0
1	C	264/272 (97%)	-0.09	5 (1%) 66 62	30, 41, 59, 102	0
2	B	268/272 (98%)	-0.24	4 (1%) 73 70	26, 34, 51, 91	0
3	P	6/13 (46%)	1.26	1 (16%) 1 1	56, 64, 77, 81	0
All	All	804/829 (96%)	-0.16	18 (2%) 62 57	26, 37, 58, 102	0

The worst 5 of 18 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	214[A]	TYR	6.5
1	C	216	ALA	6.5
2	B	276	HIS	5.7
2	B	277	HIS	5.4
1	A	276	HIS	4.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CSO	A	221	7/8	0.92	0.19	38,41,48,61	0
1	CSO	C	221	7/8	0.94	0.13	41,45,59,64	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	SO4	C	304	5/5	0.82	0.27	95,103,107,112	0
4	GOL	B	305	6/6	0.83	0.22	50,54,57,58	0
4	GOL	C	301	6/6	0.84	0.33	57,62,65,66	0
5	SO4	B	306	5/5	0.84	0.42	95,107,110,114	0
4	GOL	A	301	6/6	0.84	0.16	49,60,61,63	0
5	SO4	C	305	5/5	0.85	0.29	95,96,97,100	0
6	PGE	C	303	10/10	0.85	0.21	52,62,64,65	0
4	GOL	B	301	6/6	0.87	0.17	42,49,51,54	0
5	SO4	B	307	5/5	0.88	0.37	95,95,100,101	5
4	GOL	B	303	6/6	0.88	0.25	48,51,53,53	0
5	SO4	A	303	5/5	0.89	0.18	83,85,91,92	0
4	GOL	B	302	6/6	0.89	0.26	43,51,54,60	0
4	GOL	B	304	6/6	0.90	0.23	60,64,67,70	0
4	GOL	C	302	6/6	0.90	0.22	55,55,58,59	0
4	GOL	A	302	6/6	0.92	0.21	49,50,54,55	0

6.5 Other polymers [i](#)

There are no such residues in this entry.