

wwPDB X-ray Structure Validation Summary Report (i)

Jan 2, 2024 – 09:13 pm GMT

PDB ID : 5APU

Title : Sequence IANKEDKAD inserted between GCN4 adaptors - Structure A9b

black

Authors: Hartmann, M.D.; Mendler, C.T.; Lupas, A.N.; Hernandez Alvarez, B.

Deposited on : 2015-09-17

Resolution : 1.35 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.36

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 1.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	Similar resolution $(\# \text{Entries, resolution range}(\text{\AA}))$
R_{free}	130704	1509 (1.38-1.34)
Clashscore	141614	1551 (1.38-1.34)
Ramachandran outliers	138981	1530 (1.38-1.34)
Sidechain outliers	138945	1530 (1.38-1.34)
RSRZ outliers	127900	1487 (1.38-1.34)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	A	95	67%		28%		
1	В	95	67%		28%		
1	С	95	66%	6%	27%		



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 1932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called GENERAL CONTROL PROTEIN GCN4.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Λ	68	Total	С	N	О	S	0	4	0
1	A	00	570	365	89	114	2	0	4	
1	D	68	Total	С	N	О	S	0	9	0
1	Ъ	00	563	359	93	109	2	0	2	
1	С	69	Total	С	N	О	S	0	2	0
		09	579	369	99	109	2	U	3	U

There are 111 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-25	MET	-	expression tag	UNP P03069
A	-24	LYS	-	expression tag	UNP P03069
A	-23	HIS	-	expression tag	UNP P03069
A	-22	HIS	-	expression tag	UNP P03069
A	-21	HIS	-	expression tag	UNP P03069
A	-20	HIS	-	expression tag	UNP P03069
A	-19	HIS	-	expression tag	UNP P03069
A	-18	HIS	-	expression tag	UNP P03069
A	-17	PRO	-	expression tag	UNP P03069
A	-16	MET	-	expression tag	UNP P03069
A	-15	SER	-	expression tag	UNP P03069
A	-14	ASP	-	expression tag	UNP P03069
A	-13	TYR	-	expression tag	UNP P03069
A	-12	ASP	-	expression tag	UNP P03069
A	-11	ILE	-	expression tag	UNP P03069
A	-10	PRO	-	expression tag	UNP P03069
A	-9	THR	-	expression tag	UNP P03069
A	-8	THR	-	expression tag	UNP P03069
A	-7	GLU		expression tag	UNP P03069
A	-6	ASN	-	expression tag	UNP P03069
A	-5	LEU	-	expression tag	UNP P03069
A	-4	TYR	-	expression tag	UNP P03069
A	-3	PHE	-	expression tag	UNP P03069



Continued from previous page...

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
A	-2	GLN	-	expression tag	UNP P03069
A	-1	GLY	-	expression tag	UNP P03069
A	0	HIS	-	expression tag	UNP P03069
A	15	VAL	ASN	engineered mutation	UNP P03069
A	29	ILE	-	linker	UNP P03069
A	30	ALA	_	linker	UNP P03069
A	31	ASN	-	linker	UNP P03069
A	32	LYS	-	linker	UNP P03069
A	33	GLU	-	linker	UNP P03069
A	34	ASP	-	linker	UNP P03069
A	35	LYS	-	linker	UNP P03069
A	36	ALA	-	linker	UNP P03069
A	37	ASP	-	linker	UNP P03069
A	52	VAL	ASN	engineered mutation	UNP P03069
В	-25	MET	-	expression tag	UNP P03069
В	-24	LYS	-	expression tag	UNP P03069
В	-23	HIS	-	expression tag	UNP P03069
В	-22	HIS	-	expression tag	UNP P03069
В	-21	HIS	-	expression tag	UNP P03069
В	-20	HIS	-	expression tag	UNP P03069
В	-19	HIS	-	expression tag	UNP P03069
В	-18	HIS	-	expression tag	UNP P03069
В	-17	PRO	-	expression tag	UNP P03069
В	-16	MET	-	expression tag	UNP P03069
В	-15	SER	-	expression tag	UNP P03069
В	-14	ASP	-	expression tag	UNP P03069
В	-13	TYR	-	expression tag	UNP P03069
В	-12	ASP	_	expression tag	UNP P03069
В	-11	ILE	-	expression tag	UNP P03069
В	-10	PRO	-	expression tag	UNP P03069
В	-9	THR	_	expression tag	UNP P03069
В	-8	THR	-	expression tag	UNP P03069
В	-7	GLU	-	expression tag	UNP P03069
В	-6	ASN	-	expression tag	UNP P03069
В	-5	LEU	-	expression tag	UNP P03069
В	-4	TYR	-	expression tag	UNP P03069
В	-3	PHE	-	expression tag	UNP P03069
В	-2	GLN	-	expression tag	UNP P03069
В	-1	GLY	-	expression tag	UNP P03069
В	0	HIS	-	expression tag	UNP P03069
В	15	VAL	ASN	engineered mutation	UNP P03069
В	29	ILE	-	linker	UNP P03069



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled	Actual	Comment	Reference
В	30	ALA	-	linker	UNP P03069
В	31	ASN	-	linker	UNP P03069
В	32	LYS	-	linker	UNP P03069
В	33	GLU	_	linker	UNP P03069
В	34	ASP	-	linker	UNP P03069
В	35	LYS	-	linker	UNP P03069
В	36	ALA	-	linker	UNP P03069
В	37	ASP	-	linker	UNP P03069
В	52	VAL	ASN	engineered mutation	UNP P03069
С	-25	MET	-	expression tag	UNP P03069
С	-24	LYS	-	expression tag	UNP P03069
С	-23	HIS	-	expression tag	UNP P03069
С	-22	HIS	-	expression tag	UNP P03069
С	-21	HIS	-	expression tag	UNP P03069
С	-20	HIS	-	expression tag	UNP P03069
С	-19	HIS	-	expression tag	UNP P03069
С	-18	HIS	-	expression tag	UNP P03069
С	-17	PRO	-	expression tag	UNP P03069
С	-16	MET	_	expression tag	UNP P03069
С	-15	SER	-	expression tag	UNP P03069
С	-14	ASP	-	expression tag	UNP P03069
С	-13	TYR	-	expression tag	UNP P03069
С	-12	ASP	-	expression tag	UNP P03069
С	-11	ILE	-	expression tag	UNP P03069
С	-10	PRO	-	expression tag	UNP P03069
С	-9	THR	-	expression tag	UNP P03069
С	-8	THR	-	expression tag	UNP P03069
С	-7	GLU	-	expression tag	UNP P03069
С	-6	ASN	-	expression tag	UNP P03069
С	-5	LEU	-	expression tag	UNP P03069
С	-4	TYR	-	expression tag	UNP P03069
С	-3	PHE	-	expression tag	UNP P03069
С	-2	GLN		expression tag	UNP P03069
С	-1	GLY	-	expression tag	UNP P03069
С	0	HIS		expression tag	UNP P03069
С	15	VAL	ASN	engineered mutation	UNP P03069
С	29	ILE	-	linker	UNP P03069
С	30	ALA	-	linker	UNP P03069
С	31	ASN	-	linker	UNP P03069
С	32	LYS	-	linker	UNP P03069
С	33	GLU	-	linker	UNP P03069
С	34	ASP	-	linker	UNP P03069



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
С	35	LYS	-	linker	UNP P03069
С	36	ALA	-	linker	UNP P03069
С	37	ASP	-	linker	UNP P03069
С	52	VAL	ASN	engineered mutation	UNP P03069

 \bullet Molecule 2 is UREA (three-letter code: URE) (formula: $\mathrm{CH_4N_2O}).$



\mathbf{Mol}	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total 4	C 1	N 2	O 1	0	0

• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	92	Total O 92 92	0	0
3	В	72	Total O 72 72	0	0
3	С	52	Total O 52 52	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

● Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain A: 67% ・ 28%

● Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain B: 67% ・ 28%

● Molecule 1: GENERAL CONTROL PROTEIN GCN4

Chain C: 66% 6% 27%



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	26.20Å 37.52Å 95.00Å	Domositon
a, b, c, α , β , γ	90.00° 92.56° 90.00°	Depositor
Resolution (Å)	34.89 - 1.35	Depositor
Resolution (A)	34.89 - 1.35	EDS
% Data completeness	95.9 (34.89-1.35)	Depositor
(in resolution range)	95.9 (34.89-1.35)	EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.21 (at 1.35Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
D D	0.163 , 0.199	Depositor
R, R_{free}	0.177 , 0.210	DCC
R_{free} test set	1963 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	16.3	Xtriage
Anisotropy	0.083	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.40 , 47.6	EDS
L-test for twinning ²	$< L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	0.035 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1932	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 16.92% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: URE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.28	0/588	0.46	0/786	
1	В	0.28	0/573	0.43	0/763	
1	С	0.30	0/592	0.44	0/788	
All	All	0.28	0/1753	0.44	0/2337	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	570	0	591	5	0
1	В	563	0	594	4	0
1	С	579	0	606	5	0
2	A	4	0	4	0	0
3	A	92	0	0	0	0
3	В	72	0	0	0	0
3	С	52	0	0	1	0
All	All	1932	0	1795	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.



The worst 5 of 8 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-1 Atom-2		$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:A:12:LEU:HD23	1:C:11:LEU:HD21	1.65	0.79
1:A:21:GLU:HG3	1:B:22:VAL:HG11	1.86	0.58
1:C:60:ALA:O	1:C:64:LYS:HG2	2.07	0.54
1:B:29:ILE:HD13	1:C:29:ILE:HG12	1.91	0.52
1:A:32:LYS:HG3	1:B:29:ILE:HD11	1.94	0.50

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	71/95 (75%)	71 (100%)	0	0	100	100
1	В	68/95 (72%)	68 (100%)	0	0	100	100
1	С	70/95 (74%)	70 (100%)	0	0	100	100
All	All	209/285 (73%)	209 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mo	Chain	Analysed Rotameric O		Outliers	Percentiles	
1	A	63/89 (71%)	63 (100%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed Rotameric Outliers		Percentiles		
1	В	63/89 (71%)	62 (98%)	1 (2%)	62 30	
1	С	62/89 (70%)	61 (98%)	1 (2%)	62 30	
All	All	188/267 (70%)	186 (99%)	2 (1%)	73 45	

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	34	ASP
1	С	10	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type	
1	В	40	GLN	

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Mol Type Chain		e Chain Res Link		Bond lengths			Bond angles		
MIOI	Type	Chain	n nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	URE	A	1068	-	3,3,3	0.69	0	3,3,3	0.35	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ} {>} 2$		$OWAB(A^2)$	Q<0.9
1	A	68/95 (71%)	0.43	4 (5%)	22 25	12, 22, 39, 48	0
1	В	68/95 (71%)	0.34	4 (5%)	22 25	11, 22, 37, 43	0
1	С	69/95 (72%)	0.39	6 (8%)	10 12	11, 21, 37, 47	0
All	All	205/285 (71%)	0.39	14 (6%)	17 19	11, 22, 38, 48	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	67	GLY	6.0
1	В	0	HIS	4.3
1	С	4	LEU	4.2
1	A	1	MET	3.9
1	В	67	GLY	3.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	URE	A	1068	4/4	0.97	0.13	20,22,22,24	0

6.5 Other polymers (i)

There are no such residues in this entry.

