



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 26, 2023 – 05:09 AM EDT

PDB ID : 6ATF
Title : HLA-DRB1*1402 in complex with Vimentin59-71
Authors : Scally, S.W.; Ting, Y.T.; Rossjohn, J.
Deposited on : 2017-08-29
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	189	 2% 93% 5%
1	D	189	 2% 92% 5%
2	B	200	 84% 10% 6%
2	E	200	 87% 7% 6%
3	C	13	 85% 15%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	F	13	 85% 15%
4	G	2	 50% 50%
4	H	2	 50% 50%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 7456 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA class II histocompatibility antigen, DR alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	180	1497	968	242	282	5	0	3	0
1	D	179	1482	960	239	278	5	0	2	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	182	THR	-	expression tag	UNP P01903
A	183	SER	-	expression tag	UNP P01903
A	184	GLY	-	expression tag	UNP P01903
A	185	ASP	-	expression tag	UNP P01903
A	186	ASP	-	expression tag	UNP P01903
A	187	ASP	-	expression tag	UNP P01903
A	188	ASP	-	expression tag	UNP P01903
A	189	LYS	-	expression tag	UNP P01903
D	182	THR	-	expression tag	UNP P01903
D	183	SER	-	expression tag	UNP P01903
D	184	GLY	-	expression tag	UNP P01903
D	185	ASP	-	expression tag	UNP P01903
D	186	ASP	-	expression tag	UNP P01903
D	187	ASP	-	expression tag	UNP P01903
D	188	ASP	-	expression tag	UNP P01903
D	189	LYS	-	expression tag	UNP P01903

- Molecule 2 is a protein called MHC class II antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	189	1557	977	281	294	5	0	0	0
2	E	189	1565	982	284	294	5	0	1	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	expression tag	UNP A0A0A1I7H6
B	0	SER	-	expression tag	UNP A0A0A1I7H6
B	191	THR	-	expression tag	UNP A0A0A1I7H6
B	192	GLY	-	expression tag	UNP A0A0A1I7H6
B	193	GLY	-	expression tag	UNP A0A0A1I7H6
B	194	ASP	-	expression tag	UNP A0A0A1I7H6
B	195	ASP	-	expression tag	UNP A0A0A1I7H6
B	196	ASP	-	expression tag	UNP A0A0A1I7H6
B	197	ASP	-	expression tag	UNP A0A0A1I7H6
B	198	LYS	-	expression tag	UNP A0A0A1I7H6
E	-1	GLY	-	expression tag	UNP A0A0A1I7H6
E	0	SER	-	expression tag	UNP A0A0A1I7H6
E	191	THR	-	expression tag	UNP A0A0A1I7H6
E	192	GLY	-	expression tag	UNP A0A0A1I7H6
E	193	GLY	-	expression tag	UNP A0A0A1I7H6
E	194	ASP	-	expression tag	UNP A0A0A1I7H6
E	195	ASP	-	expression tag	UNP A0A0A1I7H6
E	196	ASP	-	expression tag	UNP A0A0A1I7H6
E	197	ASP	-	expression tag	UNP A0A0A1I7H6
E	198	LYS	-	expression tag	UNP A0A0A1I7H6

- Molecule 3 is a protein called Vimentin59-71.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	C	13	Total	C	N	O	0	1	0
			103	63	22	18			
3	F	13	Total	C	N	O	0	1	0
			103	63	22	18			

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	G	2	Total	C	N	O	0	0	0
			28	16	2	10			
4	H	2	Total	C	N	O	0	0	0
			28	16	2	10			

- Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
5	A	1	14	8	1	5	0	0
5	D	1	14	8	1	5	0	0

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	B	1	Total C O 4 2 2	0	0
6	E	1	Total C O 4 2 2	0	0
6	E	1	Total C O 4 2 2	0	0
6	E	1	Total C O 4 2 2	0	0

- Molecule 7 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total C O 10 6 4	0	0
7	B	1	Total C O 10 6 4	0	0

- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	255	Total O 255 255	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	227	Total 227	O 227	0	0
8	D	256	Total 256	O 256	0	0
8	E	236	Total 236	O 236	0	0
8	C	21	Total 21	O 21	0	0
8	F	22	Total 22	O 22	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain

Chain A: 





- Molecule 1: HLA class II histocompatibility antigen, DR alpha chain

Chain D: 



- Molecule 2: MHC class II antigen

Chain B: 




- Molecule 2: MHC class II antigen

Chain E: 




- Molecule 3: Vimentin59-71

Chain C: 



- Molecule 3: Vimentin59-71

Chain F: 



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.98Å 77.77Å 94.54Å 90.00° 109.52° 90.00°	Depositor
Resolution (Å)	33.48 – 1.90 33.48 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (33.48-1.90) 97.0 (33.48-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.32 (at 1.89Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.170 , 0.202 0.172 , 0.202	Depositor DCC
R_{free} test set	3632 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	14.5	Xtrriage
Anisotropy	0.398	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 52.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.146 for h,-k,-h-l	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	7456	wwPDB-VP
Average B, all atoms (Å ²)	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, NAG, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/1545	0.46	0/2108
1	D	0.28	0/1533	0.46	0/2090
2	B	0.28	0/1600	0.46	0/2175
2	E	0.28	0/1611	0.48	1/2189 (0.0%)
3	C	0.25	0/106	0.44	0/141
3	F	0.26	0/106	0.47	0/141
All	All	0.28	0/6501	0.47	1/8844 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	55	ARG	NE-CZ-NH1	5.56	123.08	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1497	0	1424	3	0
1	D	1482	0	1416	3	0
2	B	1557	0	1464	15	0
2	E	1565	0	1477	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	103	0	114	2	0
3	F	103	0	114	3	0
4	G	28	0	25	1	0
4	H	28	0	25	1	0
5	A	14	0	13	0	0
5	D	14	0	13	0	0
6	B	16	0	24	1	0
6	E	12	0	18	2	0
7	B	20	0	28	3	0
8	A	255	0	0	3	0
8	B	227	0	0	5	0
8	C	21	0	0	1	0
8	D	256	0	0	1	0
8	E	236	0	0	5	0
8	F	22	0	0	3	0
All	All	7456	0	6155	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 41 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:2:ASP:N	4:G:2:NAG:HO6	1.69	0.90
2:E:55:ARG:HH11	2:E:55:ARG:HG3	1.35	0.89
2:B:43:ASP:HA	7:B:205:PGE:H2	1.76	0.68
2:E:19:ASN:ND2	8:E:303:HOH:O	2.24	0.66
2:E:94[A]:ARG:NH2	8:E:304:HOH:O	2.31	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/189 (96%)	180 (99%)	1 (1%)	0	100	100
1	D	179/189 (95%)	177 (99%)	2 (1%)	0	100	100
2	B	187/200 (94%)	183 (98%)	4 (2%)	0	100	100
2	E	188/200 (94%)	184 (98%)	4 (2%)	0	100	100
3	C	12/13 (92%)	11 (92%)	1 (8%)	0	100	100
3	F	12/13 (92%)	11 (92%)	1 (8%)	0	100	100
All	All	759/804 (94%)	746 (98%)	13 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	167/173 (96%)	166 (99%)	1 (1%)	86	87
1	D	166/173 (96%)	166 (100%)	0	100	100
2	B	172/179 (96%)	171 (99%)	1 (1%)	86	87
2	E	173/179 (97%)	171 (99%)	2 (1%)	71	70
3	C	11/10 (110%)	11 (100%)	0	100	100
3	F	11/10 (110%)	11 (100%)	0	100	100
All	All	700/724 (97%)	696 (99%)	4 (1%)	86	87

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92	LEU
2	B	127	ILE
2	E	55	ARG
2	E	127	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	G	1	1,4	14,14,15	0.47	0	17,19,21	0.47	0
4	NAG	G	2	4	14,14,15	0.25	0	17,19,21	0.40	0
4	NAG	H	1	1,4	14,14,15	0.52	0	17,19,21	0.44	0
4	NAG	H	2	4	14,14,15	0.46	0	17,19,21	0.51	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	G	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	NAG	H	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	H	2	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	G	2	NAG	C8-C7-N2-C2
4	G	2	NAG	O7-C7-N2-C2
4	H	2	NAG	C8-C7-N2-C2
4	H	2	NAG	O7-C7-N2-C2


There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	G	2	NAG	1	0
4	H	2	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	EDO	E	201	-	3,3,3	0.44	0	2,2,2	0.32	0
6	EDO	E	203	-	3,3,3	0.49	0	2,2,2	0.15	0
7	PGE	B	206	-	9,9,9	0.34	0	8,8,8	0.23	0
6	EDO	B	203	-	3,3,3	0.41	0	2,2,2	0.53	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	A	500	1	14,14,15	0.30	0	17,19,21	0.44	0
6	EDO	B	201	-	3,3,3	0.44	0	2,2,2	0.37	0
6	EDO	E	202	-	3,3,3	0.45	0	2,2,2	0.38	0
6	EDO	B	204	-	3,3,3	0.46	0	2,2,2	0.14	0
7	PGE	B	205	-	9,9,9	0.34	0	8,8,8	0.42	0
5	NAG	D	500	1	14,14,15	0.31	0	17,19,21	0.44	0
6	EDO	B	202	-	3,3,3	0.48	0	2,2,2	0.15	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	E	201	-	-	1/1/1/1	-
6	EDO	E	203	-	-	0/1/1/1	-
7	PGE	B	206	-	-	3/7/7/7	-
6	EDO	B	203	-	-	0/1/1/1	-
5	NAG	A	500	1	-	0/6/23/26	0/1/1/1
6	EDO	B	201	-	-	1/1/1/1	-
6	EDO	E	202	-	-	0/1/1/1	-
6	EDO	B	204	-	-	0/1/1/1	-
7	PGE	B	205	-	-	5/7/7/7	-
5	NAG	D	500	1	-	2/6/23/26	0/1/1/1
6	EDO	B	202	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 12 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	205	PGE	O1-C1-C2-O2
6	B	201	EDO	O1-C1-C2-O2
7	B	205	PGE	O3-C5-C6-O4
5	D	500	NAG	C4-C5-C6-O6
7	B	206	PGE	C1-C2-O2-C3

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	E	201	EDO	2	0
6	B	201	EDO	1	0
7	B	205	PGE	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	180/189 (95%)	-0.20	4 (2%) 62 64	9, 16, 39, 66	0
1	D	179/189 (94%)	-0.19	3 (1%) 70 72	8, 16, 38, 54	0
2	B	189/200 (94%)	-0.30	0 100 100	9, 15, 30, 38	0
2	E	189/200 (94%)	-0.34	0 100 100	9, 15, 30, 41	0
3	C	13/13 (100%)	-0.44	0 100 100	12, 14, 18, 34	0
3	F	13/13 (100%)	-0.47	0 100 100	11, 14, 18, 34	0
All	All	763/804 (94%)	-0.27	7 (0%) 84 85	8, 15, 35, 66	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	100	ARG	2.7
1	A	3	GLU	2.5
1	A	100	ARG	2.5
1	D	158	GLU	2.4
1	D	157	THR	2.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NAG	H	2	14/15	0.67	0.33	52,58,61,61	0
4	NAG	G	2	14/15	0.70	0.34	48,52,57,57	0
4	NAG	H	1	14/15	0.82	0.17	27,34,42,46	0
4	NAG	G	1	14/15	0.86	0.19	24,30,36,42	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
7	PGE	B	205	10/10	0.46	0.29	43,48,51,51	0
7	PGE	B	206	10/10	0.61	0.23	41,43,44,45	0
6	EDO	B	202	4/4	0.67	0.27	44,44,44,44	0
6	EDO	E	202	4/4	0.68	0.20	44,45,45,45	0
6	EDO	B	203	4/4	0.75	0.22	53,53,53,53	0
6	EDO	B	204	4/4	0.83	0.19	39,39,39,40	0
5	NAG	D	500	14/15	0.83	0.22	34,40,45,46	0
5	NAG	A	500	14/15	0.84	0.25	31,39,44,46	0
6	EDO	E	201	4/4	0.90	0.11	29,30,30,30	0
6	EDO	E	203	4/4	0.91	0.14	24,24,25,26	0
6	EDO	B	201	4/4	0.92	0.14	29,29,29,29	0

6.5 Other polymers [i](#)

There are no such residues in this entry.