



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 16, 2023 – 06:10 PM EDT

PDB ID : 2AV6
Title : X-Ray studies on maltodextrin phosphorylase complexes: recognition of substrates and catalytic mechanism of phosphorylase family
Authors : Geremia, S.; Campagnolo, M.
Deposited on : 2005-08-29
Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.01 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Clashscore	141614	9178 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	796	
1	B	796	
2	C	5	
2	D	5	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 14073 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Maltodextrin phosphorylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	796	6389	4079	1128	1162	20	0	0	0
1	B	796	6389	4079	1128	1162	20	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	261	ALA	HIS	engineered mutation	UNP P00490
A	262	PHE	THR	engineered mutation	UNP P00490
A	263	GLU	ALA	engineered mutation	UNP P00490
B	261	ALA	HIS	engineered mutation	UNP P00490
B	262	PHE	THR	engineered mutation	UNP P00490
B	263	GLU	ALA	engineered mutation	UNP P00490

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	C	5	56	30	26	0	0	0
2	D	5	56	30	26	0	0	0

- Molecule 3 is NITRATE ION (three-letter code: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	N	O	0	0
			4	1	3		
3	B	1	Total	N	O	0	0
			4	1	3		

- Molecule 4 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
4	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	587	Total 587	O 587	0	0
5	B	558	Total 558	O 558	0	0


3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Maltodextrin phosphorylase



- Molecule 1: Maltodextrin phosphorylase



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain C:  20% 80%

GLC1
GLC2
GLC3
GLC4
GLC5

- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose-(1-4)-beta-D-glucopyranose

Chain D:  100%

BGC1
GLC2
GLC3
GLC4
GLC5

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	74.96Å 105.61Å 218.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.01 19.20 – 2.01	Depositor EDS
% Data completeness (in resolution range)	86.3 (15.00-2.01) 86.4 (19.20-2.01)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.63 (at 2.01Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.190 , 0.244 0.200 , 0.194	Depositor DCC
R_{free} test set	5039 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	21.7	Xtrriage
Anisotropy	0.511	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 54.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14073	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.77% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NO3, PLP, GLC, BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.93	2/6539 (0.0%)	0.93	16/8865 (0.2%)
1	B	0.92	2/6539 (0.0%)	0.95	22/8865 (0.2%)
All	All	0.92	4/13078 (0.0%)	0.94	38/17730 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	136	TYR	CD1-CE1	5.91	1.48	1.39
1	A	273	TYR	CE2-CZ	-5.15	1.31	1.38
1	B	666	GLU	CD-OE2	5.04	1.31	1.25
1	A	777	MET	SD-CE	5.02	2.06	1.77

All (38) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	259	ASP	CB-CG-OD2	10.21	127.49	118.30
1	B	786	ASP	CB-CG-OD2	8.66	126.09	118.30
1	B	491	ASP	CB-CG-OD2	8.52	125.97	118.30
1	B	492	ASP	CB-CG-OD2	8.41	125.87	118.30
1	B	95	ASP	CB-CG-OD2	8.23	125.70	118.30
1	A	492	ASP	CB-CG-OD2	8.12	125.61	118.30
1	A	658	ASP	CB-CG-OD2	7.47	125.02	118.30
1	B	562	ASP	CB-CG-OD2	7.09	124.68	118.30
1	A	760	ASP	CB-CG-OD1	7.00	124.60	118.30
1	A	743	ASP	CB-CG-OD2	6.62	124.25	118.30
1	A	7	ASP	CB-CG-OD2	6.54	124.18	118.30
1	A	755	ASP	CB-CG-OD2	6.53	124.18	118.30
1	B	59	ARG	NE-CZ-NH2	-6.45	117.07	120.30
1	A	176	ASP	CB-CG-OD2	6.35	124.02	118.30
1	B	598	ASP	CB-CG-OD2	6.35	124.02	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	323	ASP	CB-CG-OD2	6.17	123.85	118.30
1	B	717	ASP	CB-CG-OD2	6.17	123.86	118.30
1	A	491	ASP	CB-CG-OD2	5.99	123.69	118.30
1	B	743	ASP	CB-CG-OD2	5.97	123.67	118.30
1	A	598	ASP	CB-CG-OD2	5.93	123.64	118.30
1	B	426	ASP	CB-CG-OD2	5.89	123.60	118.30
1	A	151	ASP	CB-CG-OD2	5.87	123.58	118.30
1	B	658	ASP	CB-CG-OD2	5.85	123.57	118.30
1	A	389	ASP	CB-CG-OD2	5.66	123.40	118.30
1	A	603	ASP	CB-CG-OD2	5.61	123.34	118.30
1	B	299	ASP	CB-CG-OD2	5.56	123.31	118.30
1	B	7	ASP	CB-CG-OD2	5.50	123.25	118.30
1	A	235	ASP	CB-CG-OD2	5.40	123.16	118.30
1	B	235	ASP	CB-CG-OD2	5.38	123.15	118.30
1	B	100	ASP	CB-CG-OD1	5.37	123.13	118.30
1	B	151	ASP	CB-CG-OD2	5.35	123.11	118.30
1	A	237	ASP	CB-CG-OD2	5.30	123.07	118.30
1	B	170	ARG	NE-CZ-NH1	5.29	122.95	120.30
1	A	356	ASP	CB-CG-OD2	5.29	123.06	118.30
1	B	160	ASP	CB-CG-OD2	5.28	123.05	118.30
1	A	786	ASP	CB-CG-OD2	5.20	122.98	118.30
1	B	755	ASP	CB-CG-OD2	5.17	122.95	118.30
1	B	700	ASP	CB-CG-OD2	5.05	122.84	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6389	0	6333	47	0
1	B	6389	0	6333	43	0
2	C	56	0	48	0	0
2	D	56	0	48	1	0
3	A	4	0	0	0	0
3	B	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	15	0	7	0	0
4	B	15	0	6	1	0
5	A	587	0	0	10	0
5	B	558	0	0	2	0
All	All	14073	0	12775	87	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (87) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:777:MET:CE	1:A:777:MET:SD	2.06	1.43
1:A:123:LEU:HD11	1:A:217:LEU:HD13	1.55	0.87
1:B:97:ASN:HD22	1:B:100:ASP:H	1.24	0.84
1:A:785:ARG:HG2	5:A:2546:HOH:O	1.76	0.83
1:B:188:ASP:C	1:B:188:ASP:OD1	2.22	0.76
1:A:470:LYS:NZ	1:A:498:GLN:HE22	1.85	0.74
1:B:470:LYS:NZ	1:B:498:GLN:HE22	1.86	0.73
1:B:551:LEU:O	1:B:555:ILE:HG23	1.88	0.72
1:A:97:ASN:HD22	1:A:100:ASP:H	1.36	0.72
1:B:470:LYS:HZ3	1:B:498:GLN:HE22	1.36	0.72
1:B:739:LEU:HB3	1:B:742:ALA:HB3	1.76	0.66
1:A:520:ILE:HD11	5:A:2551:HOH:O	1.96	0.64
1:A:551:LEU:O	1:A:555:ILE:HG23	1.98	0.62
1:A:488:LYS:HE3	5:A:2498:HOH:O	2.02	0.59
1:A:466:ALA:HB2	5:A:2481:HOH:O	2.02	0.58
1:A:47:ARG:HD3	5:A:2512:HOH:O	2.07	0.55
1:A:470:LYS:HZ2	1:A:498:GLN:HE22	1.53	0.55
1:A:470:LYS:HZ3	1:A:498:GLN:HE22	1.51	0.55
1:A:470:LYS:NZ	1:A:498:GLN:NE2	2.55	0.54
1:A:21:TYR:O	1:B:172:ASN:HB2	2.07	0.54
1:B:728:HIS:ND1	1:B:733:GLN:HG2	2.23	0.54
1:A:679:THR:OG1	1:A:682:GLN:HG3	2.08	0.53
1:B:109:ALA:HB1	1:B:143:GLY:HA3	1.91	0.52
1:A:555:ILE:HD13	1:A:604:LYS:HD3	1.92	0.52
1:B:123:LEU:HD11	1:B:217:LEU:HD13	1.92	0.51
1:B:7:ASP:O	1:B:11:GLN:HG2	2.11	0.51
1:B:679:THR:OG1	1:B:682:GLN:HG3	2.12	0.50
1:B:557:GLU:O	1:B:558:ASN:HB2	2.12	0.49
1:B:97:ASN:ND2	1:B:100:ASP:H	2.00	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123:LEU:CD1	1:A:217:LEU:HD13	2.35	0.49
1:A:507:LYS:NZ	1:A:526:ALA:O	2.38	0.49
1:A:700:ASP:HB3	1:A:703:LEU:HB3	1.94	0.49
1:B:425:SER:OG	1:B:446:ASN:ND2	2.36	0.49
1:B:470:LYS:NZ	1:B:498:GLN:NE2	2.60	0.48
1:B:67:GLU:HB2	1:B:111:GLY:HA2	1.95	0.48
1:A:109:ALA:HB1	1:A:143:GLY:HA3	1.96	0.47
1:B:188:ASP:OD1	1:B:189:GLY:N	2.47	0.47
1:B:544:ASN:C	1:B:544:ASN:HD22	2.18	0.47
1:B:655:GLY:O	1:B:675:ILE:HA	2.15	0.47
1:A:468:LEU:CD2	1:A:486:LEU:HD11	2.44	0.47
1:A:246:ILE:HD13	1:B:239:LEU:HD12	1.97	0.47
1:A:587:ALA:HA	1:A:723:PHE:CE1	2.49	0.47
1:B:700:ASP:HB3	1:B:703:LEU:HB3	1.96	0.47
1:B:723:PHE:O	1:B:727:LEU:HD22	2.15	0.46
1:B:532:ILE:HG21	1:B:621:LEU:HD13	1.98	0.46
1:B:703:LEU:HD11	1:B:741:MET:CE	2.45	0.45
1:A:544:ASN:ND2	5:A:2049:HOH:O	2.48	0.45
1:A:590:LYS:HA	1:A:590:LYS:HD2	1.82	0.45
1:A:237:ASP:OD1	5:A:2565:HOH:O	2.21	0.45
1:B:590:LYS:HA	1:B:590:LYS:HD2	1.78	0.45
1:B:483:LEU:O	1:B:486:LEU:HB2	2.17	0.45
1:A:529:ASP:OD1	1:A:629:GLU:OE2	2.34	0.44
1:B:3:PRO:HB3	1:B:49:GLN:OE1	2.17	0.44
1:A:635:GLY:N	1:A:680:VAL:HG22	2.32	0.44
1:A:220:TRP:CE2	1:A:278:CYS:HB3	2.53	0.44
1:A:220:TRP:CD2	1:A:278:CYS:HB3	2.53	0.44
1:B:112:ASN:HD22	2:D:3:GLC:H61	1.82	0.44
1:A:7:ASP:O	1:A:11:GLN:HG2	2.18	0.44
1:B:421:ALA:C	1:B:446:ASN:HD21	2.21	0.43
1:A:531:GLN:HE22	1:A:541:GLN:HA	1.82	0.43
1:A:657:LEU:HD22	1:A:662:VAL:HB	2.00	0.43
1:B:613:TYR:CE2	4:B:900:PLP:H2A2	2.53	0.43
1:A:671:GLU:H	1:A:671:GLU:CD	2.22	0.43
1:A:246:ILE:CD1	1:B:239:LEU:HD12	2.49	0.43
1:B:170:ARG:NH2	5:B:3092:HOH:O	2.51	0.43
1:B:555:ILE:HD13	1:B:604:LYS:HD3	2.00	0.43
1:A:544:ASN:C	1:A:544:ASN:HD22	2.23	0.42
1:B:691:TYR:CE2	1:B:739:LEU:HD22	2.54	0.42
1:B:187:LYS:O	1:B:188:ASP:HB3	2.19	0.42
1:B:309:HIS:N	1:B:310:PRO:CD	2.83	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:455:ARG:HA	1:A:459:GLN:HE21	1.84	0.41
1:A:532:ILE:HG21	1:A:621:LEU:HD13	2.02	0.41
1:B:47:ARG:HD3	5:B:3203:HOH:O	2.20	0.41
1:A:509:ARG:HD3	5:A:2548:HOH:O	2.20	0.41
1:B:559:PRO:HB2	1:B:560:GLN:NE2	2.35	0.41
1:A:691:TYR:CE2	1:A:739:LEU:HD22	2.55	0.41
1:B:531:GLN:HE22	1:B:541:GLN:HA	1.85	0.41
1:A:523:ASN:HA	1:A:524:PRO:HD3	1.98	0.41
1:A:606:LYS:HG3	5:A:2091:HOH:O	2.20	0.41
1:A:655:GLY:O	1:A:675:ILE:HA	2.21	0.41
1:B:452:THR:HA	1:B:453:PRO:HD3	1.91	0.41
1:A:447:VAL:HG11	1:A:787:TYR:CD2	2.57	0.40
1:B:433:PHE:N	1:B:434:PRO:CD	2.84	0.40
1:A:400:HIS:HD2	5:A:2241:HOH:O	2.05	0.40
1:A:452:THR:HA	1:A:453:PRO:HD3	1.86	0.40
1:A:529:ASP:HA	1:A:627:ILE:HB	2.02	0.40
1:B:468:LEU:CD2	1:B:486:LEU:HD11	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	794/796 (100%)	770 (97%)	23 (3%)	1 (0%)	51	49
1	B	794/796 (100%)	765 (96%)	26 (3%)	3 (0%)	34	30
All	All	1588/1592 (100%)	1535 (97%)	49 (3%)	4 (0%)	41	37

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	188	ASP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	559	PRO
1	B	558	ASN
1	A	533	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	667/667 (100%)	631 (95%)	36 (5%)	22	18
1	B	667/667 (100%)	632 (95%)	35 (5%)	23	19
All	All	1334/1334 (100%)	1263 (95%)	71 (5%)	22	18

All (71) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92	LYS
1	A	117	ARG
1	A	139	ASN
1	A	237	ASP
1	A	289	LEU
1	A	363	LEU
1	A	392	VAL
1	A	395	LYS
1	A	401	ASP
1	A	423	LEU
1	A	438	GLN
1	A	443	LYS
1	A	473	GLN
1	A	494	LYS
1	A	516	VAL
1	A	525	GLN
1	A	535	LEU
1	A	538	TYR
1	A	542	HIS
1	A	544	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	555	ILE
1	A	562	ASP
1	A	564	VAL
1	A	578	TYR
1	A	590	LYS
1	A	609	PHE
1	A	657	LEU
1	A	689	LYS
1	A	695	LYS
1	A	699	LYS
1	A	727	LEU
1	A	732	LYS
1	A	739	LEU
1	A	755	ASP
1	A	785	ARG
1	A	796	ARG
1	B	4	ILE
1	B	117	ARG
1	B	139	ASN
1	B	188	ASP
1	B	190	ARG
1	B	237	ASP
1	B	289	LEU
1	B	363	LEU
1	B	392	VAL
1	B	395	LYS
1	B	401	ASP
1	B	423	LEU
1	B	438	GLN
1	B	443	LYS
1	B	454	ARG
1	B	494	LYS
1	B	516	VAL
1	B	525	GLN
1	B	533	LYS
1	B	535	LEU
1	B	542	HIS
1	B	544	ASN
1	B	555	ILE
1	B	564	VAL
1	B	578	TYR
1	B	609	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	657	LEU
1	B	689	LYS
1	B	695	LYS
1	B	699	LYS
1	B	727	LEU
1	B	732	LYS
1	B	739	LEU
1	B	755	ASP
1	B	785	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (29) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	GLN
1	A	9	GLN
1	A	97	ASN
1	A	112	ASN
1	A	139	ASN
1	A	178	GLN
1	A	221	GLN
1	A	260	ASN
1	A	400	HIS
1	A	446	ASN
1	A	459	GLN
1	A	498	GLN
1	A	525	GLN
1	A	531	GLN
1	A	544	ASN
1	B	2	GLN
1	B	9	GLN
1	B	97	ASN
1	B	112	ASN
1	B	139	ASN
1	B	178	GLN
1	B	260	ASN
1	B	446	ASN
1	B	459	GLN
1	B	498	GLN
1	B	525	GLN
1	B	531	GLN
1	B	544	ASN
1	B	560	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	BGC	C	1	2	12,12,12	0.84	0	17,17,17	0.78	0
2	GLC	C	2	2	11,11,12	1.14	1 (9%)	15,15,17	1.04	2 (13%)
2	GLC	C	3	2	11,11,12	1.46	1 (9%)	15,15,17	0.94	1 (6%)
2	GLC	C	4	2	11,11,12	0.90	1 (9%)	15,15,17	1.77	4 (26%)
2	GLC	C	5	2	11,11,12	0.88	1 (9%)	15,15,17	2.54	3 (20%)
2	BGC	D	1	2	12,12,12	0.80	1 (8%)	17,17,17	0.69	0
2	GLC	D	2	2	11,11,12	0.68	0	15,15,17	1.21	2 (13%)
2	GLC	D	3	2	11,11,12	0.75	0	15,15,17	0.71	0
2	GLC	D	4	2	11,11,12	0.90	0	15,15,17	2.03	3 (20%)
2	GLC	D	5	2	11,11,12	0.82	0	15,15,17	2.24	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	1	2	-	0/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	C	3	2	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	4	2	-	2/2/19/22	0/1/1/1
2	GLC	C	5	2	-	0/2/19/22	0/1/1/1
2	BGC	D	1	2	-	1/2/22/22	0/1/1/1
2	GLC	D	2	2	-	0/2/19/22	0/1/1/1
2	GLC	D	3	2	-	0/2/19/22	0/1/1/1
2	GLC	D	4	2	-	2/2/19/22	0/1/1/1
2	GLC	D	5	2	-	0/2/19/22	0/1/1/1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	3	GLC	O5-C1	-3.12	1.38	1.43
2	C	2	GLC	C2-C3	2.94	1.56	1.52
2	C	5	GLC	C2-C3	2.19	1.55	1.52
2	C	4	GLC	C2-C3	2.10	1.55	1.52
2	D	1	BGC	O5-C5	-2.02	1.39	1.44

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	5	GLC	C1-O5-C5	6.49	120.98	112.19
2	D	5	GLC	C1-C2-C3	6.28	117.38	109.67
2	C	5	GLC	C1-C2-C3	6.16	117.24	109.67
2	D	4	GLC	C1-O5-C5	6.06	120.40	112.19
2	C	4	GLC	C1-O5-C5	5.19	119.22	112.19
2	D	5	GLC	C1-O5-C5	4.45	118.23	112.19
2	D	4	GLC	O4-C4-C3	3.29	117.96	110.35
2	D	4	GLC	O2-C2-C3	3.14	116.42	110.14
2	D	2	GLC	C1-C2-C3	2.50	112.73	109.67
2	C	2	GLC	C1-C2-C3	2.35	112.55	109.67
2	C	2	GLC	O5-C5-C6	2.29	110.80	107.20
2	C	4	GLC	O4-C4-C3	2.27	115.61	110.35
2	C	5	GLC	O5-C5-C6	2.21	110.66	107.20
2	C	4	GLC	O2-C2-C3	2.20	114.54	110.14
2	D	2	GLC	O5-C5-C6	2.18	110.62	107.20
2	C	3	GLC	O2-C2-C1	2.15	113.56	109.15
2	C	4	GLC	O5-C1-C2	2.06	113.95	110.77

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	4	GLC	C4-C5-C6-O6
2	C	4	GLC	O5-C5-C6-O6
2	D	4	GLC	C4-C5-C6-O6
2	D	4	GLC	O5-C5-C6-O6
2	D	1	BGC	C4-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	3	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry i

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NO3	A	1999	-	1,3,3	3.68	1 (100%)	0,3,3	-	-
4	PLP	B	900	1	15,15,16	1.24	1 (6%)	20,22,23	1.80	7 (35%)
3	NO3	B	2999	-	1,3,3	4.21	1 (100%)	0,3,3	-	-
4	PLP	A	900	1	15,15,16	1.69	3 (20%)	20,22,23	2.05	7 (35%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PLP	B	900	1	-	2/6/6/8	0/1/1/1
4	PLP	A	900	1	-	1/6/6/8	0/1/1/1

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	900	PLP	C3-C2	-4.24	1.36	1.40
3	B	2999	NO3	O1-N	4.21	1.43	1.24
3	A	1999	NO3	O1-N	3.68	1.41	1.24
4	A	900	PLP	C4A-C4	-2.99	1.45	1.51
4	B	900	PLP	C5-C4	-2.48	1.37	1.40
4	A	900	PLP	C2-N1	2.21	1.38	1.33

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	900	PLP	C3-C4-C5	4.88	124.02	118.74
4	B	900	PLP	C3-C4-C5	3.49	122.51	118.74
4	A	900	PLP	C4A-C4-C3	-3.29	114.92	120.50

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	900	PLP	C3-C2-N1	-3.27	116.54	120.77
4	A	900	PLP	C6-C5-C4	-3.23	115.61	118.16
4	B	900	PLP	C6-C5-C4	-3.23	115.61	118.16
4	B	900	PLP	C2A-C2-C3	2.74	124.27	120.89
4	A	900	PLP	C6-N1-C2	2.49	123.78	119.17
4	A	900	PLP	C3-C2-N1	-2.49	117.56	120.77
4	A	900	PLP	O4P-P-O1P	2.37	113.11	106.47
4	B	900	PLP	C6-N1-C2	2.36	123.55	119.17
4	A	900	PLP	C5-C6-N1	-2.22	120.12	123.82
4	B	900	PLP	O2P-P-O1P	2.02	118.60	110.68
4	B	900	PLP	C4A-C4-C3	-2.00	117.11	120.50

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	900	PLP	C5A-O4P-P-O1P
4	A	900	PLP	C6-C5-C5A-O4P
4	B	900	PLP	C5A-O4P-P-O3P

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	900	PLP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	796/796 (100%)	-0.09	13 (1%) 72 70	14, 25, 44, 59	0
1	B	796/796 (100%)	-0.05	22 (2%) 53 51	14, 25, 46, 59	0
All	All	1592/1592 (100%)	-0.07	35 (2%) 62 60	14, 25, 45, 59	0

All (35) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	562	ASP	5.2
1	B	560	GLN	4.9
1	B	57	ASN	4.5
1	B	559	PRO	3.6
1	A	51	PHE	3.6
1	B	187	LYS	3.6
1	B	561	ALA	3.2
1	B	579	TYR	3.1
1	B	718	GLY	3.1
1	B	688	ALA	2.7
1	A	699	LYS	2.6
1	A	289	LEU	2.6
1	B	188	ASP	2.6
1	A	701	LYS	2.6
1	A	85	GLN	2.6
1	B	716	SER	2.5
1	B	717	ASP	2.5
1	B	4	ILE	2.4
1	A	1	SER	2.4
1	B	390	GLU	2.4
1	B	689	LYS	2.3
1	B	391	LYS	2.3
1	A	688	ALA	2.3
1	A	762	GLU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	555	ILE	2.3
1	A	795	LYS	2.2
1	B	600	LEU	2.2
1	B	190	ARG	2.2
1	B	51	PHE	2.1
1	B	695	LYS	2.1
1	A	733	GLN	2.1
1	A	396	LEU	2.1
1	A	516	VAL	2.0
1	B	384	LYS	2.0
1	B	85	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	C	5	11/12	0.91	0.15	31,35,37,37	0
2	BGC	D	1	12/12	0.92	0.21	32,34,37,42	0
2	GLC	D	2	11/12	0.92	0.12	20,24,28,28	0
2	BGC	C	1	12/12	0.94	0.13	33,34,37,42	0
2	GLC	D	4	11/12	0.94	0.10	24,27,30,31	0
2	GLC	D	5	11/12	0.94	0.13	32,35,36,37	0
2	GLC	C	2	11/12	0.95	0.09	22,23,27,28	0
2	GLC	D	3	11/12	0.95	0.13	20,23,24,24	0
2	GLC	C	4	11/12	0.96	0.08	24,27,29,31	0
2	GLC	C	3	11/12	0.97	0.08	19,23,24,24	0




The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Chain C:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Chain D:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	NO3	B	2999	4/4	0.83	0.23	42,45,46,46	0
3	NO3	A	1999	4/4	0.94	0.15	42,45,45,46	0
4	PLP	B	900	15/16	0.97	0.08	14,19,32,35	0
4	PLP	A	900	15/16	0.98	0.09	15,20,32,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.