

wwPDB NMR Structure Validation Summary Report (i)

Jun 6, 2023 – 04:49 pm BST

PDB ID : 7AVB BMRB ID : 34573

Title: Solution structure of the fluorogen-activating protein FAST in the apo state

Authors: Mineev, K.S.; Goncharuk, S.A.; Baranov, M.S.

Deposited on : 2020-11-05

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

wwPDB-RCI : v 1n 11 5 13 A (Berjanski et al., 2005)

PANAV : Wang et al. (2010)

 $\begin{array}{ccc} wwPDB\text{-}ShiftChecker &: & v1.2 \\ BMRB \ Restraints \ Analysis &: & v1.2 \\ \end{array}$

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.33

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 93%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$egin{array}{c} { m NMR \ archive} \ (\#{ m Entries}) \end{array}$	
Clashscore	158937	12864	
Ramachandran outliers	154571	11451	
Sidechain outliers	154315	11428	

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain			
1	A	138	38%	13%	49%	



2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 2 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: fewest violations.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues						
Well-defined core	Backbone RMSD (Å)	Medoid model				
1	A:29-A:42, A:58-A:94,	0.43	2			
	A:105-A:124 (71)					

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 3 single-model clusters were found.

Cluster number	Models
1	3, 6, 7, 9, 10, 14, 15
2	2, 4, 11, 16, 20
3	1, 8, 18
4	17, 19
Single-model clusters	5; 12; 13



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2106 atoms, of which 1033 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called FAST.

Mol	Chain	Residues	Atoms				Trace		
1	٨	138	Total	С	Н	N	О	S	0
1	А	130	2106	680	1033	187	201	5	U




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.






4.2 Residue scores for the representative (medoid) model from the NMR ensemble


The representative model is number 2. Colouring as in section 4.1 above.

• Molecule 1: FAST





5 Refinement protocol and experimental data overview (i)



The models were refined using the following method: torsion angle dynamics.

Of the 200 calculated structures, 20 were deposited, based on the following criterion: structures with the least restraint violations.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	3.98.13

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1574
Number of shifts mapped to atoms	1574
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	93%



6 Model quality (i)

6.1 Standard geometry (i)

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	562	547	547	11±3
All	All	11240	10940	10940	223

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

5 of 40 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:83:VAL:HG22	1:A:118:TYR:CZ	0.67	2.24	4	13
1:A:63:PHE:HA	1:A:67:ALA:HB3	0.67	1.67	18	20
1:A:30:ALA:HB2	1:A:121:PHE:CE2	0.66	2.25	16	8
1:A:30:ALA:HB2	1:A:121:PHE:CD1	0.63	2.28	14	1
1:A:79:PHE:O	1:A:83:VAL:HG23	0.62	1.95	18	20

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	71/138 (51%)	67±1 (94±1%)	3±1 (5±1%)	1±1 (1±1%)	16	63
All	All	1420/2760 (51%)	1334 (94%)	68 (5%)	18 (1%)	16	63

All 2 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	87	ASN	15
1	A	116	ASP	3

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed Rotameric		Outliers	Percentiles		
1	A	58/111 (52%)	51±2 (88±4%)	7±2 (12±4%)	9	52	
All	All	1160/2220 (52%)	1025 (88%)	135 (12%)	9	52	

5 of 28 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	117	SER	18
1	A	94	TRP	16
1	A	72	SER	14
1	A	111	LYS	8
1	A	123	LYS	8

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 93% for the well-defined parts and 87% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: assigned_chem_shift_list_1

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1574
Number of shifts mapped to atoms	1574
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	2

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	${\rm Correction} \pm {\rm precision}, ppm$	Suggested action
$^{13}\mathrm{C}_{\alpha}$	134	0.15 ± 0.08	None needed ($< 0.5 \text{ ppm}$)
$^{13}C_{\beta}$	120	0.09 ± 0.17	None needed ($< 0.5 \text{ ppm}$)
¹³ C′	112	0.42 ± 0.08	None needed (< 0.5 ppm)
^{15}N	106	-0.10 ± 0.20	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 93%, i.e. 884 atoms were assigned a chemical shift out of a possible 954. 0 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}{ m H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Backbone	346/360 (96%)	144/149 (97%)	138/142 (97%)	64/69 (93%)
Sidechain	444/476 (93%)	302/308 (98%)	136/151 (90%)	6/17 (35%)

Continued on next page...



Continued from previous page...

	Total	$^{1}\mathrm{H}$	$^{13}\mathbf{C}$	$^{15}{ m N}$
Aromatic	94/118 (80%)	56/58 (97%)	36/57~(63%)	2/3 (67%)
Overall	884/954 (93%)	502/515 (97%)	310/350~(89%)	72/89 (81%)

7.1.4 Statistically unusual chemical shifts (i)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	32	GLN	HB3	-0.13	0.71 - 3.33	-8.2
1	A	32	GLN	HG3	0.77	0.91 - 3.68	-5.5

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports random coil index values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1033
Intra-residue ($ i-j =0$)	245
Sequential ($ i-j =1$)	275
Medium range ($ i-j >1$ and $ i-j <5$)	131
Long range (i-j ≥5)	382
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	7.5
Number of long range restraints per residue ¹	2.8

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	1.2	0.18
0.2-0.5 (Medium)	0.1	0.23
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations



9 Distance violation analysis (i)

9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Doctroints type	Count	% ¹	Vio	lated	3	Consis	tentl	${ m y~Violated^4}$
Restraints type	Count	70	Count	$\%^2$	$\%^1$	Count	$ \%^2 $	$\%^1$
Intra-residue (i-j =0)	245	23.7	3	1.2	0.3	0	0.0	0.0
Backbone-Backbone	4	0.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	221	21.4	3	1.4	0.3	0	0.0	0.0
Sidechain-Sidechain	20	1.9	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	275	26.6	5	1.8	0.5	0	0.0	0.0
Backbone-Backbone	65	6.3	1	1.5	0.1	0	0.0	0.0
Backbone-Sidechain	198	19.2	3	1.5	0.3	0	0.0	0.0
Sidechain-Sidechain	12	1.2	1	8.3	0.1	0	0.0	0.0
Medium range ($ i-j >1 \& i-j <5$)	131	12.7	2	1.5	0.2	0	0.0	0.0
Backbone-Backbone	49	4.7	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	57	5.5	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	25	2.4	2	8.0	0.2	0	0.0	0.0
Long range ($ i-j \ge 5$)	382	37.0	1	0.3	0.1	0	0.0	0.0
Backbone-Backbone	72	7.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	179	17.3	1	0.6	0.1	0	0.0	0.0
Sidechain-Sidechain	131	12.7	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1033	100.0	11	1.1	1.1	0	0.0	0.0
Backbone-Backbone	190	18.4	1	0.5	0.1	0	0.0	0.0
Backbone-Sidechain	655	63.4	7	1.1	0.7	0	0.0	0.0
Sidechain-Sidechain	188	18.2	3	1.6	0.3	0	0.0	0.0

 $^{^1}$ percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models



9.1.1 Bar chart: Distribution of distance restraints and violations (i)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

MadalID		Nun	nber o	f viola	ations	5	M (8)	N/ (${ m SD}^6$ (Å)	Madian (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (Å)	Max (Å)	$SD^*(A)$	Median (Å)
1	0	0	1	0	0	1	0.13	0.13	0.0	0.13
2	0	0	0	0	0	0	0.0	0.0	0.0	0.0
3	0	0	1	0	0	1	0.11	0.11	0.0	0.11
4	0	0	0	0	0	0	0.0	0.0	0.0	0.0
5	0	1	0	0	0	1	0.17	0.17	0.0	0.17
6	0	0	1	0	0	1	0.12	0.12	0.0	0.12
7	0	0	1	0	0	1	0.15	0.15	0.0	0.15
8	0	0	1	0	0	1	0.11	0.11	0.0	0.11
9	0	1	1	0	0	2	0.14	0.14	0.01	0.14
10	1	0	1	0	0	2	0.12	0.13	0.01	0.12
11	0	0	0	0	0	0	0.0	0.0	0.0	0.0

Continued on next page...




Continued from previous page...

Model ID		Nun	nber o	f viola	ations	3	Mean (Å)	Max (Å)	${ m SD}^6$ (Å)	Median (Å)
Model 1D	IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Mean (A)	Max (A)	$SD^*(A)$	Median (A)
12	0	0	1	0	0	1	0.12	0.12	0.0	0.12
13	2	2	0	0	0	4	0.16	0.23	0.05	0.14
14	1	1	2	0	0	4	0.15	0.18	0.03	0.15
15	0	1	1	0	0	2	0.14	0.15	0.01	0.14
16	2	0	0	0	0	2	0.12	0.13	0.01	0.12
17	1	0	0	1	0	2	0.18	0.22	0.04	0.18
18	0	0	0	0	0	0	0.0	0.0	0.0	0.0
19	0	0	0	1	0	1	0.17	0.17	0.0	0.17
20	0	0	0	0	0	0	0.0	0.0	0.0	0.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Standard deviation

9.2.1 Bar graph: Distance Violation statistics for each model (i)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right



9.3 Distance violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1022(IR:242, SQ:270, MR:129, LR:381, IC:0) restraints are not violated in the ensemble.

Nu	$\overline{\mathbf{mber}}$	of vio	lated	Fraction of the ensemble			
IR^1	SQ^2	MR^3	LR^4	IC^5	Total	Count ⁶	%
1	4	1	0	0	6	1	5.0
1	1	0	1	0	3	2	10.0
0	0	0	0	0	0	3	15.0
1	0	0	0	0	1	4	20.0
0	0	0	0	0	0	5	25.0
0	0	0	0	0	0	6	30.0
0	0	0	0	0	0	7	35.0
0	0	0	0	0	0	8	40.0
0	0	0	0	0	0	9	45.0
0	0	1	0	0	1	10	50.0
0	0	0	0	0	0	11	55.0
0	0	0	0	0	0	12	60.0
0	0	0	0	0	0	13	65.0
0	0	0	0	0	0	14	70.0
0	0	0	0	0	0	15	75.0
0	0	0	0	0	0	16	80.0
0	0	0	0	0	0	17	85.0
0	0	0	0	0	0	18	90.0
0	0	0	0	0	0	19	95.0
0	0	0	0	0	0	20	100.0

 $^{^1}$ Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph: Distance violation statistics for the ensemble (i)




9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram: Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,1003)	1:A:105:VAL:HG21	1:A:107:VAL:HG11	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG21	1:A:107:VAL:HG12	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG21	1:A:107:VAL:HG13	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG21	1:A:107:VAL:HG21	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG21	1:A:107:VAL:HG22	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG21	1:A:107:VAL:HG23	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG22	1:A:107:VAL:HG11	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG22	1:A:107:VAL:HG12	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG22	1:A:107:VAL:HG13	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG22	1:A:107:VAL:HG21	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG22	1:A:107:VAL:HG22	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG22	1:A:107:VAL:HG23	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG23	1:A:107:VAL:HG11	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG23	1:A:107:VAL:HG12	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG23	1:A:107:VAL:HG13	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG23	1:A:107:VAL:HG21	10	0.12	0.01	0.12

Continued on next page...



Continued from previous page...


Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,1003)	1:A:105:VAL:HG23	1:A:107:VAL:HG22	10	0.12	0.01	0.12
(1,1003)	1:A:105:VAL:HG23	1:A:107:VAL:HG23	10	0.12	0.01	0.12
(1,814)	1:A:23:LEU:HA	1:A:23:LEU:HD11	4	0.12	0.01	0.12
(1,814)	1:A:23:LEU:HA	1:A:23:LEU:HD12	4	0.12	0.01	0.12
(1,814)	1:A:23:LEU:HA	1:A:23:LEU:HD13	4	0.12	0.01	0.12
(1,814)	1:A:23:LEU:HA	1:A:23:LEU:HD21	4	0.12	0.01	0.12
(1,814)	1:A:23:LEU:HA	1:A:23:LEU:HD22	4	0.12	0.01	0.12
(1,814)	1:A:23:LEU:HA	1:A:23:LEU:HD23	4	0.12	0.01	0.12
(1,210)	1:A:90:THR:H	1:A:109:MET:HG2	2	0.2	0.02	0.2
(1,210)	1:A:90:THR:H	1:A:109:MET:HG3	2	0.2	0.02	0.2
(1,989)	1:A:100:ARG:HB2	1:A:101:GLY:H	2	0.16	0.02	0.16
(1,989)	1:A:100:ARG:HB3	1:A:101:GLY:H	2	0.16	0.02	0.16
(1,373)	1:A:24:ASP:H	1:A:24:ASP:HB2	2	0.14	0.01	0.14
(1,373)	1:A:24:ASP:H	1:A:24:ASP:HB3	2	0.14	0.01	0.14

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





9.5.2 Table: All distance violations (i)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,372)	1:A:23:LEU:HA	1:A:24:ASP:H	13	0.23
(1,210)	1:A:90:THR:H	1:A:109:MET:HG2	17	0.22
(1,210)	1:A:90:THR:H	1:A:109:MET:HG3	17	0.22
(1,989)	1:A:100:ARG:HB2	1:A:101:GLY:H	14	0.18
(1,989)	1:A:100:ARG:HB3	1:A:101:GLY:H	14	0.18
(1,986)	1:A:98:THR:HG21	1:A:100:ARG:HB2	14	0.18
(1,986)	1:A:98:THR:HG21	1:A:100:ARG:HB3	14	0.18
(1,986)	1:A:98:THR:HG22	1:A:100:ARG:HB2	14	0.18
(1,986)	1:A:98:THR:HG22	1:A:100:ARG:HB3	14	0.18
(1,986)	1:A:98:THR:HG23	1:A:100:ARG:HB2	14	0.18
(1,986)	1:A:98:THR:HG23	1:A:100:ARG:HB3	14	0.18
(1,905)	1:A:58:ILE:HG12	1:A:59:GLY:H	5	0.17
(1,905)	1:A:58:ILE:HG13	1:A:59:GLY:H	5	0.17
(1,210)	1:A:90:THR:H	1:A:109:MET:HG2	19	0.17
(1,210)	1:A:90:THR:H	1:A:109:MET:HG3	19	0.17
(1,989)	1:A:100:ARG:HB2	1:A:101:GLY:H	15	0.15
(1,989)	1:A:100:ARG:HB3	1:A:101:GLY:H	15	0.15
(1,373)	1:A:24:ASP:H	1:A:24:ASP:HB2	13	0.15
(1,373)	1:A:24:ASP:H	1:A:24:ASP:HB3	13	0.15
(1,1003)	1:A:105:VAL:HG21	1:A:107:VAL:HG11	7	0.15
(1,1003)	1:A:105:VAL:HG21	1:A:107:VAL:HG12	7	0.15
(1,1003)	1:A:105:VAL:HG21	1:A:107:VAL:HG13	7	0.15
(1,1003)	1:A:105:VAL:HG21	1:A:107:VAL:HG21	7	0.15
(1,1003)	1:A:105:VAL:HG21	1:A:107:VAL:HG22	7	0.15
(1,1003)	1:A:105:VAL:HG21	1:A:107:VAL:HG23	7	0.15
(1,1003)	1:A:105:VAL:HG22	1:A:107:VAL:HG11	7	0.15
(1,1003)	1:A:105:VAL:HG22	1:A:107:VAL:HG12	7	0.15
(1,1003)	1:A:105:VAL:HG22	1:A:107:VAL:HG13	7	0.15
(1,1003)	1:A:105:VAL:HG22	1:A:107:VAL:HG21	7	0.15
(1,1003)	1:A:105:VAL:HG22	1:A:107:VAL:HG22	7	0.15
(1,1003)	1:A:105:VAL:HG22	1:A:107:VAL:HG23	7	0.15
(1,1003)	1:A:105:VAL:HG23	1:A:107:VAL:HG11	7	0.15
(1,1003)	1:A:105:VAL:HG23	1:A:107:VAL:HG12	7	0.15
(1,1003)	1:A:105:VAL:HG23	1:A:107:VAL:HG13	7	0.15
(1,1003)	1:A:105:VAL:HG23	1:A:107:VAL:HG21	7	0.15
(1,1003)	1:A:105:VAL:HG23	1:A:107:VAL:HG22	7	0.15
(1,1003)	1:A:105:VAL:HG23	1:A:107:VAL:HG23	7	0.15
(1,426)	1:A:77:GLY:H	1:A:78:LYS:HB2	9	0.14



10 Dihedral-angle violation analysis (i)

Dihedral angle analysis failed due to data error in the dihedral angle restraints, possibly missing target value

