

wwPDB X-ray Structure Validation Summary Report (i)

May 24, 2022 – 10:11 am BST

PDB ID : 7AVO

Title: Structure of marine actinobacteria clade rhodopsin (MacR) in orange form in

P1211 space group

Authors: Gushchin, I.; Polovinkin, V.; Kovalev, K.; Shevchenko, V.; Gordeliy, V.

Deposited on : 2020-11-05

Resolution : 1.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.4, CSD as 541 be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.28.1buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0267$

CCP4 : 7.1.010 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.28.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar \ resolution} \\ (\#{\rm Entries, \ resolution \ range(\AA)}) \end{array}$
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	220	81%	16%	:
1	В	220	6% 82%	15%	••



2 Entry composition (i)


There are 5 unique types of molecules in this entry. The entry contains 3841 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Bacteriorhodopsin.

M	[o]	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
	1	Λ	217	Total	С	Ν	О	S	0	0	0
-	1	Λ	211	1737	1147	274	303	13	0	9	U
	1	B	217	Total	С	N	О	S	2	6	0
-	1	D	211	1706	1126	269	299	12	3	0	

• Molecule 2 is EICOSANE (three-letter code: LFA) (formula: $C_{20}H_{42}$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C 10 10	0	0
2	A	1	Total C 16 16	0	0
2	A	1	Total C 10 10	0	0
2	A	1	Total C 10 10	0	0

Continued on next page...




Continued from previous page...

Mol		$oxed{\mathbf{Residues}}$	Atoms	ZeroOcc	AltConf
2	A	1	Total C	0	0
			16 16 Total C		
2	A	1	16 16	0	0
2	A	1	Total C	0	0
	Λ	1	8 8	0	0
2	A	1	Total C 8 8	0	0
0	Λ	1	Total C	0	0
2	A	1	6 6	0	0
2	A	1	Total C	0	0
			10 10 Total C		
2	A	1	10 10	0	0
2	A	1	Total C	0	0
	11	1	6 6		
2	В	1	Total C 10 10	0	0
9	D	1	Total C	0	0
2	В	1	16 16	0	0
2	В	1	Total C 14 14	0	0
			14 14 Total C		
2	В	1	5 5	0	0
2	В	1	Total C	0	0
		1	9 9 Total C		
2	В	1	Total C 16 16	0	0
9	D	1	Total C	0	0
2	В	1	8 8	0	0
2	В	1	Total C	0	0
			12 12 Total C		
2	В	1	16 16	0	0
2	В	1	Total C	0	0
		1	10 10		
2	В	1	Total C 10 10	0	0
	D	1	Total C	0	0
2	В	1	6 6	0	0


 \bullet Molecule 3 is (2S)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLB) (formula: $C_{21}H_{40}O_4).$





Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
3	A	1	Total 15	C 11	O 4	0	0

 \bullet Molecule 4 is (2R)-2,3-dihydroxypropyl (9Z)-octadec-9-enoate (three-letter code: OLC) (formula: $C_{21}H_{40}O_4).$



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
4	В	1	Total 10	C 6	O 4	0	0

• Molecule 5 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	56	Total O 56 56	0	0
5	В	59	Total O 59 59	0	1



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Bacteriorhodopsin





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	40.69Å 102.01Å 56.60Å	Donositor
a, b, c, α , β , γ	90.00° 99.75° 90.00°	Depositor
Resolution (Å)	48.94 - 1.70	Depositor
Resolution (A)	48.94 - 1.70	EDS
% Data completeness	97.8 (48.94-1.70)	Depositor
(in resolution range)	97.9 (48.94-1.70)	EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.22 (at 1.70Å)	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
D D.	0.197 , 0.222	Depositor
R, R_{free}	0.207 , 0.230	DCC
R_{free} test set	2481 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	22.8	Xtriage
Anisotropy	0.314	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	(Not available), (Not available)	EDS
L-test for twinning ²	$ < L > = 0.58, < L^2 > = 0.43$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3841	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 44.41 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5334e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: OLB, LFA, LYR, OLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.63	0/1770	0.63	0/2407
1	В	0.63	0/1733	0.62	0/2360
All	All	0.63	0/3503	0.62	0/4767

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1737	0	1760	39	0
1	В	1706	0	1712	37	0
2	A	126	0	237	18	0
2	В	132	0	249	8	0
3	A	15	0	19	2	0
4	В	10	0	9	4	0
5	A	56	0	0	10	0
5	В	59	0	0	10	0
All	All	3841	0	3986	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.



The worst 5 of 88 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} & & & & & & & & & & & & & & & & & & &$	Clash overlap (Å)
1:A:143:LEU:HD23	5:A:424:HOH:O	1.43	1.17
4:B:313:OLC:O25	5:B:401:HOH:O	1.67	1.10
2:A:304:LFA:C1	2:A:309:LFA:H11	1.79	1.10
2:A:304:LFA:H11	2:A:309:LFA:H11	1.32	1.10
1:A:209[A]:GLN:HE21	2:A:306:LFA:C16	1.71	1.02

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Favoured Allowed C		Outliers	Perce	ntiles	
1	A	222/220 (101%)	221 (100%)	1 (0%)	0	100	100
1	В	220/220 (100%)	220 (100%)	0	0	100	100
All	All	442/440 (100%)	441 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/172 (100%)	170 (99%)	2 (1%)	71 59	

Continued on next page...



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	В	167/172 (97%)	164 (98%)	3 (2%)	59	43	
All	All	339/344 (98%)	334 (98%)	5 (2%)	71	51	

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1[A]	MET
1	A	1[B]	MET
1	В	187	ILE
1	В	188	ARG
1	В	208	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	В	34	HIS
1	В	142	ASN
1	В	154	GLN
1	A	142	ASN
1	A	123	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

١,	Mol	Type	Chain	Dec	Tiple	Bo	ond leng	an	B	ond angl	es
	VIOI	туре	Cham	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
	1	LYR	A	200	1	27,29,30	1.27	4 (14%)	30,37,39	1.89	8 (26%)



Mol	Type	Chain	Pog	Link	Bo	ond leng	$ ag{ths}$	В	ond ang	les
WIOI	туре	Chain	rtes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	LYR	В	200	1	27,29,30	1.39	4 (14%)	30,37,39	1.87	7 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	LYR	A	200	1	-	2/22/40/42	0/1/1/1
1	LYR	В	200	1	-	2/22/40/42	0/1/1/1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	Ideal(Å)
1	В	200	LYR	C7-C80	3.80	1.40	1.35
1	A	200	LYR	C7-C80	3.02	1.39	1.35
1	В	200	LYR	C2-C3	2.49	1.40	1.33
1	В	200	LYR	C12-C11	2.42	1.38	1.34
1	A	200	LYR	C12-C11	2.40	1.38	1.34

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	200	LYR	C1-NZ-CE	5.34	121.80	113.33
1	A	200	LYR	C1-NZ-CE	4.77	120.88	113.33
1	A	200	LYR	C13-C12-C11	-3.69	120.38	124.53
1	В	200	LYR	C10-C9-C80	-3.39	121.11	126.23
1	A	200	LYR	C10-C9-C80	-3.31	121.24	126.23

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	200	LYR	C2-C1-NZ-CE
1	В	200	LYR	CD-CE-NZ-C1
1	A	200	LYR	CD-CE-NZ-C1
1	В	200	LYR	C2-C1-NZ-CE

There are no ring outliers.

2 monomers are involved in 13 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	200	LYR	6	0
1	В	200	LYR	7	0

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trino	Chain	Res	Link	Во	Bond lengths			ond ang	les
IVIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	LFA	A	301	-	9,9,19	0.30	0	8,8,18	0.40	0
2	LFA	В	301	-	9,9,19	0.30	0	8,8,18	0.59	0
2	LFA	В	307	-	7,7,19	0.39	0	6,6,18	0.49	0
2	LFA	A	310	-	9,9,19	0.33	0	8,8,18	0.59	0
2	LFA	A	306	-	15,15,19	0.35	0	14,14,18	0.51	0
2	LFA	В	306	-	15,15,19	0.29	0	14,14,18	0.52	0
2	LFA	A	302	-	15,15,19	0.27	0	14,14,18	0.68	0
2	LFA	A	305	-	15,15,19	0.35	0	14,14,18	0.68	0
2	LFA	A	311	-	9,9,19	0.37	0	8,8,18	0.51	0
2	LFA	В	302	-	15,15,19	0.43	0	14,14,18	0.44	0
2	LFA	A	304	-	9,9,19	0.35	0	8,8,18	0.54	0
2	LFA	В	304	-	4,4,19	0.39	0	3,3,18	0.39	0
2	LFA	В	309	-	15,15,19	0.43	0	14,14,18	0.45	0
2	LFA	A	312	-	5,5,19	0.39	0	4,4,18	0.32	0
2	LFA	A	303	-	9,9,19	0.29	0	8,8,18	0.52	0
2	LFA	A	308	ı	7,7,19	0.38	0	6,6,18	0.41	0
2	LFA	В	312	-	5,5,19	0.36	0	4,4,18	0.42	0
3	OLB	A	313	-	14,14,24	1.14	1 (7%)	15,15,25	0.99	1 (6%)
4	OLC	В	313	-	9,9,24	1.47	1 (11%)	10,10,25	1.45	1 (10%)
2	LFA	В	305	-	8,8,19	0.31	0	7,7,18	0.47	0
2	LFA	В	310	-	9,9,19	0.38	0	8,8,18	0.45	0
2	LFA	В	311	-	9,9,19	0.31	0	8,8,18	0.66	0



Mol	Mol Type Chain Res		Link	Bond lengths			В	ond ang	les	
Mol Type C	Cham	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
2	LFA	A	307	-	7,7,19	0.36	0	6,6,18	0.46	0
2	LFA	В	308	-	11,11,19	0.38	0	10,10,18	0.44	0
2	LFA	В	303	-	13,13,19	0.27	0	12,12,18	0.66	0
2	LFA	A	309	-	5,5,19	0.30	0	4,4,18	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	LFA	A	301	-	-	4/7/7/17	-
2	LFA	В	301	-	-	2/7/7/17	-
2	LFA	В	307	-	-	2/5/5/17	-
2	LFA	A	310	-	-	4/7/7/17	-
2	LFA	A	306	-	-	8/13/13/17	-
2	LFA	В	306	-	-	7/13/13/17	_
2	LFA	A	302	-	-	8/13/13/17	-
2	LFA	A	305	-	-	7/13/13/17	-
2	LFA	A	311	-	-	4/7/7/17	-
2	LFA	В	302	-	-	7/13/13/17	-
2	LFA	A	304	_	-	4/7/7/17	-
2	LFA	В	304	_	-	0/2/2/17	-
2	LFA	В	309	-	-	8/13/13/17	-
2	LFA	A	312	-	-	1/3/3/17	-
2	LFA	A	303	_	-	3/7/7/17	_
2	LFA	A	308	-	-	2/5/5/17	_
2	LFA	В	312	-	-	1/3/3/17	-
3	OLB	A	313	-	-	7/14/14/24	-
4	OLC	В	313	-	-	4/9/9/24	-
2	LFA	В	305	-	-	5/6/6/17	-
2	LFA	В	310	-	-	3/7/7/17	-
2	LFA	В	311	-	-	3/7/7/17	-
2	LFA	A	307	-	-	2/5/5/17	-
2	LFA	В	308	-	-	4/9/9/17	-
2	LFA	В	303	-	-	6/11/11/17	-
2	LFA	A	309	-	-	0/3/3/17	-



All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
4	В	313	OLC	O20-C1	4.25	1.45	1.33
3	A	313	OLB	O20-C1	4.10	1.45	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
4	В	313	OLC	O20-C1-C2	3.77	121.28	111.38
3	A	313	OLB	O20-C1-C2	2.47	119.66	111.91

There are no chirality outliers.

5 of 106 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	313	OLB	O20-C21-C22-O23
3	A	313	OLB	O20-C21-C22-C24
2	В	301	LFA	C2-C3-C4-C5
4	В	313	OLC	O20-C21-C22-O23
3	A	313	OLB	C1-C2-C3-C4

There are no ring outliers.


13 monomers are involved in 29 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	310	LFA	1	0
2	A	306	LFA	3	0
2	В	306	LFA	4	0
2	A	305	LFA	3	0
2	A	311	LFA	2	0
2	В	302	LFA	1	0
2	A	304	LFA	7	0
2	В	309	LFA	1	0
2	A	312	LFA	2	0
3	A	313	OLB	2	0
4	В	313	OLC	4	0
2	В	303	LFA	2	0
2	A	309	LFA	7	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is



within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.










Ligand L	FA A 311
~~~	~~~
Bond lengths	Bond angles
<b>✓</b>	
Torsions	Rings
Ligand L	FA B 302
~~~~	~~~~
Bond lengths	Bond angles
~~~~	
Torsions	Rings
Ligand L	FA A 304
~~~	~~~
Bond lengths	Bond angles
~~~	
Torsions	Rings
Ligand L	FA B 309
~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	~~~~
Bond lengths	Bond angles
~~~~	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
Torsions	Rings



Ligand L	FA A 312
Bond lengths	Bond angles
Torsions	Rings
Ligand L	FA A 303
~~~	~~~
Bond lengths	Bond angles
<b>\\\\</b>	
Torsions	Rings
Ligand L	FA A 308
<b>✓</b>	<b>✓</b>
Bond lengths	Bond angles
Torsions	Rings









Torsions

Rings



Ligand LFA B 305						
<b>\\\</b>						
Bond lengths	Bond angles					
Torsions	Rings					





Ligand L	FA B 311
<b>~~~</b>	
Bond lengths	Bond angles
Torsions	Rings
Ligand L	FA A 307
<b>✓</b>	
Bond lengths	Bond angles
Torsions	Rings
Ligand L	FA B 308
~~~~	~~~
Bond lengths	Bond angles
~~~	
Torsions	Rings
Ligand L	FA B 303
	~~~~
Bond lengths	Bond angles
~~~~	\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\
Torsions	Rings



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q<0.9
1	A	216/220 (98%)	0.20	14 (6%) 18 21	14, 27, 52, 71	4 (1%)
1	В	$216/220 \ (98\%)$	0.25	14 (6%) 18 21	14, 27, 56, 69	3 (1%)
All	All	432/440 (98%)	0.23	28 (6%) 18 21	14, 27, 53, 71	7 (1%)

The worst 5 of 28 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	181	VAL	8.4
1	A	181	VAL	7.3
1	В	184	PHE	6.8
1	A	1[A]	MET	5.8
1	A	184	PHE	5.1

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	LYR	A	200	29/30	0.95	0.09	14,16,19,19	0
1	LYR	В	200	29/30	0.95	0.10	14,17,19,20	0

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}({ ext{\AA}}^2)$	Q < 0.9
2	LFA	A	307	8/20	0.55	0.28	49,51,52,53	0
2	LFA	A	312	6/20	0.67	0.16	49,56,57,59	0
2	LFA	A	311	10/20	0.70	0.20	42,61,78,78	0
2	LFA	В	312	6/20	0.70	0.18	55,56,58,62	0
3	OLB	A	313	15/25	0.74	0.17	53,61,75,75	0
2	LFA	В	307	8/20	0.76	0.24	45,50,56,59	0
2	LFA	В	310	10/20	0.77	0.17	50,55,68,68	0
2	LFA	A	308	8/20	0.77	0.19	36,45,50,50	0
2	LFA	В	309	16/20	0.77	0.19	37,50,54,57	0
2	LFA	A	306	16/20	0.78	0.20	38,46,70,70	0
2	LFA	В	302	16/20	0.79	0.19	31,46,55,56	0
2	LFA	A	305	16/20	0.79	0.18	33,38,50,51	0
2	LFA	В	303	14/20	0.81	0.23	41,44,51,52	0
2	LFA	A	302	16/20	0.81	0.21	42,46,57,59	0
2	LFA	В	305	9/20	0.83	0.19	34,40,44,47	0
2	LFA	В	308	12/20	0.83	0.20	39,48,56,57	0
2	LFA	В	311	10/20	0.84	0.20	36,55,58,58	0
2	LFA	A	310	10/20	0.84	0.14	41,47,60,61	0
2	LFA	A	304	10/20	0.84	0.18	41,43,48,48	0
4	OLC	В	313	10/25	0.84	0.16	27,42,47,62	0
2	LFA	A	303	10/20	0.86	0.17	37,43,48,48	0
2	LFA	В	306	16/20	0.87	0.15	36,39,46,49	0
2	LFA	A	301	10/20	0.88	0.16	37,38,46,50	0
2	LFA	A	309	6/20	0.89	0.16	36,37,42,44	0
2	LFA	В	304	5/20	0.91	0.18	38,41,43,46	0
2	LFA	В	301	10/20	0.92	0.15	35,38,41,51	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around LFA A 307:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA A 312:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around LFA A 311:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA B 312:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around OLB A 313:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA B 307:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around LFA B 310:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA A 308:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around LFA B 309:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA A 306:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around LFA B 302:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA A 305:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around LFA B 303:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA A 302:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











Electron density around LFA B 305: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around LFA B 308: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around LFA B 311:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA A 310:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around LFA A 304:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around OLC B 313:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around LFA A 303:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA B 306:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around LFA A 301:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around LFA B 301:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

