



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 17, 2023 – 09:14 am GMT

PDB ID : 4B1M
Title : CARBOHYDRATE BINDING MODULE CBM66 FROM BACILLUS SUBTILIS
Authors : Cuskin, F.; Flint, J.E.; Morland, C.; Basle, A.; Henrissat, B.; Countinho, P.M.; Strazzulli, A.; Solzehinkin, A.; Davies, G.J.; Gilbert, H.J.; Gloster, T.M.
Deposited on : 2012-07-11
Resolution : 1.10 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1619 (1.14-1.06)
Clashscore	141614	1671 (1.14-1.06)
Ramachandran outliers	138981	1615 (1.14-1.06)
Sidechain outliers	138945	1613 (1.14-1.06)
RSRZ outliers	127900	1588 (1.14-1.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	185	 3% 82% 7% 11%
1	B	185	 3% 83% 5% 12%
1	C	185	 3% 77% 9% 13%
2	D	2	 50% 50%
2	E	2	 100%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	2	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 4469 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LEVANASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	165	1315	829	216	264	6	0	6	0
1	B	163	1299	818	212	265	4	0	7	0
1	C	161	1308	835	211	259	3	0	11	0

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	493	MET	-	expression tag	UNP P05656
A	494	GLY	-	expression tag	UNP P05656
A	495	HIS	-	expression tag	UNP P05656
A	496	HIS	-	expression tag	UNP P05656
A	497	HIS	-	expression tag	UNP P05656
A	498	HIS	-	expression tag	UNP P05656
A	499	HIS	-	expression tag	UNP P05656
A	500	HIS	-	expression tag	UNP P05656
A	501	HIS	-	expression tag	UNP P05656
A	502	HIS	-	expression tag	UNP P05656
A	503	HIS	-	expression tag	UNP P05656
A	504	HIS	-	expression tag	UNP P05656
A	505	SER	-	expression tag	UNP P05656
A	506	SER	-	expression tag	UNP P05656
A	507	GLY	-	expression tag	UNP P05656
A	508	HIS	-	expression tag	UNP P05656
A	509	ILE	-	expression tag	UNP P05656
A	510	GLU	-	expression tag	UNP P05656
A	511	GLY	-	expression tag	UNP P05656
A	512	ARG	-	expression tag	UNP P05656
A	513	HIS	-	expression tag	UNP P05656
A	514	MET	-	expression tag	UNP P05656
B	493	MET	-	expression tag	UNP P05656

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	494	GLY	-	expression tag	UNP P05656
B	495	HIS	-	expression tag	UNP P05656
B	496	HIS	-	expression tag	UNP P05656
B	497	HIS	-	expression tag	UNP P05656
B	498	HIS	-	expression tag	UNP P05656
B	499	HIS	-	expression tag	UNP P05656
B	500	HIS	-	expression tag	UNP P05656
B	501	HIS	-	expression tag	UNP P05656
B	502	HIS	-	expression tag	UNP P05656
B	503	HIS	-	expression tag	UNP P05656
B	504	HIS	-	expression tag	UNP P05656
B	505	SER	-	expression tag	UNP P05656
B	506	SER	-	expression tag	UNP P05656
B	507	GLY	-	expression tag	UNP P05656
B	508	HIS	-	expression tag	UNP P05656
B	509	ILE	-	expression tag	UNP P05656
B	510	GLU	-	expression tag	UNP P05656
B	511	GLY	-	expression tag	UNP P05656
B	512	ARG	-	expression tag	UNP P05656
B	513	HIS	-	expression tag	UNP P05656
B	514	MET	-	expression tag	UNP P05656
C	493	MET	-	expression tag	UNP P05656
C	494	GLY	-	expression tag	UNP P05656
C	495	HIS	-	expression tag	UNP P05656
C	496	HIS	-	expression tag	UNP P05656
C	497	HIS	-	expression tag	UNP P05656
C	498	HIS	-	expression tag	UNP P05656
C	499	HIS	-	expression tag	UNP P05656
C	500	HIS	-	expression tag	UNP P05656
C	501	HIS	-	expression tag	UNP P05656
C	502	HIS	-	expression tag	UNP P05656
C	503	HIS	-	expression tag	UNP P05656
C	504	HIS	-	expression tag	UNP P05656
C	505	SER	-	expression tag	UNP P05656
C	506	SER	-	expression tag	UNP P05656
C	507	GLY	-	expression tag	UNP P05656
C	508	HIS	-	expression tag	UNP P05656
C	509	ILE	-	expression tag	UNP P05656
C	510	GLU	-	expression tag	UNP P05656
C	511	GLY	-	expression tag	UNP P05656
C	512	ARG	-	expression tag	UNP P05656
C	513	HIS	-	expression tag	UNP P05656

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	514	MET	-	expression tag	UNP P05656

- Molecule 2 is an oligosaccharide called beta-D-fructofuranose-(2-6)-beta-D-fructofuranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
2	D	2	23	12	11	0	0	0
2	E	2	23	12	11	0	0	0
2	F	2	23	12	11	0	0	0

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Na	0	0
			2	2		
3	B	1	Total	Na	0	0
			1	1		
3	C	1	Total	Na	0	0
			1	1		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	177	Total	O	0	0
			177	177		
5	B	139	Total	O	0	0
			139	139		
5	C	153	Total	O	0	0
			153	153		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: LEVANASE



- Molecule 1: LEVANASE



- Molecule 1: LEVANASE



- Molecule 2: beta-D-fructofuranose-(2-6)-beta-D-fructofuranose



- Molecule 2: beta-D-fructofuranose-(2-6)-beta-D-fructofuranose



- Molecule 2: beta-D-fructofuranose-(2-6)-beta-D-fructofuranose



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	66.90Å 36.91Å 87.36Å 90.00° 90.61° 90.00°	Depositor
Resolution (Å)	87.35 – 1.10 33.45 – 1.10	Depositor EDS
% Data completeness (in resolution range)	91.1 (87.35-1.10) 91.1 (33.45-1.10)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.83 (at 1.10Å)	Xtrriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.156 , 0.181 0.163 , 0.189	Depositor DCC
R_{free} test set	7887 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	7.8	Xtrriage
Anisotropy	1.187	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.018 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4469	wwPDB-VP
Average B, all atoms (Å ²)	12.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, FRU, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.70	0/1360	0.80	1/1832 (0.1%)
1	B	0.67	0/1346	0.81	1/1816 (0.1%)
1	C	0.66	0/1374	0.83	4/1854 (0.2%)
All	All	0.68	0/4080	0.81	6/5502 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	522	MET	CG-SD-CE	-9.17	85.53	100.20
1	A	636	ARG	NE-CZ-NH2	-6.66	116.97	120.30
1	C	636	ARG	NE-CZ-NH2	-5.54	117.53	120.30
1	B	635	ASP	CB-CG-OD1	5.30	123.08	118.30
1	C	617[A]	LYS	CD-CE-NZ	-5.17	99.82	111.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	599	LEU	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1315	0	1260	7	0
1	B	1299	0	1245	6	0
1	C	1308	0	1294	13	0
2	D	23	0	22	0	0
2	E	23	0	22	0	0
2	F	23	0	20	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
4	B	5	0	0	0	0
5	A	177	0	0	7	0
5	B	139	0	0	2	1
5	C	153	0	0	2	0
All	All	4469	0	3863	26	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 26 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:2044:HOH:O	1:B:635:ASP:OD2	1.87	0.90
1:A:556:SER:OG	5:A:2068:HOH:O	1.84	0.89
1:A:583:LYS:HG2	5:A:2100:HOH:O	1.76	0.84
1:A:527:THR:HG21	5:A:2028:HOH:O	1.88	0.73
5:A:2044:HOH:O	1:B:652[B]:SER:OG	2.02	0.73

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:B:2010:HOH:O	5:B:2018:HOH:O[2_556]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	169/185 (91%)	165 (98%)	4 (2%)	0	100	100
1	B	168/185 (91%)	165 (98%)	3 (2%)	0	100	100
1	C	172/185 (93%)	169 (98%)	3 (2%)	0	100	100
All	All	509/555 (92%)	499 (98%)	10 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	142/153 (93%)	139 (98%)	3 (2%)	53	14
1	B	141/153 (92%)	139 (99%)	2 (1%)	67	30
1	C	145/153 (95%)	143 (99%)	2 (1%)	67	30
All	All	428/459 (93%)	421 (98%)	7 (2%)	60	25

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	662	ASN
1	B	675	LYS
1	C	675	LYS
1	C	662	ASN
1	A	675	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

6 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FRU	D	1	2	11,12,12	1.25	1 (9%)	10,18,18	1.15	1 (10%)
2	FRU	D	2	2	11,11,12	0.67	0	15,15,18	0.62	0
2	FRU	E	1	2	11,12,12	1.51	1 (9%)	10,18,18	2.38	4 (40%)
2	FRU	E	2	2	11,11,12	0.75	0	15,15,18	0.98	1 (6%)
2	FRU	F	1	2	11,12,12	1.54	1 (9%)	10,18,18	3.05	2 (20%)
2	FRU	F	2	2	11,11,12	0.55	0	15,15,18	0.73	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FRU	D	1	2	-	3/5/24/24	0/1/1/1
2	FRU	D	2	2	-	0/4/20/24	0/1/1/1
2	FRU	E	1	2	-	3/5/24/24	0/1/1/1
2	FRU	E	2	2	-	0/4/20/24	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FRU	F	1	2	-	1/5/24/24	0/1/1/1
2	FRU	F	2	2	-	0/4/20/24	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	1	FRU	O5-C2	-4.20	1.36	1.43
2	D	1	FRU	O2-C2	3.34	1.46	1.40
2	F	1	FRU	O4-C4	-3.13	1.35	1.43

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1	FRU	O1-C1-C2	-8.70	93.37	111.86
2	E	1	FRU	C6-C5-C4	4.29	125.43	115.09
2	E	1	FRU	O3-C3-C4	-4.26	98.59	113.32
2	F	1	FRU	O2-C2-O5	-3.56	102.63	109.50
2	E	1	FRU	O6-C6-C5	3.39	122.94	111.29

There are no chirality outliers.



5 of 7 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	D	1	FRU	O1-C1-C2-C3
2	D	1	FRU	O1-C1-C2-O2
2	D	1	FRU	O1-C1-C2-O5
2	E	1	FRU	O1-C1-C2-C3
2	E	1	FRU	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	B	1681	-	4,4,4	0.26	0	6,6,6	0.66	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	165/185 (89%)	0.06	2 (1%) 79 76	7, 9, 15, 22	0
1	B	163/185 (88%)	0.14	6 (3%) 41 38	7, 10, 19, 31	0
1	C	161/185 (87%)	0.14	5 (3%) 49 46	7, 10, 18, 31	0
All	All	489/555 (88%)	0.11	13 (2%) 54 51	7, 10, 18, 31	0

The worst 5 of 13 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	569	GLY	8.0
1	B	570	ASN	4.9
1	C	570	ASN	4.6
1	B	571	GLY	3.6
1	B	568	ASP	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FRU	F	1	12/12	0.90	0.14	10,23,34,37	0
2	FRU	E	1	12/12	0.92	0.19	13,25,41,48	0
2	FRU	D	1	12/12	0.92	0.18	12,27,37,43	0


Continued on next page...


Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	FRU	E	2	11/12	0.98	0.06	8,9,12,13	0
2	FRU	D	2	11/12	0.98	0.08	7,8,11,12	0
2	FRU	F	2	11/12	0.98	0.08	7,8,9,10	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.







6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	B	1681	5/5	0.96	0.23	20,22,27,29	0
3	NA	B	1680	1/1	0.99	0.06	14,14,14,14	0
3	NA	C	1680	1/1	0.99	0.04	11,11,11,11	0
3	NA	A	1681	1/1	0.99	0.28	21,21,21,21	0
3	NA	A	1680	1/1	1.00	0.04	9,9,9,9	0

6.5 Other polymers [i](#)

There are no such residues in this entry.