



wwPDB X-ray Structure Validation Summary Report ⓘ

May 16, 2020 – 01:40 am BST

PDB ID : 4B1Z
Title : Structure of the Phactr1 RPEL domain bound to G-actin
Authors : Mouilleron, S.; Wiezlak, M.; O'Reilly, N.; Treisman, R.; McDonald, N.Q.
Deposited on : 2012-07-12
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1149 (3.34-3.26)
Clashscore	141614	1205 (3.34-3.26)
Ramachandran outliers	138981	1183 (3.34-3.26)
Sidechain outliers	138945	1182 (3.34-3.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	376	80% 10% 11%
1	B	376	85% 10% 5%
1	C	376	88% 10% .
1	D	376	84% 11% 5%
1	E	376	84% 12% 5%
1	F	376	82% 13% 5%
2	M	115	74% 18% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	N	115	 78% 12% 7%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 17941 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called ACTIN, ALPHA SKELETAL MUSCLE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	335	Total 2495	C 1588	N 415	O 474	S 18	0	0	0
1	B	357	Total 2711	C 1721	N 448	O 523	S 19	0	0	0
1	C	370	Total 2815	C 1787	N 476	O 532	S 20	0	0	0
1	D	356	Total 2648	C 1675	N 446	O 508	S 19	0	0	0
1	E	359	Total 2682	C 1701	N 450	O 512	S 19	0	0	0
1	F	357	Total 2682	C 1706	N 444	O 513	S 19	0	0	0

- Molecule 2 is a protein called PHOSPHATASE AND ACTIN REGULATOR 1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	M	110	Total 846	C 517	N 166	O 163	0	0	0
2	N	107	Total 821	C 505	N 159	O 157	0	0	0

- Molecule 3 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula: $C_{10}H_{16}N_5O_{13}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	Total 31	10	5	13	3	0	0
3	B	1	Total 31	10	5	13	3	0	0
3	C	1	Total 31	10	5	13	3	0	0
3	D	1	Total 31	10	5	13	3	0	0
3	E	1	Total 31	10	5	13	3	0	0
3	F	1	Total 31	10	5	13	3	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
4	D	1	Total 1	1	0	0
4	E	1	Total 1	1	0	0
4	B	1	Total 1	1	0	0
4	C	1	Total 1	1	0	0
4	A	1	Total 1	1	0	0
4	F	1	Total 1	1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0
5	C	1	Total C O 6 3 3	0	0
5	E	1	Total C O 6 3 3	0	0
5	F	1	Total C O 6 3 3	0	0
5	N	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	E	1	Total O 1 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ACTIN, ALPHA SKELETAL MUSCLE




- Molecule 1: ACTIN, ALPHA SKELETAL MUSCLE



- Molecule 1: ACTIN, ALPHA SKELETAL MUSCLE





- Molecule 1: ACTIN, ALPHA SKELETAL MUSCLE







- Molecule 1: ACTIN, ALPHA SKELETAL MUSCLE




- Molecule 1: ACTIN, ALPHA SKELETAL MUSCLE



- Molecule 2: PHOSPHATASE AND ACTIN REGULATOR 1



- Molecule 2: PHOSPHATASE AND ACTIN REGULATOR 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	127.06Å 142.89Å 184.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	74.62 – 3.30 74.62 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.4 (74.62-3.30) 99.4 (74.62-3.30)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.34 (at 3.33Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE: 1.8.1_1168)	Depositor
R, R_{free}	0.213 , 0.236 0.565 , 0.511	Depositor DCC
R_{free} test set	2580 reflections (2.53%)	wwPDB-VP
Wilson B-factor (Å ²)	70.7	Xtrriage
Anisotropy	0.685	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 55.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.28$, $\langle L^2 \rangle = 0.12$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.50	EDS
Total number of atoms	17941	wwPDB-VP
Average B, all atoms (Å ²)	87.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 1.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/2552	0.47	0/3480
1	B	0.24	0/2772	0.46	0/3774
1	C	0.24	0/2878	0.47	0/3911
1	D	0.23	0/2706	0.47	0/3685
1	E	0.24	0/2742	0.47	0/3738
1	F	0.25	0/2742	0.48	0/3736
2	M	0.21	0/851	0.43	0/1148
2	N	0.21	0/827	0.41	0/1115
All	All	0.24	0/18070	0.47	0/24587

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2495	0	2353	19	0
1	B	2711	0	2582	26	0
1	C	2815	0	2724	27	0
1	D	2648	0	2476	26	0
1	E	2682	0	2525	34	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2682	0	2550	33	0
2	M	846	0	814	13	0
2	N	821	0	787	10	0
3	A	31	0	12	1	0
3	B	31	0	12	1	0
3	C	31	0	12	0	0
3	D	31	0	12	2	0
3	E	31	0	12	1	0
3	F	31	0	12	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
4	E	1	0	0	0	0
4	F	1	0	0	0	0
5	A	12	0	16	0	0
5	B	6	0	8	0	0
5	C	12	0	16	0	0
5	E	6	0	8	1	0
5	F	6	0	8	0	0
5	N	6	0	8	0	0
6	E	1	0	0	0	0
All	All	17941	0	16947	167	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 167 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:146:GLY:O	2:N:465:ARG:NH1	2.04	0.89
1:F:371:HIS:HB3	2:N:527:GLU:HG3	1.58	0.84
1:D:62:ARG:HB2	1:D:67:LEU:HD11	1.63	0.80
1:C:41:GLN:O	1:E:28:ARG:NH2	2.14	0.80
1:C:39:ARG:HG2	1:C:66:THR:HG23	1.67	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/376 (88%)	326 (98%)	5 (2%)	0	100	100
1	B	353/376 (94%)	347 (98%)	6 (2%)	0	100	100
1	C	368/376 (98%)	359 (98%)	8 (2%)	1 (0%)	41	71
1	D	352/376 (94%)	346 (98%)	6 (2%)	0	100	100
1	E	355/376 (94%)	348 (98%)	7 (2%)	0	100	100
1	F	353/376 (94%)	346 (98%)	7 (2%)	0	100	100
2	M	108/115 (94%)	105 (97%)	3 (3%)	0	100	100
2	N	105/115 (91%)	102 (97%)	3 (3%)	0	100	100
All	All	2325/2486 (94%)	2279 (98%)	45 (2%)	1 (0%)	100	100

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	52	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/319 (79%)	250 (100%)	1 (0%)	91	95
1	B	281/319 (88%)	281 (100%)	0	100	100
1	C	292/319 (92%)	291 (100%)	1 (0%)	92	96
1	D	264/319 (83%)	262 (99%)	2 (1%)	81	89

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	270/319 (85%)	268 (99%)	2 (1%)	84	90
1	F	274/319 (86%)	273 (100%)	1 (0%)	91	95
2	M	83/111 (75%)	73 (88%)	10 (12%)	5	20
2	N	80/111 (72%)	71 (89%)	9 (11%)	6	22
All	All	1795/2136 (84%)	1769 (99%)	26 (1%)	67	82

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	M	448	ASP
2	M	485	ASN
2	N	462	LEU
2	M	468	GLN
2	M	481	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	88	HIS
1	B	371	HIS
1	E	371	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 6 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	GOL	A	1379	-	5,5,5	0.37	0	5,5,5	0.28	0
3	ATP	C	1376	4	26,33,33	0.92	1 (3%)	31,52,52	1.43	5 (16%)
5	GOL	B	1378	-	5,5,5	0.36	0	5,5,5	0.22	0
5	GOL	C	1378	-	5,5,5	0.38	0	5,5,5	0.30	0
5	GOL	F	1378	-	5,5,5	0.38	0	5,5,5	0.22	0
5	GOL	A	1378	-	5,5,5	0.38	0	5,5,5	0.21	0
5	GOL	E	1378	-	5,5,5	0.36	0	5,5,5	0.31	0
5	GOL	N	1529	-	5,5,5	0.37	0	5,5,5	0.29	0
3	ATP	A	1376	4	26,33,33	0.92	1 (3%)	31,52,52	1.44	5 (16%)
3	ATP	B	1376	4	26,33,33	0.90	1 (3%)	31,52,52	1.48	5 (16%)
3	ATP	E	1376	4	26,33,33	0.93	1 (3%)	31,52,52	1.38	5 (16%)
3	ATP	F	1376	4	26,33,33	0.91	1 (3%)	31,52,52	1.40	5 (16%)
3	ATP	D	1376	4	26,33,33	0.93	1 (3%)	31,52,52	1.40	5 (16%)
5	GOL	C	1379	-	5,5,5	0.37	0	5,5,5	0.29	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	A	1379	-	-	0/4/4/4	-
3	ATP	C	1376	4	-	0/18/38/38	0/3/3/3
5	GOL	B	1378	-	-	2/4/4/4	-
5	GOL	C	1378	-	-	2/4/4/4	-
5	GOL	F	1378	-	-	2/4/4/4	-
5	GOL	A	1378	-	-	2/4/4/4	-
5	GOL	E	1378	-	-	2/4/4/4	-
5	GOL	N	1529	-	-	2/4/4/4	-
3	ATP	A	1376	4	-	0/18/38/38	0/3/3/3
3	ATP	B	1376	4	-	2/18/38/38	0/3/3/3
3	ATP	E	1376	4	-	4/18/38/38	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ATP	F	1376	4	-	0/18/38/38	0/3/3/3
3	ATP	D	1376	4	-	0/18/38/38	0/3/3/3
5	GOL	C	1379	-	-	4/4/4/4	-

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1376	ATP	C5-C4	2.50	1.47	1.40
3	D	1376	ATP	C5-C4	2.48	1.47	1.40
3	C	1376	ATP	C5-C4	2.48	1.47	1.40
3	A	1376	ATP	C5-C4	2.46	1.47	1.40
3	F	1376	ATP	C5-C4	2.45	1.47	1.40

The worst 5 of 30 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	F	1376	ATP	N3-C2-N1	-3.30	123.52	128.68
3	B	1376	ATP	N3-C2-N1	-3.26	123.58	128.68
3	C	1376	ATP	N3-C2-N1	-3.23	123.63	128.68
3	B	1376	ATP	PA-O3A-PB	-3.17	121.94	132.83
3	A	1376	ATP	N3-C2-N1	-3.16	123.73	128.68

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	N	1529	GOL	C1-C2-C3-O3
5	B	1378	GOL	C1-C2-C3-O3
5	C	1378	GOL	C1-C2-C3-O3
5	F	1378	GOL	C1-C2-C3-O3
5	A	1378	GOL	C1-C2-C3-O3

There are no ring outliers.

6 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	E	1378	GOL	1	0
3	A	1376	ATP	1	0
3	B	1376	ATP	1	0
3	E	1376	ATP	1	0
3	F	1376	ATP	1	0


Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	1376	ATP	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [\(i\)](#)

6.1 Protein, DNA and RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates [\(i\)](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [\(i\)](#)




Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around ATP A 1376:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ATP B 1376:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ATP E 1376:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ATP F 1376:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

Unable to reproduce the depositor's R factor - this section is therefore empty.