

wwPDB X-ray Structure Validation Summary Report (i)

Oct 30, 2023 – 01:08 PM JST

PDB ID : 5B1I

Title: Crystal structure of K42A mutant of cystathionine beta-synthase from Lacto-

bacillus plantarum in a complex with L-methionine

Authors: Matoba, Y.; Sugiyama, M.

Deposited on : 2015-12-04

Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.orgA user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar \ resolution} \\ (\#{\rm Entries, \ resolution \ range(\AA)}) \end{array}$		
R_{free}	130704	1149 (3.34-3.26)		
Clashscore	141614	1205 (3.34-3.26)		
Ramachandran outliers	138981	1183 (3.34-3.26)		
Sidechain outliers	138945	1182 (3.34-3.26)		
RSRZ outliers	127900	1115 (3.34-3.26)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	311	55%	37%	5% •			
1	В	311	60%	31%	7% •			
1	С	311	59%	33%	5% •			
1	D	311	53%	38%	6% •			
1	Е	311	53%	37%	7% •			
1	F	311	57%	35%	5% •			





Continued from previous page...

Mol	Chain	Length	Quality of chain				
1	G	311	53%	39%	5% •		
1	Н	311	34%	42%	6% •		



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 18392 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Cystathionine beta-synthase.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	303	Total	С	N	О	S	0	0	0
1		303	2265	1438	394	426	7	0	0	
1	В	303	Total	С	N	О	S	0	0	0
1	Б	303	2265	1438	394	426	7	0	0	
1	С	303	Total	С	N	О	S	0	0	0
1		303	2265	1438	394	426	7	U	0	
1	D	303	Total	С	N	О	S	0	0	0
1	D	303	2265	1438	394	426	7			
1	Е	303	Total	С	N	О	S	0	0	0
1	12	303	2265	1438	394	426	7	0		
1	F	303	Total	С	N	О	S	0	0	0
1	Г	303	2265	1438	394	426	7	0	U	
1	G	303	Total	С	Ν	Ο	S	0	0	0
1	ı G	505	2265	1438	394	426	7	0	0	
1	Н	303	Total	С	N	О	S	0	0	0
1	1 H	303	2265	1438	394	426	7		U	

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	42	ALA	LYS	engineered mutation	UNP F9UT54
A	304	LEU	-	expression tag	UNP F9UT54
A	305	GLU	-	expression tag	UNP F9UT54
A	306	HIS	-	expression tag	UNP F9UT54
A	307	HIS	-	expression tag	UNP F9UT54
A	308	HIS	-	expression tag	UNP F9UT54
A	309	HIS	-	expression tag	UNP F9UT54
A	310	HIS	-	expression tag	UNP F9UT54
A	311	HIS	-	expression tag	UNP F9UT54
В	42	ALA	LYS	engineered mutation	UNP F9UT54
В	304	LEU	-	expression tag	UNP F9UT54
В	305	GLU	-	expression tag	UNP F9UT54
В	306	HIS	-	expression tag	UNP F9UT54



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled	Actual	Comment	Reference
В	307	HIS	-	expression tag	UNP F9UT54
В	308	HIS	-	expression tag	UNP F9UT54
В	309	HIS	-	expression tag	UNP F9UT54
В	310	HIS	-	expression tag	UNP F9UT54
В	311	HIS	-	expression tag	UNP F9UT54
С	42	ALA	LYS	engineered mutation	UNP F9UT54
С	304	LEU	-	expression tag	UNP F9UT54
С	305	GLU	-	expression tag	UNP F9UT54
С	306	HIS	-	expression tag	UNP F9UT54
С	307	HIS	-	expression tag	UNP F9UT54
С	308	HIS	-	expression tag	UNP F9UT54
С	309	HIS	-	expression tag	UNP F9UT54
С	310	HIS	-	expression tag	UNP F9UT54
С	311	HIS	-	expression tag	UNP F9UT54
D	42	ALA	LYS	engineered mutation	UNP F9UT54
D	304	LEU	-	expression tag	UNP F9UT54
D	305	GLU	-	expression tag	UNP F9UT54
D	306	HIS	-	expression tag	UNP F9UT54
D	307	HIS	-	expression tag	UNP F9UT54
D	308	HIS	-	expression tag	UNP F9UT54
D	309	HIS	-	expression tag	UNP F9UT54
D	310	HIS	-	expression tag	UNP F9UT54
D	311	HIS	-	expression tag	UNP F9UT54
Е	42	ALA	LYS	engineered mutation	UNP F9UT54
Е	304	LEU	-	expression tag	UNP F9UT54
Е	305	GLU	-	expression tag	UNP F9UT54
Е	306	HIS	-	expression tag	UNP F9UT54
Е	307	HIS	-	expression tag	UNP F9UT54
Е	308	HIS	-	expression tag	UNP F9UT54
E	309	HIS	-	expression tag	UNP F9UT54
E	310	HIS	-	expression tag	UNP F9UT54
Е	311	HIS	-	expression tag	UNP F9UT54
F	42	ALA	LYS	engineered mutation	UNP F9UT54
F	304	LEU	-	expression tag	UNP F9UT54
F	305	GLU	-	expression tag	UNP F9UT54
F	306	HIS	-	expression tag	UNP F9UT54
F	307	HIS	-	expression tag	UNP F9UT54
F	308	HIS	-	expression tag	UNP F9UT54
F	309	HIS	-	expression tag	UNP F9UT54
F	310	HIS	-	expression tag	UNP F9UT54
F	311	HIS	-	expression tag	UNP F9UT54
G	42	ALA	LYS	engineered mutation	UNP F9UT54



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	304	LEU	-	expression tag	UNP F9UT54
G	305	GLU	-	expression tag	UNP F9UT54
G	306	HIS	-	expression tag	UNP F9UT54
G	307	HIS	-	expression tag	UNP F9UT54
G	308	HIS	-	expression tag	UNP F9UT54
G	309	HIS	-	expression tag	UNP F9UT54
G	310	HIS	-	expression tag	UNP F9UT54
G	311	HIS	-	expression tag	UNP F9UT54
Н	42	ALA	LYS	engineered mutation	UNP F9UT54
Н	304	LEU	-	expression tag	UNP F9UT54
Н	305	GLU	-	expression tag	UNP F9UT54
Н	306	HIS	-	expression tag	UNP F9UT54
Н	307	HIS	-	expression tag	UNP F9UT54
Н	308	HIS	-	expression tag	UNP F9UT54
Н	309	HIS	-	expression tag	UNP F9UT54
Н	310	HIS	-	expression tag	UNP F9UT54
Н	311	HIS	_	expression tag	UNP F9UT54

• Molecule 2 is N-[(3-HYDROXY-2-METHYL-5-{[(TRIHYDROXYPHOSPHORANYL)OXY]METHYL}PYRIDIN-4-YL)METHYLENE]METHIONINE (three-letter code: AA5) (formula: $C_{13}H_{19}N_2O_7PS$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
2	٨	1	Total	С	N	О	Р	S	0	0
	\mathcal{L} \mathcal{A}		24	13	2	7	1	1	0	
9	D	1	Total	С	N	О	Р	S	0	0
	В	B 1	24	13	2	7	1	1		. 0



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
2	С	1	Total	С	N	О	Р	S	0	0	
		1	24	13	2	7	1	1	0	0	
2	D	1	Total	С	N	О	Р	S	0	0	
	ע	1	24	13	2	7	1	1	0	0	
2	Е	1	Total	С	N	О	Р	S	0	0	
	<u> 1</u> 2	1	24	13	2	7	1	1	0		
2	F	1	Total	С	N	О	Р	S	0	0	
	I'	1	24	13	2	7	1	1	0		
2	G	1	Total	С	N	О	Р	S	0	0	
	2 G	1	24	13	2	7	1	1	U		
2	Н	1	Total	С	N	О	Р	S	0	0	
	11	1	24	13	2	7	1	1			

 \bullet Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	A	1	Total O S 5 4 1	0	0
3	В	1	Total O S 5 4 1	0	0
3	В	1	Total O S 5 4 1	0	0
3	С	1	Total O S 5 4 1	0	0



Continued from previous page...


Mol	Chain	Residues	Ato	oms		ZeroOcc	AltConf
3	С	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total 5	O 4	S 1	0	0
3	D	1	Total	О	S	0	0
	D	1	5	4	1	O	U
3	Е	1	Total	Ο	\mathbf{S}	0	0
3	l Li	1	5	4	1	0	
3	Е	1	Total	О	S	0	0
3	12	1	5	4	1	0	O
3	F	1	Total	О	S	0	0
3	Г	1	5	4	1	0	
3	F	1	Total	О	S	0	0
3	Г	1	5	4	1	0	U
3	G	1	Total	О	S	0	0
3	G	1	5	4	1	0	U
3	G	1	Total	О	S	0	0
3	G	1	5	4	1	U	U
3	Н	1	Total	О	S	0	0
	11	1	5	4	1	U	U
3	Н	1	Total	О	S	0	0
	11	1	5	4	1		U




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Cystathionine beta-synthase




• Molecule 1: Cystathionine beta-synthase




• Molecule 1: Cystathionine beta-synthase

Chain C: 59% 33% 5%.






• Molecule 1: Cystathionine beta-synthase




• Molecule 1: Cystathionine beta-synthase



• Molecule 1: Cystathionine beta-synthase











4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants	165.94Å 165.94Å 259.49Å	Donogitor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.98 - 3.30	Depositor
resolution (A)	48.84 - 3.30	EDS
% Data completeness	97.7 (29.98-3.30)	Depositor
(in resolution range)	99.8 (48.84-3.30)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.71 (at 3.33Å)	Xtriage
Refinement program	CNS 1.2	Depositor
R, R_{free}	0.238 , 0.283	Depositor
it, it free	0.235 , 0.284	DCC
R_{free} test set	2797 reflections (5.08%)	wwPDB-VP
Wilson B-factor (A^2)	43.9	Xtriage
Anisotropy	0.369	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.33, 41.4	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.87	EDS
Total number of atoms	18392	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 1.92% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, AA5

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.39	0/2314	0.62	0/3152
1	В	0.44	0/2314	0.66	0/3152
1	С	0.44	0/2314	0.64	0/3152
1	D	0.43	0/2314	0.63	0/3152
1	Е	0.41	0/2314	0.62	0/3152
1	F	0.44	0/2314	0.64	0/3152
1	G	0.38	0/2314	0.59	0/3152
1	Н	0.41	0/2314	0.60	0/3152
All	All	0.42	0/18512	0.63	0/25216

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2265	0	2275	92	0
1	В	2265	0	2275	89	0
1	С	2265	0	2275	88	0
1	D	2265	0	2275	121	1
1	Е	2265	0	2275	108	1



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	F	2265	0	2275	92	0
1	G	2265	0	2275	98	0
1	Н	2265	0	2275	130	0
2	A	24	0	15	5	0
2	В	24	0	16	5	0
2	С	24	0	16	3	0
2	D	24	0	16	2	0
2	E	24	0	16	3	0
2	F	24	0	16	2	0
2	G	24	0	16	2	0
2	Н	24	0	16	4	0
3	A	10	0	0	0	0
3	В	10	0	0	1	0
3	С	10	0	0	0	0
3	D	10	0	0	0	0
3	E	10	0	0	0	0
3	F	10	0	0	0	0
3	G	10	0	0	0	0
3	Н	10	0	0	0	0
All	All	18392	0	18327	751	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 751 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:148:ASN:HB3	1:A:149:PRO:HD3	1.46	0.97
1:H:148:ASN:HB3	1:H:149:PRO:HD3	1.45	0.94
1:B:45:LEU:HD22	1:B:49:LEU:HG	1.51	0.93
1:A:25:HIS:NE2	1:C:230:ASP:HB2	1.83	0.93
1:G:45:LEU:HD22	1:G:49:LEU:HG	1.48	0.92

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	Clash overlap (Å)
1:D:210:GLY:N	1:E:58:ARG:NH2[1_455]	2.11	0.09



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	301/311 (97%)	271 (90%)	26 (9%)	4 (1%)	12	40
1	В	301/311 (97%)	276 (92%)	23 (8%)	2 (1%)	22	54
1	С	301/311 (97%)	278 (92%)	22 (7%)	1 (0%)	41	71
1	D	301/311 (97%)	276 (92%)	25 (8%)	0	100	100
1	E	301/311 (97%)	274 (91%)	24 (8%)	3 (1%)	15	46
1	F	301/311 (97%)	275 (91%)	25 (8%)	1 (0%)	41	71
1	G	301/311 (97%)	271 (90%)	28 (9%)	2 (1%)	22	54
1	Н	301/311 (97%)	276 (92%)	24 (8%)	1 (0%)	41	71
All	All	2408/2488 (97%)	2197 (91%)	197 (8%)	14 (1%)	25	57

5 of 14 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Е	57	GLY
1	В	192	ASP
1	Е	97	SER
1	Е	192	ASP
1	F	97	SER

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric Outliers		Percentiles		
1	A	235/243~(97%)	199 (85%)	36 (15%)	2 12		



Continued from previous page...

Mol	Chain	nain Analysed Rotameric Outliers		Outliers	Percentiles		
1	В	235/243 (97%)	199 (85%)	36 (15%)	2 12		
1	С	235/243 (97%)	198 (84%)	37 (16%)	2 12		
1	D	235/243 (97%)	198 (84%)	37 (16%)	2 12		
1	E	235/243 (97%)	201 (86%)	34 (14%)	3 14		
1	F	235/243 (97%)	201 (86%)	34 (14%)	3 14		
1	G	235/243 (97%)	200 (85%)	35 (15%)	3 13		
1	Н	235/243 (97%)	199 (85%)	36 (15%)	2 12		
All	All	1880/1944 (97%)	1595 (85%)	285 (15%)	3 13		

5 of 285 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	155	THR
1	G	225	ILE
1	Н	111	ILE
1	С	222	VAL
1	С	203	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 44 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	215	HIS
1	G	243	ASN
1	F	236	GLN
1	G	213	HIS
1	Н	55	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

24 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Trino	Chain	Dag	Link	Вс	ond leng	ths	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	D	402	-	4,4,4	0.26	0	6,6,6	0.18	0
3	SO4	Н	403	-	4,4,4	0.28	0	6,6,6	0.07	0
3	SO4	С	403	-	4,4,4	0.28	0	6,6,6	0.13	0
3	SO4	A	402	_	4,4,4	0.29	0	6,6,6	0.11	0
3	SO4	Н	402	-	4,4,4	0.30	0	6,6,6	0.09	0
2	AA5	D	401	_	24,24,24	1.43	4 (16%)	29,33,33	1.15	3 (10%)
3	SO4	В	403	-	4,4,4	0.26	0	6,6,6	0.12	0
2	AA5	Н	401	-	24,24,24	1.79	6 (25%)	29,33,33	1.51	4 (13%)
2	AA5	В	401	-	24,24,24	1.51	5 (20%)	29,33,33	1.32	3 (10%)
3	SO4	G	402	-	4,4,4	0.28	0	6,6,6	0.05	0
2	AA5	A	401	-	24,24,24	1.22	3 (12%)	29,33,33	1.11	3 (10%)
3	SO4	G	403	-	4,4,4	0.27	0	6,6,6	0.10	0
3	SO4	Е	402	-	4,4,4	0.26	0	6,6,6	0.15	0
2	AA5	F	401	-	24,24,24	1.43	4 (16%)	29,33,33	1.21	3 (10%)
3	SO4	D	403	-	4,4,4	0.28	0	6,6,6	0.12	0
3	SO4	В	402	-	4,4,4	0.26	0	6,6,6	0.28	0
3	SO4	Е	403	_	4,4,4	0.26	0	6,6,6	0.09	0
2	AA5	С	401	_	24,24,24	1.37	3 (12%)	29,33,33	1.24	4 (13%)
3	SO4	A	403	-	4,4,4	0.35	0	6,6,6	0.14	0
3	SO4	F	403	-	4,4,4	0.32	0	6,6,6	0.06	0
3	SO4	F	402	-	4,4,4	0.22	0	6,6,6	0.18	0
2	AA5	G	401	-	24,24,24	1.48	4 (16%)	29,33,33	1.20	3 (10%)
2	AA5	Е	401	-	24,24,24	1.66	5 (20%)	29,33,33	1.20	3 (10%)
3	SO4	С	402	-	4,4,4	0.28	0	6,6,6	0.22	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AA5	A	401	-	-	4/19/19/19	0/1/1/1
2	AA5	D	401	-	-	3/19/19/19	0/1/1/1
2	AA5	С	401	-	-	2/19/19/19	0/1/1/1
2	AA5	F	401	-	-	1/19/19/19	0/1/1/1
2	AA5	Н	401	-	-	2/19/19/19	0/1/1/1
2	AA5	В	401	-	-	4/19/19/19	0/1/1/1
2	AA5	G	401	-	-	4/19/19/19	0/1/1/1
2	AA5	Е	401	-	-	2/19/19/19	0/1/1/1

The worst 5 of 34 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
2	Н	401	AA5	C4-C3	4.64	1.47	1.40
2	В	401	AA5	C4-C3	4.32	1.47	1.40
2	Е	401	AA5	C4-C3	4.14	1.47	1.40
2	F	401	AA5	C4-C3	4.10	1.47	1.40
2	Н	401	AA5	P-O1P	3.59	1.62	1.50

The worst 5 of 26 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	Н	401	AA5	CA-N-C4A	4.95	124.45	117.31
2	В	401	AA5	CA-N-C4A	3.55	122.43	117.31
2	В	401	AA5	C4-C3-C2	-3.41	118.08	120.19
2	С	401	AA5	CA-N-C4A	3.24	121.99	117.31
2	Ε	401	AA5	CA-N-C4A	3.23	121.97	117.31

There are no chirality outliers.

5 of 22 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	AA5	CB-CA-N-C4A
2	В	401	AA5	C-CA-N-C4A
2	В	401	AA5	CB-CA-N-C4A
2	С	401	AA5	C-CA-N-C4A
2	С	401	AA5	CB-CA-N-C4A




There are no ring outliers.

9 monomers are involved in 27 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	401	AA5	2	0
3	В	403	SO4	1	0
2	Н	401	AA5	4	0
2	В	401	AA5	5	0
2	A	401	AA5	5	0
2	F	401	AA5	2	0
2	С	401	AA5	3	0
2	G	401	AA5	2	0
2	Е	401	AA5	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ} {>} 2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	303/311 (97%)	0.12	6 (1%) 65 64	14, 44, 65, 71	0
1	В	303/311 (97%)	-0.24	0 100 100	11, 26, 49, 63	0
1	С	303/311 (97%)	-0.30	0 100 100	7, 26, 47, 62	0
1	D	303/311 (97%)	-0.16	6 (1%) 65 64	16, 33, 54, 64	0
1	E	303/311 (97%)	0.11	3 (0%) 82 82	19, 45, 66, 75	0
1	F	303/311 (97%)	-0.08	3 (0%) 82 82	14, 31, 50, 63	0
1	G	303/311 (97%)	0.26	7 (2%) 60 59	24, 54, 83, 94	0
1	Н	303/311 (97%)	1.65	107 (35%) 0 0	46, 93, 114, 118	0
All	All	2424/2488 (97%)	0.17	132 (5%) 25 24	7, 39, 99, 118	0

The worst 5 of 132 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Н	221	GLY	7.5
1	Н	88	THR	6.0
1	Н	109	ALA	5.6
1	Н	193	SER	5.3
1	Н	125	ARG	4.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q<0.9
3	SO4	Н	403	5/5	0.67	0.37	145,146,146,146	0
3	SO4	G	403	5/5	0.74	0.24	105,106,106,107	0
2	AA5	Н	401	24/24	0.74	0.32	85,87,88,89	0
3	SO4	A	403	5/5	0.85	0.25	85,85,86,87	0
3	SO4	G	402	5/5	0.86	0.29	121,121,121,121	0
3	SO4	Н	402	5/5	0.89	0.30	103,104,104,104	0
3	SO4	D	403	5/5	0.90	0.20	75,75,76,76	0
3	SO4	F	403	5/5	0.90	0.24	86,86,87,87	0
3	SO4	Е	403	5/5	0.91	0.19	88,88,89,89	0
2	AA5	Е	401	24/24	0.91	0.26	45,53,57,57	0
3	SO4	Е	402	5/5	0.91	0.17	78,78,79,79	0
3	SO4	В	403	5/5	0.93	0.21	68,68,69,69	0
3	SO4	С	403	5/5	0.93	0.17	69,69,70,70	0
2	AA5	G	401	24/24	0.94	0.20	45,47,48,49	0
3	SO4	В	402	5/5	0.94	0.21	63,64,64,64	0
2	AA5	A	401	24/24	0.94	0.22	39,42,49,53	0
2	AA5	F	401	24/24	0.95	0.17	18,23,27,29	0
3	SO4	A	402	5/5	0.95	0.13	62,63,63,64	0
3	SO4	D	402	5/5	0.96	0.16	65,66,66,67	0
2	AA5	D	401	24/24	0.96	0.17	25,30,31,33	0
3	SO4	С	402	5/5	0.96	0.16	59,60,61,61	0
2	AA5	В	401	24/24	0.96	0.16	6,17,21,25	0
3	SO4	F	402	5/5	0.96	0.13	61,62,63,64	0
2	AA5	С	401	24/24	0.97	0.15	16,21,24,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







































6.5 Other polymers (i)

There are no such residues in this entry.

