



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 21, 2023 – 01:49 pm BST

PDB ID : 8BBK
Title : Crystal structure of human Sirt3 in complex with a fragment of the human AROS protein
Authors : Steegborn, C.; Weiss, S.
Deposited on : 2022-10-13
Resolution : 3.27 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.27 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1177 (3.32-3.24)
Clashscore	141614	1044 (3.30-3.26)
Ramachandran outliers	138981	1026 (3.30-3.26)
Sidechain outliers	138945	1025 (3.30-3.26)
RSRZ outliers	127900	1141 (3.32-3.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	399	
1	B	399	
1	C	399	
1	D	399	
1	E	399	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	399	 % 51% 17% 31%
2	G	136	 98%
2	H	136	 98%
2	I	136	 98%
2	J	136	 98%
2	K	136	 98%
2	L	136	 98%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 13135 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NAD-dependent protein deacetylase sirtuin-3, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	274	Total 2148	C 1384	N 370	O 385	S 9	0	0	0
1	B	274	Total 2159	C 1393	N 371	O 386	S 9	0	1	0
1	C	274	Total 2159	C 1393	N 371	O 386	S 9	0	1	0
1	D	274	Total 2159	C 1393	N 371	O 386	S 9	0	1	0
1	E	274	Total 2159	C 1393	N 371	O 386	S 9	0	1	0
1	F	274	Total 2159	C 1393	N 371	O 386	S 9	0	1	0

- Molecule 2 is a protein called Active regulator of SIRT1.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	G	3	Total 31	C 18	N 10	O 3	0	0	0
2	H	3	Total 31	C 18	N 10	O 3	0	0	0
2	I	3	Total 31	C 18	N 10	O 3	0	0	0
2	J	3	Total 31	C 18	N 10	O 3	0	0	0
2	K	3	Total 31	C 18	N 10	O 3	0	0	0
2	L	3	Total 31	C 18	N 10	O 3	0	0	0


- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	Zn 1	0	0
3	B	1	Total 1	Zn 1	0	0
3	C	1	Total 1	Zn 1	0	0
3	D	1	Total 1	Zn 1	0	0
3	E	1	Total 1	Zn 1	0	0
3	F	1	Total 1	Zn 1	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NAD-dependent protein deacetylase sirtuin-3, mitochondrial




- Molecule 1: NAD-dependent protein deacetylase sirtuin-3, mitochondrial




- Molecule 1: NAD-dependent protein deacetylase sirtuin-3, mitochondrial






● Molecule 1: NAD-dependent protein deacetylase sirtuin-3, mitochondrial



● Molecule 1: NAD-dependent protein deacetylase sirtuin-3, mitochondrial



● Molecule 1: NAD-dependent protein deacetylase sirtuin-3, mitochondrial



MET	LEU	G123	D967	ASP
ALA	ASP	K122	GLY	LYS
TRP	PRO	L123	R279	
GLY	ALA	Q126	T284	
TRP	ARG	K65	V292	
ARG	LEU	R136	F294	
ALA	GLN	A134	R294	
ALA	ARG	R139	G295	
ALA	PRO	V140	E296	
ALA	PRO	V141	P297	
LEU	ARG	V142	L298	
ARG	PRO	G145	F309	
LEU	VAL	L164	A312	
TRP	PRO	I179	D313	
GLY	ARG	P183	L314	
GLY	ARG	F184	L315	
GLY	ARG	H187	I317	
GLY	ARG	M188	L318	
GLY	ALA	P191	L319	
GLY	ALA	L195	L320	
VAL	PRO	A196	S221	
VAL	PRO	P200	L322	
VAL	PRO	P201	E223	
CYS	ILE	L213	V324	
CYS	LYS	L216	E325	
CYS	GLY	H217	P326	
LEU	GLY	L222	F327	
VAL	ARG	Y226	L342	
LEU	ILE	Q228	I343	
ARG	SER	R235	N344	
ASP	PHE	I239	V348	
ASP	SER	L244	G349	
VAL	SER	V245	L363	
VAL	VAL	H248	V374	
GLY	VAL	Q256	E375	
GLY	VAL	T257	L376	
HIS	GLY	V258	L377	
GLY	GLY	P264	G378	
ALA	GLY		W379	
ARG	GLY		E382	
GLY	GLY		M383	
ALA	SER		R384	
ARG	ASP		D385	
GLY	GLY		Q388	
GLY	GLY		L394	
GLY	ASP		ASP	
PRO	GLY		GLY	
PRO	LYS		PRO	

Molecule 2: Active regulator of SIRT1




MET	SER	ASP	PHE	
ALA	GLY	TYR	THR	
ALA	ALA	R64	GLU	
LEU	LEU	K65	GLU	
LEU	ARG	R66	ASP	
ARG	ARG	GLU	PHE	
GLY	GLY	CYS	GLN	
GLY	GLY	LYS	LYS	
LEU	GLY	ARG	PHE	
LEU	GLY	ASP	GLN	
LEU	GLY	HIS	GLN	
LEU	LEU	LEU	GLU	
LEU	LEU	ARG	GLU	
ALA	VAL	VAL	TYR	
ALA	ASN	ASN	PHE	
SER	LEU	LEU	GLY	
GLY	SER	LYS	SER	
ALA	PHE	LEU	THR	
PRO	ALA	THR	ARG	
PRO	ARG	ASP	ASP	
PRO	PRO	THR	THR	
PRO	ARG	THR	THR	
GLY	GLY	SER	THR	
GLY	GLY	THR	THR	
GLN	THR	VAL	VAL	
ALA	ALA	ALA	ALA	
LYS	LYS	LYS	LYS	
PRO	PRO	GLU	GLU	
ARG	ARG	SER	SER	
GLY	VAL	VAL	VAL	
ALA	SER	SER	SER	
PRO	ALA	ALA	ALA	
VAL	VAL	GLN	GLN	
LYS	ILE	ILE	ILE	
ARG	ARG	ARG	ARG	
PRO	PRO	PRO	PRO	
ARG	ARG	GLN	GLN	
ARG	ARG	ASN	ASN	
LYS	LYS	ARG	THR	
THR	LYS	GLY	LYS	
ALA	ALA	ILE	ILE	
GLN	GLN	LYS	LYS	
ALA	ALA	CYS	CYS	
GLN	GLN	ASP	ASP	
LYS	LYS	ARG	ARG	
LEU	LEU	PRO	PRO	
ARG	ARG	VAL	VAL	
ASN	ASN	ALA	ALA	
SER	SER	LYS	LYS	
ALA	ALA	THR	THR	
LYS	LYS	LYS	LYS	
GLY	GLY	LYS	LYS	
VAL	VAL	LYS	LYS	
PRO	PRO	ALA	ALA	
LYS	LYS	GLU	GLU	
ALA	ALA	GLY	GLY	
LEU	LEU	THR	THR	

Molecule 2: Active regulator of SIRT1



MET	SER	ASP	PHE	
ALA	GLY	TYR	THR	
ALA	ALA	R64	GLU	
LEU	LEU	K65	GLU	
LEU	ARG	R66	ASP	
ARG	ARG	GLU	PHE	
GLY	GLY	CYS	GLN	
GLY	GLY	LYS	LYS	
LEU	GLY	ARG	PHE	
LEU	GLY	ASP	GLN	
LEU	GLY	HIS	GLN	
LEU	LEU	LEU	GLU	
LEU	LEU	ARG	GLU	
ALA	VAL	VAL	TYR	
ALA	ASN	ASN	PHE	
SER	LEU	LEU	GLY	
GLY	SER	LYS	SER	
ALA	PHE	LEU	THR	
PRO	ALA	THR	ARG	
PRO	ARG	ASP	ASP	
PRO	PRO	THR	THR	
PRO	ARG	THR	THR	
GLY	GLY	SER	THR	
GLY	GLY	THR	THR	
GLN	THR	VAL	VAL	
ALA	ALA	ALA	ALA	
LYS	LYS	LYS	LYS	
PRO	PRO	GLU	GLU	
ARG	ARG	SER	SER	
GLY	VAL	VAL	VAL	
ALA	SER	SER	SER	
PRO	ALA	ALA	ALA	
VAL	VAL	GLN	GLN	
LYS	ILE	ILE	ILE	
ARG	ARG	ARG	ARG	
PRO	PRO	PRO	PRO	
ARG	ARG	GLN	GLN	
ARG	ARG	ASN	ASN	
LYS	LYS	ARG	THR	
THR	LYS	GLY	LYS	
ALA	ALA	ILE	ILE	
GLN	GLN	LYS	LYS	
ALA	ALA	CYS	CYS	
GLN	GLN	ASP	ASP	
LYS	LYS	ARG	ARG	
LEU	LEU	PRO	PRO	
ARG	ARG	VAL	VAL	
ASN	ASN	ALA	ALA	
SER	SER	LYS	LYS	
ALA	ALA	THR	THR	
LYS	LYS	LYS	LYS	
GLY	GLY	LYS	LYS	
VAL	VAL	LYS	LYS	
PRO	PRO	ALA	ALA	
LYS	LYS	GLU	GLU	
ALA	ALA	GLY	GLY	
LEU	LEU	THR	THR	

Molecule 2: Active regulator of SIRT1



MET	SER	ASP	PHE	
ALA	GLY	TYR	THR	
ALA	ALA	R64	GLU	
LEU	LEU	K65	GLU	
LEU	ARG	R66	ASP	
ARG	ARG	GLU	PHE	
GLY	GLY	CYS	GLN	
GLY	GLY	LYS	LYS	
LEU	GLY	ARG	PHE	
LEU	GLY	ASP	GLN	
LEU	GLY	HIS	GLN	
LEU	LEU	LEU	GLU	
LEU	LEU	ARG	GLU	
ALA	VAL	VAL	TYR	
ALA	ASN	ASN	PHE	
SER	LEU	LEU	GLY	
GLY	SER	LYS	SER	
ALA	PHE	LEU	THR	
PRO	ALA	THR	ARG	
PRO	ARG	ASP	ASP	
PRO	PRO	THR	THR	
PRO	ARG	THR	THR	
GLY	GLY	SER	THR	
GLY	GLY	THR	THR	
GLN	THR	VAL	VAL	
ALA	ALA	ALA	ALA	
LYS	LYS	LYS	LYS	
PRO	PRO	GLU	GLU	
ARG	ARG	SER	SER	
GLY	VAL	VAL	VAL	
ALA	SER	SER	SER	
PRO	ALA	ALA	ALA	
VAL	VAL	GLN	GLN	
LYS	ILE	ILE	ILE	
ARG	ARG	ARG	ARG	
PRO	PRO	PRO	PRO	
ARG	ARG	GLN	GLN	
ARG	ARG	ASN	ASN	
LYS	LYS	ARG	THR	
THR	LYS	GLY	LYS	
ALA	ALA	ILE	ILE	
GLN	GLN	LYS	LYS	
ALA	ALA	CYS	CYS	
GLN	GLN	ASP	ASP	
LYS	LYS	ARG	ARG	
LEU	LEU	PRO	PRO	
ARG	ARG	VAL	VAL	
ASN	ASN	ALA	ALA	
SER	SER	LYS	LYS	
ALA	ALA	THR	THR	
LYS	LYS	LYS	LYS	
GLY	GLY	LYS	LYS	
VAL	VAL	LYS	LYS	
PRO	PRO	ALA	ALA	
LYS	LYS	GLU	GLU	
ALA	ALA	GLY	GLY	
LEU	LEU	THR	THR	

4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	110.22Å 110.22Å 344.36Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.19 – 3.27 49.19 – 3.27	Depositor EDS
% Data completeness (in resolution range)	99.7 (49.19-3.27) 99.7 (49.19-3.27)	Depositor EDS
R_{merge}	0.32	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.08 (at 3.25Å)	Xtrriage
Refinement program	PHENIX 1.18.2_3874	Depositor
R, R_{free}	0.202 , 0.261 0.203 , 0.262	Depositor DCC
R_{free} test set	1929 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	103.0	Xtrriage
Anisotropy	0.294	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 52.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.037 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	13135	wwPDB-VP
Average B, all atoms (Å ²)	95.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.57	0/2203	0.78	3/3001 (0.1%)
1	B	0.54	0/2215	0.79	0/3017
1	C	0.52	0/2215	0.76	0/3017
1	D	0.53	0/2215	0.76	2/3017 (0.1%)
1	E	0.58	1/2215 (0.0%)	0.78	0/3017
1	F	0.56	1/2215 (0.0%)	0.80	1/3017 (0.0%)
2	G	0.67	0/30	1.11	0/36
2	H	0.58	0/30	1.13	0/36
2	I	1.03	0/30	1.40	1/36 (2.8%)
2	J	0.97	0/30	2.01	1/36 (2.8%)
2	K	0.68	0/30	0.94	0/36
2	L	0.81	0/30	1.54	0/36
All	All	0.55	2/13458 (0.0%)	0.79	8/18302 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	256	CYS	CB-SG	-5.67	1.72	1.81
1	E	256	CYS	CB-SG	-5.02	1.73	1.81

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	65	LYS	CD-CE-NZ	8.25	130.67	111.70
1	A	139	ARG	CD-NE-CZ	8.19	135.07	123.60
1	A	358	ARG	NE-CZ-NH2	-7.93	116.33	120.30
1	A	139	ARG	CG-CD-NE	-6.42	98.32	111.80
2	I	65	LYS	CD-CE-NZ	5.82	125.09	111.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2148	0	2163	54	0
1	B	2159	0	2171	46	1
1	C	2159	0	2171	48	0
1	D	2159	0	2171	50	0
1	E	2159	0	2171	48	1
1	F	2159	0	2171	58	0
2	G	31	0	38	4	0
2	H	31	0	38	4	0
2	I	31	0	38	1	0
2	J	31	0	38	3	0
2	K	31	0	38	4	0
2	L	31	0	38	6	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
All	All	13135	0	13246	300	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 300 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:248:HIS:HD2	2:G:65:LYS:HE3	1.06	1.11
1:A:248:HIS:CD2	2:G:65:LYS:HE3	1.88	1.07
1:F:293:PHE:N	1:F:296:GLU:OE1	2.01	0.92
1:D:248:HIS:CD2	2:J:65:LYS:HE2	2.07	0.88
1:F:292:VAL:HG12	2:L:65:LYS:HD2	1.55	0.86

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:337:SER:OG	1:E:130:GLU:OE1[6_554]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/399 (68%)	261 (96%)	11 (4%)	0	100	100
1	B	273/399 (68%)	264 (97%)	8 (3%)	1 (0%)	34	67
1	C	273/399 (68%)	259 (95%)	13 (5%)	1 (0%)	34	67
1	D	273/399 (68%)	264 (97%)	8 (3%)	1 (0%)	34	67
1	E	273/399 (68%)	264 (97%)	8 (3%)	1 (0%)	34	67
1	F	273/399 (68%)	261 (96%)	9 (3%)	3 (1%)	14	46
2	G	1/136 (1%)	1 (100%)	0	0	100	100
2	H	1/136 (1%)	0	1 (100%)	0	100	100
2	I	1/136 (1%)	1 (100%)	0	0	100	100
2	J	1/136 (1%)	0	1 (100%)	0	100	100
2	K	1/136 (1%)	1 (100%)	0	0	100	100
2	L	1/136 (1%)	1 (100%)	0	0	100	100
All	All	1643/3210 (51%)	1577 (96%)	59 (4%)	7 (0%)	34	67

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	294	PHE
1	F	183	PRO
1	C	183	PRO
1	D	183	PRO
1	E	183	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	237/327 (72%)	237 (100%)	0	100	100
1	B	238/327 (73%)	238 (100%)	0	100	100
1	C	238/327 (73%)	238 (100%)	0	100	100
1	D	238/327 (73%)	238 (100%)	0	100	100
1	E	238/327 (73%)	238 (100%)	0	100	100
1	F	238/327 (73%)	237 (100%)	1 (0%)	91	95
2	G	3/114 (3%)	3 (100%)	0	100	100
2	H	3/114 (3%)	3 (100%)	0	100	100
2	I	3/114 (3%)	3 (100%)	0	100	100
2	J	3/114 (3%)	2 (67%)	1 (33%)	0	0
2	K	3/114 (3%)	3 (100%)	0	100	100
2	L	3/114 (3%)	3 (100%)	0	100	100
All	All	1445/2646 (55%)	1443 (100%)	2 (0%)	93	97

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	J	66	ARG
1	F	294	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	248	HIS
1	D	169	GLN
1	D	248	HIS
1	F	354	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	274/399 (68%)	-0.25	1 (0%) 92 93	62, 92, 132, 155	0
1	B	274/399 (68%)	-0.21	1 (0%) 92 93	58, 91, 126, 142	0
1	C	274/399 (68%)	-0.25	0 100 100	66, 94, 126, 140	0
1	D	274/399 (68%)	-0.23	1 (0%) 92 93	67, 99, 138, 162	0
1	E	274/399 (68%)	-0.27	1 (0%) 92 93	71, 87, 112, 127	0
1	F	274/399 (68%)	-0.35	2 (0%) 87 88	30, 92, 118, 146	0
2	G	3/136 (2%)	-0.04	0 100 100	98, 98, 100, 114	0
2	H	3/136 (2%)	0.24	0 100 100	100, 100, 101, 115	0
2	I	3/136 (2%)	-0.32	0 100 100	95, 95, 95, 109	0
2	J	3/136 (2%)	0.29	0 100 100	93, 93, 108, 122	0
2	K	3/136 (2%)	0.45	0 100 100	100, 100, 112, 122	0
2	L	3/136 (2%)	-0.34	0 100 100	95, 95, 108, 119	0
All	All	1662/3210 (51%)	-0.26	6 (0%) 92 93	30, 92, 127, 162	0

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	121	GLY	3.4
1	D	121	GLY	3.2
1	F	394	LEU	2.3
1	B	347	LEU	2.1
1	F	257	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ZN	F	401	1/1	0.98	0.14	101,101,101,101	0
3	ZN	B	401	1/1	0.99	0.15	81,81,81,81	0
3	ZN	C	401	1/1	0.99	0.17	133,133,133,133	0
3	ZN	D	401	1/1	0.99	0.18	92,92,92,92	0
3	ZN	E	401	1/1	0.99	0.22	87,87,87,87	0
3	ZN	A	401	1/1	0.99	0.20	73,73,73,73	0

6.5 Other polymers [i](#)

There are no such residues in this entry.