



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 13, 2021 – 08:02 pm BST

PDB ID : 7BG1
Title : Structure of anti-FLAG M2 Fab domain remodeled based on proteomic sequencing
Authors : Pronker, M.F.; Snijder, J.
Deposited on : 2021-01-05
Resolution : 1.86 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.22
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.22


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.86 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	222	
2	L	220	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	CL	H	302	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 3497 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called anti-FLAG M2 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	210	1600	1016	259	318	7	0	0	0

- Molecule 2 is a protein called anti-FLAG M2 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	210	1636	1028	276	326	6	0	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
3	H	1	5	4	1	0	0
3	L	1	5	4	1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	H	1	Total Cl 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	H	143	Total O 143 143	0	0
5	L	107	Total O 107 107	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: anti-FLAG M2 heavy chain



- Molecule 2: anti-FLAG M2 light chain



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	87.36Å 133.76Å 41.48Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.52 – 1.86 41.52 – 1.86	Depositor EDS
% Data completeness (in resolution range)	95.7 (41.52-1.86) 95.7 (41.52-1.86)	Depositor EDS
R_{merge}	0.03	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.12 (at 1.86Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.217 , 0.255 0.217 , 0.255	Depositor DCC
R_{free} test set	2008 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	30.7	Xtrriage
Anisotropy	0.071	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 56.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	3497	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	H	0.29	0/1645	0.53	0/2252
2	L	0.27	0/1674	0.49	0/2270
All	All	0.28	0/3319	0.51	0/4522

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1600	0	1543	9	0
2	L	1636	0	1583	13	0
3	H	5	0	0	0	0
3	L	5	0	0	0	0
4	H	1	0	0	2	0
5	H	143	0	0	1	0
5	L	107	0	0	0	0
All	All	3497	0	3126	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:20:MET:HE1	1:H:113:LEU:HD22	1.86	0.58
2:L:200:GLU:HG2	2:L:211:VAL:HG22	1.85	0.57
1:H:101:PHE:N	4:H:302:CL:CL	2.75	0.54
2:L:135:ALA:HB3	2:L:186:LEU:HD12	1.92	0.52
1:H:38:LYS:HB2	1:H:48:ILE:HD11	1.92	0.52
2:L:127:SER:O	2:L:131:THR:HG23	2.14	0.47
2:L:151:VAL:HG21	2:L:180:MET:HE1	1.97	0.47
1:H:9:ALA:HB2	1:H:155:PRO:HD2	1.96	0.47
2:L:128:GLU:OE2	2:L:128:GLU:N	2.42	0.46
1:H:1:GLN:N	5:H:401:HOH:O	2.33	0.46
2:L:59:ARG:HG2	2:L:63:VAL:HB	1.99	0.45
1:H:123:PRO:HB3	1:H:151:TYR:HB3	1.98	0.45
1:H:100:LYS:N	4:H:302:CL:CL	2.85	0.45
1:H:143:THR:HG22	1:H:196:THR:OG1	2.18	0.43
2:L:212:LYS:HD3	2:L:212:LYS:HA	1.74	0.43
2:L:24:ARG:HA	2:L:74:THR:O	2.18	0.43
1:H:39:GLN:NE2	2:L:43:LEU:HD21	2.34	0.42
2:L:64:PRO:HG2	2:L:67:PHE:CE1	2.54	0.41
2:L:29:ILE:HG13	2:L:36:THR:HG23	2.02	0.41
2:L:118:PRO:HB3	2:L:144:PHE:CD2	2.56	0.41
2:L:147:LYS:HD2	2:L:178:TYR:CE1	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	206/222 (93%)	202 (98%)	4 (2%)	0	100	100
2	L	206/220 (94%)	200 (97%)	5 (2%)	1 (0%)	29	15
All	All	412/442 (93%)	402 (98%)	9 (2%)	1 (0%)	47	33

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	L	73	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	180/189 (95%)	174 (97%)	6 (3%)	38	21
2	L	188/197 (95%)	182 (97%)	6 (3%)	39	22
All	All	368/386 (95%)	356 (97%)	12 (3%)	38	21

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	10	GLU
1	H	65	LYS
1	H	89	GLU
1	H	191	LEU
1	H	209	CYS
1	H	210	ASN
2	L	11	LEU
2	L	15	LEU
2	L	78	LEU
2	L	168	TRP
2	L	186	LEU
2	L	214	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	L	301	-	4,4,4	0.14	0	6,6,6	0.10	0
3	SO4	H	301	-	4,4,4	0.16	0	6,6,6	0.12	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	H	210/222 (94%)	0.52	20 (9%) 8 7	21, 36, 80, 112	0
2	L	210/220 (95%)	1.06	50 (23%) 0 0	23, 52, 121, 142	0
All	All	420/442 (95%)	0.79	70 (16%) 1 1	21, 42, 108, 142	0

All (70) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	189	ASP	6.5
2	L	208	SER	6.3
2	L	122	ILE	6.2
2	L	214	PHE	5.9
2	L	158	SER	5.6
2	L	199	CYS	4.8
1	H	197	VAL	4.2
2	L	202	THR	4.1
1	H	143	THR	4.1
2	L	210	ILE	4.1
1	H	225	VAL	4.1
2	L	130	LEU	4.0
1	H	227	ARG	4.0
1	H	201	THR	3.9
1	H	226	PRO	3.8
2	L	207	THR	3.7
2	L	211	VAL	3.7
1	H	141	MET	3.6
2	L	118	PRO	3.6
1	H	202	TRP	3.6
2	L	155	ILE	3.6
2	L	73	GLY	3.5
2	L	24	ARG	3.4
1	H	142	VAL	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	209	PRO	3.4
2	L	153	TRP	3.2
2	L	151	VAL	3.2
2	L	120	VAL	3.1
2	L	198	THR	3.0
2	L	157	GLY	3.0
2	L	201	ALA	3.0
2	L	204	LYS	3.0
2	L	134	GLY	3.0
2	L	117	ALA	3.0
1	H	102	TYR	2.9
1	H	140	SER	2.9
2	L	139	CYS	2.9
2	L	194	HIS	2.9
1	H	205	GLN	2.9
1	H	206	THR	2.8
1	H	1	GLN	2.8
2	L	123	PHE	2.8
1	H	129	ALA	2.7
1	H	130	PRO	2.7
2	L	150	ASN	2.7
2	L	115	ASP	2.6
2	L	205	THR	2.6
1	H	128	LEU	2.6
2	L	187	THR	2.6
2	L	148	ASP	2.5
2	L	186	LEU	2.5
2	L	156	ASP	2.5
2	L	193	ARG	2.5
2	L	152	LYS	2.5
2	L	141	LEU	2.5
2	L	124	PRO	2.4
1	H	204	SER	2.4
1	H	101	PHE	2.4
2	L	168	TRP	2.4
2	L	206	SER	2.3
2	L	197	TYR	2.3
2	L	191	TYR	2.2
2	L	131	THR	2.2
2	L	137	VAL	2.2
2	L	184	LEU	2.2
2	L	32	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	185	THR	2.1
1	H	198	PRO	2.1
2	L	183	THR	2.1
2	L	159	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	CL	H	302	1/1	0.62	0.12	83,83,83,83	0
3	SO4	H	301	5/5	0.94	0.10	49,50,56,61	0
3	SO4	L	301	5/5	0.96	0.12	57,64,65,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.