



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 28, 2024 – 03:20 pm GMT

PDB ID : 8BIX
Title : Cystathionine gamma-lyase N360S mutant from *Toxoplasma gondii* in complex with cystathionine
Authors : Fernandez-Rodriguez, C.; Conter, C.; Oyenarte, I.; Favretto, F.; Quintana, I.; Martinez-Chantar, M.L.; Astegno, A.; Martinez-Cruz, L.A.
Deposited on : 2022-11-02
Resolution : 1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	417	
1	C	417	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6579 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Cystathionine beta-lyase, putative.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	C	396	Total 3055	C 1949	N 517	O 568	S 21	0	1	0
1	A	396	Total 3072	C 1958	N 520	O 573	S 21	0	3	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	360	SER	ASN	engineered mutation	UNP B6K8Y1
A	360	SER	ASN	engineered mutation	UNP B6K8Y1

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	C	1	Total 15	C 8	N 1	O 5	P 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	15	8	1	5	1	0	0

- Molecule 3 is 2-AMINO-4-MERCAPTO-BUTYRIC ACID (three-letter code: HCS) (formula: C₄H₉NO₂S) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	8	4	1	2	1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	204	Total O 204 204	0	0
4	A	210	Total O 210 210	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Cystathionine beta-lyase, putative



- Molecule 1: Cystathionine beta-lyase, putative



4 Data and refinement statistics i

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	158.43Å 158.43Å 93.83Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	55.38 – 1.59 55.38 – 1.59	Depositor EDS
% Data completeness (in resolution range)	97.7 (55.38-1.59) 97.7 (55.38-1.59)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.51 (at 1.59Å)	Xtrriage
Refinement program	PHENIX 1.18	Depositor
R, R_{free}	0.172 , 0.190 0.172 , 0.190	Depositor DCC
R_{free} test set	8741 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å ²)	23.0	Xtrriage
Anisotropy	0.021	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 50.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.052 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6579	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HCS, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.78	2/3146 (0.1%)	0.80	2/4269 (0.0%)
1	C	0.73	1/3126 (0.0%)	0.79	0/4242
All	All	0.76	3/6272 (0.0%)	0.80	2/8511 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	195	CYS	CB-SG	-6.96	1.70	1.82
1	C	179	CYS	CB-SG	-5.67	1.72	1.81
1	A	179	CYS	CB-SG	-5.35	1.73	1.81

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	150	ASP	CB-CG-OD1	6.03	123.73	118.30
1	A	187	ASP	CB-CG-OD2	5.54	123.28	118.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	142	LEU	Peptide
1	C	22	VAL	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3072	0	3037	29	1
1	C	3055	0	3026	26	1
2	A	15	0	7	0	0
2	C	15	0	7	1	0
3	A	8	0	6	0	0
4	A	210	0	0	2	0
4	C	204	0	0	6	0
All	All	6579	0	6083	46	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:253:LYS:HE2	4:C:604:HOH:O	1.85	0.76
1:C:68:VAL:HG21	1:A:345:ILE:HG23	1.67	0.75
1:A:337:VAL:HG11	1:A:344:GLY:HA2	1.71	0.73
1:A:249:GLU:OE2	4:A:601:HOH:O	2.08	0.72
1:C:359:GLU:OE1	1:A:77:SER:HB2	1.92	0.69
1:A:189[A]:GLU:HG3	1:A:219:LEU:HB3	1.76	0.66
1:A:55:PRO:HG3	1:A:274:LEU:HD13	1.78	0.65
1:A:341:THR:OG1	1:A:391:ASP:HB3	1.98	0.63
1:C:278[B]:LYS:HE3	4:C:791:HOH:O	1.97	0.63
1:A:119:PHE:CE1	1:A:253:LYS:HE2	2.35	0.60
1:C:55:PRO:HB2	4:C:626:HOH:O	2.02	0.58
1:C:373:VAL:HG11	1:A:68:VAL:HG13	1.84	0.57
1:C:347:MET:SD	1:C:348:MET:HE2	2.45	0.56
1:C:275:LEU:HG	4:C:724:HOH:O	2.05	0.56
1:C:347:MET:SD	1:C:348:MET:CE	2.95	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:230:LYS:HD2	1:A:361:LEU:HG	1.88	0.54
1:C:198:LYS:NZ	4:C:603:HOH:O	2.27	0.52
1:C:347:MET:HE1	1:C:394:ILE:HG13	1.90	0.52
1:C:261:MET:CE	1:A:113:VAL:HG11	2.38	0.52
1:A:135:GLY:HA2	1:A:138[A]:ARG:NH2	2.26	0.51
1:C:68:VAL:HG22	1:A:349:ASN:OD1	2.11	0.50
1:C:137:ASN:OD1	1:C:141:ARG:HD3	2.12	0.49
1:A:348:MET:HA	1:A:348:MET:CE	2.44	0.48
1:C:92:LYS:HE3	4:C:783:HOH:O	2.13	0.48
1:C:137:ASN:OD1	1:C:141:ARG:NH2	2.48	0.47
1:A:337:VAL:HG11	1:A:344:GLY:CA	2.43	0.47
1:C:261:MET:HE1	1:A:113:VAL:HG11	1.96	0.47
1:A:351:VAL:HG11	1:A:368:ILE:HG21	1.96	0.46
1:A:138[B]:ARG:HG3	1:A:142:LEU:HD12	1.97	0.46
1:A:40:VAL:HA	1:A:278[B]:LYS:HG2	1.98	0.46
1:A:131:CYS:HB3	4:A:686:HOH:O	2.17	0.45
1:A:379:MET:HB3	1:A:384:ARG:HG2	1.98	0.45
1:C:337:VAL:HG11	1:C:344:GLY:HA2	1.99	0.45
1:A:181:PRO:HD2	1:A:182:THR:HG23	1.99	0.44
1:C:261:MET:HE1	1:A:113:VAL:CG1	2.48	0.43
1:C:305:GLU:HG3	1:C:306:HIS:CE1	2.54	0.43
1:C:387:VAL:HG23	1:C:389:ILE:HD12	1.99	0.42
1:C:207:THR:HB	2:C:501:PLP:H2A2	2.01	0.42
1:C:127:LEU:HD23	1:C:152:GLU:HB3	2.01	0.42
1:C:261:MET:HE2	1:A:113:VAL:HG11	2.01	0.42
1:A:253:LYS:HE3	1:A:253:LYS:HB3	1.61	0.41
1:C:138:ARG:O	1:C:138:ARG:HG2	2.19	0.41
1:C:373:VAL:HG11	1:A:68:VAL:CG1	2.51	0.41
1:A:140:ALA:HB1	1:A:151:PHE:CG	2.56	0.41
1:A:137:ASN:HD21	1:A:141:ARG:HH11	1.68	0.41
1:A:138[B]:ARG:HH12	1:A:378:ASN:HB2	1.85	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:23:LYS:NZ	1:A:349:ASN:O[6_555]	2.12	0.08

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	397/417 (95%)	387 (98%)	10 (2%)	0	100	100
1	C	395/417 (95%)	381 (96%)	14 (4%)	0	100	100
All	All	792/834 (95%)	768 (97%)	24 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	336/354 (95%)	332 (99%)	4 (1%)	71	54
1	C	334/354 (94%)	332 (99%)	2 (1%)	86	77
All	All	670/708 (95%)	664 (99%)	6 (1%)	78	65

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	139	CYS
1	C	253	LYS
1	A	139	CYS
1	A	302	HIS
1	A	347	MET
1	A	370	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such

sidechains are listed below:

Mol	Chain	Res	Type
1	C	378	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PLP	C	501	1	15,15,16	7.86	5 (33%)	20,22,23	2.01	6 (30%)
3	HCS	A	502	1	6,7,7	0.89	0	7,8,8	2.65	3 (42%)
2	PLP	A	501	1	15,15,16	7.39	6 (40%)	20,22,23	2.55	10 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	C	501	1	-	0/6/6/8	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HCS	A	502	1	-	5/7/7/7	-
2	PLP	A	501	1	-	0/6/6/8	0/1/1/1

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	PLP	C3-C2	-23.43	1.17	1.40
2	A	501	PLP	C3-C2	-21.80	1.19	1.40
2	C	501	PLP	C5-C4	-17.07	1.21	1.40
2	A	501	PLP	C5-C4	-15.80	1.23	1.40
2	A	501	PLP	C6-N1	6.81	1.48	1.34
2	C	501	PLP	C6-N1	6.65	1.48	1.34
2	A	501	PLP	C3-C4	5.32	1.51	1.40
2	C	501	PLP	C3-C4	4.82	1.50	1.40
2	C	501	PLP	C2A-C2	2.44	1.54	1.50
2	A	501	PLP	C2A-C2	2.38	1.54	1.50
2	A	501	PLP	P-O2P	-2.28	1.46	1.54

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	PLP	C5-C6-N1	-5.82	114.12	123.82
2	A	501	PLP	C2A-C2-C3	5.66	127.88	120.89
3	A	502	HCS	CG-CB-CA	-5.11	104.80	113.14
2	C	501	PLP	C5-C6-N1	-5.00	115.49	123.82
2	C	501	PLP	C5A-C5-C6	-3.72	113.25	119.37
2	A	501	PLP	C5A-C5-C6	-3.59	113.47	119.37
2	C	501	PLP	C2A-C2-C3	3.19	124.83	120.89
2	A	501	PLP	C6-C5-C4	3.14	120.63	118.16
3	A	502	HCS	OXT-C-CA	2.93	123.36	113.38
2	C	501	PLP	C6-C5-C4	2.78	120.35	118.16
3	A	502	HCS	CB-CG-SD	-2.65	110.98	113.74
2	A	501	PLP	O3-C3-C2	2.35	122.61	117.49
2	A	501	PLP	C3-C4-C5	2.27	121.19	118.74
2	A	501	PLP	C4A-C4-C3	-2.16	116.84	120.50
2	A	501	PLP	C2A-C2-N1	-2.11	113.55	117.67
2	C	501	PLP	C4-C3-C2	2.10	123.18	120.07
2	A	501	PLP	O4P-P-O1P	-2.10	100.59	106.47
2	A	501	PLP	C6-N1-C2	2.03	122.93	119.17
2	C	501	PLP	C4A-C4-C3	-2.03	117.06	120.50

There are no chirality outliers.

All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	502	HCS	O-C-CA-N
3	A	502	HCS	CA-CB-CG-SD
3	A	502	HCS	OXT-C-CA-N
3	A	502	HCS	C-CA-CB-CG
3	A	502	HCS	N-CA-CB-CG


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	501	PLP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	396/417 (94%)	0.36	41 (10%) 6 5	16, 25, 55, 81	0
1	C	396/417 (94%)	0.50	50 (12%) 3 3	17, 26, 59, 100	0
All	All	792/834 (94%)	0.43	91 (11%) 4 4	16, 25, 59, 100	0

All (91) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	387	VAL	12.0
1	C	374	PHE	11.7
1	C	380	LEU	11.3
1	C	377	ALA	10.5
1	A	387	VAL	8.6
1	A	377	ALA	8.0
1	C	385	LEU	7.7
1	C	378	ASN	7.7
1	C	379	MET	7.7
1	A	391	ASP	7.2
1	A	380	LEU	7.2
1	C	131	CYS	7.0
1	A	379	MET	6.9
1	A	131	CYS	6.3
1	A	378	ASN	6.2
1	C	373	VAL	6.0
1	C	391	ASP	5.7
1	A	374	PHE	5.7
1	C	376	HIS	5.6
1	C	22	VAL	5.5
1	C	375	THR	5.5
1	C	386	LYS	5.2
1	C	132	SER	5.2
1	C	372	ALA	5.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	385	LEU	5.0
1	C	388	GLY	4.9
1	A	372	ALA	4.9
1	A	375	THR	4.9
1	A	383	ASP	4.7
1	C	23	LYS	4.6
1	C	383	ASP	4.6
1	C	370	CYS	4.2
1	C	141	ARG	4.1
1	A	390	THR	4.1
1	A	392	GLY	4.1
1	A	133	TYR	4.0
1	C	389	ILE	4.0
1	A	386	LYS	4.0
1	C	341	THR	3.9
1	C	390	THR	3.7
1	C	381	ARG	3.6
1	A	381	ARG	3.5
1	C	384	ARG	3.5
1	C	137	ASN	3.5
1	A	153	PHE	3.5
1	C	134	GLY	3.4
1	A	345	ILE	3.3
1	A	384	ARG	3.3
1	C	133	TYR	3.3
1	A	137	ASN	3.2
1	C	142	LEU	3.2
1	A	142	LEU	3.1
1	A	373	VAL	3.1
1	C	138	ARG	3.1
1	C	392	GLY	3.1
1	A	168	PRO	3.0
1	A	388	GLY	3.0
1	A	376	HIS	2.9
1	C	382	GLU	2.8
1	C	68	VAL	2.7
1	C	193	GLN	2.7
1	A	68	VAL	2.6
1	C	69	GLU	2.6
1	C	55	PRO	2.6
1	A	141	ARG	2.6
1	C	393	PHE	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	417	ALA	2.6
1	A	382	GLU	2.6
1	C	345	ILE	2.6
1	C	302	HIS	2.5
1	C	271	PHE	2.5
1	C	306	HIS	2.5
1	A	57	ILE	2.4
1	A	132	SER	2.4
1	A	69	GLU	2.4
1	A	341	THR	2.4
1	C	340	GLY	2.3
1	A	275	LEU	2.3
1	C	371	PRO	2.3
1	A	342	GLU	2.3
1	A	271	PHE	2.2
1	A	389	ILE	2.2
1	A	239	LEU	2.1
1	A	136	THR	2.1
1	C	343	ALA	2.1
1	C	337	VAL	2.1
1	C	70	ASN	2.1
1	A	160	THR	2.1
1	A	371	PRO	2.0
1	C	57	ILE	2.0
1	C	270	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	HCS	A	502	8/8	0.84	0.14	35,60,62,85	0
2	PLP	A	501	15/16	0.98	0.09	18,20,24,30	0
2	PLP	C	501	15/16	0.98	0.08	18,21,29,31	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around PLP A 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around PLP C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.