



wwPDB EM Validation Summary Report ⓘ

Nov 20, 2022 – 06:01 pm GMT

PDB ID : 4BON
EMDB ID : EMD-2378
Title : The structure and super-organization of acetylcholine receptor-rapsyn complexes class B
Authors : Zuber, B.; Unwin, N.
Deposited on : 2013-05-21
Resolution : 40.00 Å(reported)
Based on initial model : 2BG9

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.2


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 40.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	461	15% 6% 51% 20% 20%
1	D	461	14% 7% 51% 21% 20%
2	B	493	7% 5% 50% 19% 25%
3	C	522	11% 7% 43% 19% 29%
4	E	505	7% 6% 47% 18% 27%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 14924 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ACETYLCHOLINE RECEPTOR SUBUNIT ALPHA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	370	Total	C	N	O	S	0	0
			2991	1954	478	540	19		
1	D	370	Total	C	N	O	S	0	0
			2991	1954	478	540	19		

- Molecule 2 is a protein called ACETYLCHOLINE RECEPTOR BETA SUBUNIT.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	370	Total	C	N	O	S	0	0
			2972	1938	465	554	15		

- Molecule 3 is a protein called ACETYLCHOLINE RECEPTOR DELTA SUBUNIT.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	C	370	Total	C	N	O	S	0	1
			2983	1944	489	536	14		


- Molecule 4 is a protein called ACETYLCHOLINE RECEPTOR GAMMA SUBUNIT.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	E	371	Total	C	N	O	S	0	1
			2987	1948	478	551	10		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ACETYLCHOLINE RECEPTOR SUBUNIT ALPHA





- Molecule 1: ACETYLCHOLINE RECEPTOR SUBUNIT ALPHA






● Molecule 2: ACETYLCHOLINE RECEPTOR BETA SUBUNIT





● Molecule 3: ACETYLCHOLINE RECEPTOR DELTA SUBUNIT



● Molecule 4: ACETYLCHOLINE RECEPTOR GAMMA SUBUNIT



E44	Y104	G164	N224	K284	GLU	ASP	A464
K45	A105	GLU	I225	Y285	LYS	LEU	I465
E46	N106	VAL	I226	L286	PRO	ALA	F466
E47	V107	VAL	A227	I287	GLN	ASN	G469
A48	L108	GLU	P228	F288	PRO	PHE	H470
L49	V109	TRP	C229	V289	ARG	ALA	L471
T50	Y110	ILE	V230	M290	ARG	PRO	H472
T51	M111	HIS	I231	F291	ARG	GLU	M473
N52	D112	HIS	L232	V292	SER	ILE	Q473
V53	G113	I172	I233	L293	SER	LYS	W474
W54	G114	D173	S233	S293	PHE	S414	P475
I55	S114	P174	S234	L294	GLY	C415	W476
E56	M115	P175	L235	V295	ILE	W416	E477
I57	Y116	E175	V236	I296	MET	E417	F477
Q58	W117	D176	V237	V297	ILE	A418	PRO
Q59	L118	T178	L238	T298	ILE	C419	PHE
N60	P119	E179	V239	N299	LYS	N420	PRO
D61	P120	W180	Y240	C300	ALA	F421	PRO
Y62	A121	N180	F241	V301	GLU	I422	GLY
R63	I122	G181	L242	I302	GLU	A423	ASP
L64	Y123	W183	P243	V303	TYR	K424	PRO
S65	R124	T184	A244	L304	ILE	S425	ARG
W66	S125	I185	Q245	N305	LEU	S426	LYS
M67	T126	I186	A246	V306	LYS	T426	TYR
T68	C127	H187	G247	S307	LYS	K427	VAL
S69	P128	R188	G248	L308	PRO	E428	PRO
E70	L129	P189	Q249	R309	ARG	Q429	
Y71	A130	A190	K250	T310	SER	H430	
E72	V131	K191	C251	P311	GLU	D431	
G73	T132	K192	T252	N312	LEU	S432	
I74	Y133	Y193	L253	T313	MET	C433	
D75	F134	Y194	S254	H314	PHE	S434	
L76	F135	N195	I255	SER	GLU	E435	
W77	F136	W196	S256	LEU	GLU	N436	
R78	D137	Q197	V257	SER	GLN	E437	
I79	W138	L198	L258	GLU	LYS	N438	
P80	Q139	T199	L259	LYS	ASP	W439	
S81	M140	K200	A260	ILE	ARG	V440	
E82	C141	D201	T262	LYS	HIS	L441	
L83	S142	D202	I263	LEU	GLY	I442	
L84	L143	L203	F264	PHE	LYS	C443	
W85	V144	D204	L265	ARG	ARG	K444	
P87	F145	F205	Q206	GLU	VAL	V445	
V88	R146	Q207	L267	PHE	ASN	I446	
V90	S147	E207	I268	LEU	LYS	D447	
L91	Q148	I208	A269	PRO	MET	K448	
E92	T149	I209	Q270	LYS	THR	A449	
N93	Y150	F210	K271	TYR	SER	C450	
N94	M151	L211	V272	LEU	ASP	F451	
V95	A152	L212	P273	GLY	ILE	W452	
D96	H153	I213	E274	MET	ILE	I453	
G97	E154	I214	T275	HIS	GLY	A454	
Q98	V155	Q215	S276	LEU	THR	L455	
F99	M156	R216	K277	GLU	THR	L456	
E100	L157	P218	V278	PRO	VAL	L457	
V101	Q158	L219	V279	SER	ASP	F458	
A102	S160	F220	P280	GLU	LEU	S459	
E162	A161	Y221	L281	GLU	TYR	L460	
E163	E162	I222	I282	THR	LYS	G461	
		I223	G283	PRO		T462	
						L463	

4 Experimental information

Property	Value	Source
EM reconstruction method	TOMOGRAPHY	Depositor
Imposed symmetry	POINT, Not provided	
Number of tilted images used	3564	Depositor
Resolution determination method	Not provided	
CTF correction method	Not provided	
Microscope	FEI TECNAI F30	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	3000	Depositor
Maximum defocus (nm)	6000	Depositor
Magnification	80213	Depositor
Image detector	GATAN ULTRASCAN 4000 (4k x 4k)	Depositor
Maximum voxel value	1.798	Depositor
Minimum voxel value	-0.712	Depositor
Average voxel value	0.000	Depositor
Voxel value standard deviation	0.080	Depositor
Recommended contour level	0.362	Depositor
Tomogram size (\AA)	448.8, 448.8, 448.8	wwPDB
Tomogram dimensions	60, 60, 60	wwPDB
Tomogram angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Grid spacing (\AA)	7.48, 7.48, 7.48	Depositor

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	3/3069 (0.1%)	1.03	10/4186 (0.2%)
1	D	0.74	2/3069 (0.1%)	1.01	6/4186 (0.1%)
2	B	0.76	2/3048 (0.1%)	0.99	4/4162 (0.1%)
3	C	0.74	2/3059 (0.1%)	1.03	9/4175 (0.2%)
4	E	0.73	6/3057 (0.2%)	1.01	9/4174 (0.2%)
All	All	0.74	15/15302 (0.1%)	1.01	38/20883 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	2
3	C	0	2
All	All	0	4

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	129	THR	C-N	-8.42	1.14	1.34
1	A	118	TRP	CB-CG	7.92	1.64	1.50
1	D	208	GLN	C-N	7.56	1.51	1.34
4	E	8	GLU	CB-CG	6.52	1.64	1.52
3	C	265	LEU	C-N	6.19	1.48	1.34

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	266	ALA	N-CA-CB	10.39	124.65	110.10
4	E	198	LEU	CA-CB-CG	7.20	131.86	115.30
3	C	315	ARG	NE-CZ-NH2	7.16	123.88	120.30
1	A	209	ARG	NE-CZ-NH2	7.10	123.85	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	263	ILE	CG1-CB-CG2	-6.67	96.73	111.40

There are no chirality outliers.

All (4) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	C	63	TYR	Sidechain
3	C	74	TYR	Sidechain
1	D	277	TYR	Sidechain
1	D	72	TYR	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2991	0	3005	1066	0
1	D	2991	0	3006	1057	0
2	B	2972	0	2953	1081	0
3	C	2983	0	2987	1155	0
4	E	2987	0	2994	1100	0
All	All	14924	0	14945	5204	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 174.

The worst 5 of 5204 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:E:183:TRP:CB	4:E:216:ARG:HG2	1.33	1.56
2:B:134:TYR:CE1	2:B:213:ILE:HG13	1.44	1.49
1:A:167:LEU:HD12	1:A:178:MET:CB	1.43	1.46
1:A:167:LEU:CD1	1:A:178:MET:HB2	1.46	1.44
3:C:316:THR:CG2	3:C:447:ASN:HB3	1.53	1.36

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	366/461 (79%)	289 (79%)	49 (13%)	28 (8%)	1	13
1	D	366/461 (79%)	294 (80%)	41 (11%)	31 (8%)	1	12
2	B	364/493 (74%)	273 (75%)	59 (16%)	32 (9%)	1	11
3	C	364/522 (70%)	288 (79%)	58 (16%)	18 (5%)	2	20
4	E	365/505 (72%)	281 (77%)	58 (16%)	26 (7%)	1	14
All	All	1825/2442 (75%)	1425 (78%)	265 (14%)	135 (7%)	2	14

5 of 135 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	2	GLU
1	A	27	HIS
1	A	76	LYS
1	A	83	ASP
1	A	102	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	343/427 (80%)	248 (72%)	95 (28%)	0	3
1	D	343/427 (80%)	258 (75%)	85 (25%)	0	3
2	B	340/449 (76%)	262 (77%)	78 (23%)	1	4
3	C	335/475 (70%)	243 (72%)	92 (28%)	0	3

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	E	337/463 (73%)	250 (74%)	87 (26%)	0	3
All	All	1698/2241 (76%)	1261 (74%)	437 (26%)	2	3

5 of 437 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	C	274	THR
1	D	94	ASN
4	E	184	THR
3	C	296	MET
3	C	471	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
3	C	200	ASN
4	E	156	ASN
1	D	42	ASN
4	E	153	HIS
4	E	206	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	B	1


All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	B	129:THR	C	130:ILE	N	1.14

6 Tomogram visualisation [i](#)


This section contains visualisations of the EMDB entry EMD-2378. These allow visual inspection of the internal detail of the tomogram and identification of artifacts.

6.1 Orthogonal projections [i](#)




The images above show the tomogram projected in three orthogonal directions.

6.2 Central slices [i](#)




The images above show central slices of the tomogram in three orthogonal directions.

6.3 Largest variance slices [i](#)



X Index: 33



Y Index: 30



Z Index: 22

The images above show the largest variance slices of the tomogram in three orthogonal directions.


6.4 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Tomogram analysis [i](#)

This section contains the results of statistical analysis of the tomogram.

7.1 Voxel-value distribution [i](#)



The voxel-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic.


8 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-2378 and PDB model 4BON. Per-residue inclusion information can be found in section 3 on page 4.

8.1 Map-model overlay [i](#)

This section was not generated.

8.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

8.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.













8.4 Atom inclusion [i](#)



At the recommended contour level, 86% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

8.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.362) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8462	 0.0420
A	 0.8042	 0.0400
B	 0.8805	 0.0400
C	 0.8374	 0.0420
D	 0.8110	 0.0360
E	 0.8981	 0.0540

