



# wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 9, 2024 – 12:45 pm GMT

PDB ID : 5BPU  
Title : Crystal structure of Norrin, a Wnt signalling activator, Crystal Form I  
Authors : Chang, T.-H.; Hsieh, F.-L.; Harlos, K.; Jones, E.Y.  
Deposited on : 2015-05-28  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	122	 14% 74% 16% • 9%
1	B	122	 10% 72% 9% 19%
1	C	122	 20% 67% 16% • 16%
1	D	122	 11% 76% 7% • 16%
1	E	122	 16% 59% 19% • 18%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	122	
2	H	7	
3	I	4	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GGL	H	1	X	-	-	-
3	GGL	I	1	X	-	-	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5087 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Norrin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	111	Total 866	C 526	N 166	O 159	S 15	0	0	0
1	B	99	Total 782	C 477	N 153	O 138	S 14	0	0	0
1	C	103	Total 807	C 490	N 157	O 146	S 14	0	0	0
1	D	102	Total 804	C 489	N 156	O 144	S 15	0	0	0
1	E	100	Total 788	C 480	N 154	O 140	S 14	0	0	0
1	F	99	Total 782	C 477	N 153	O 138	S 14	0	0	0

There are 78 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	22	GLY	-	expression tag	UNP Q00604
A	23	PRO	-	expression tag	UNP Q00604
A	24	GLY	-	expression tag	UNP Q00604
A	134	GLY	-	expression tag	UNP Q00604
A	135	THR	-	expression tag	UNP Q00604
A	136	GLU	-	expression tag	UNP Q00604
A	137	THR	-	expression tag	UNP Q00604
A	138	SER	-	expression tag	UNP Q00604
A	139	GLN	-	expression tag	UNP Q00604
A	140	VAL	-	expression tag	UNP Q00604
A	141	ALA	-	expression tag	UNP Q00604
A	142	PRO	-	expression tag	UNP Q00604
A	143	ALA	-	expression tag	UNP Q00604
B	22	GLY	-	expression tag	UNP Q00604
B	23	PRO	-	expression tag	UNP Q00604
B	24	GLY	-	expression tag	UNP Q00604
B	134	GLY	-	expression tag	UNP Q00604

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	135	THR	-	expression tag	UNP Q00604
B	136	GLU	-	expression tag	UNP Q00604
B	137	THR	-	expression tag	UNP Q00604
B	138	SER	-	expression tag	UNP Q00604
B	139	GLN	-	expression tag	UNP Q00604
B	140	VAL	-	expression tag	UNP Q00604
B	141	ALA	-	expression tag	UNP Q00604
B	142	PRO	-	expression tag	UNP Q00604
B	143	ALA	-	expression tag	UNP Q00604
C	22	GLY	-	expression tag	UNP Q00604
C	23	PRO	-	expression tag	UNP Q00604
C	24	GLY	-	expression tag	UNP Q00604
C	134	GLY	-	expression tag	UNP Q00604
C	135	THR	-	expression tag	UNP Q00604
C	136	GLU	-	expression tag	UNP Q00604
C	137	THR	-	expression tag	UNP Q00604
C	138	SER	-	expression tag	UNP Q00604
C	139	GLN	-	expression tag	UNP Q00604
C	140	VAL	-	expression tag	UNP Q00604
C	141	ALA	-	expression tag	UNP Q00604
C	142	PRO	-	expression tag	UNP Q00604
C	143	ALA	-	expression tag	UNP Q00604
D	22	GLY	-	expression tag	UNP Q00604
D	23	PRO	-	expression tag	UNP Q00604
D	24	GLY	-	expression tag	UNP Q00604
D	134	GLY	-	expression tag	UNP Q00604
D	135	THR	-	expression tag	UNP Q00604
D	136	GLU	-	expression tag	UNP Q00604
D	137	THR	-	expression tag	UNP Q00604
D	138	SER	-	expression tag	UNP Q00604
D	139	GLN	-	expression tag	UNP Q00604
D	140	VAL	-	expression tag	UNP Q00604
D	141	ALA	-	expression tag	UNP Q00604
D	142	PRO	-	expression tag	UNP Q00604
D	143	ALA	-	expression tag	UNP Q00604
E	22	GLY	-	expression tag	UNP Q00604
E	23	PRO	-	expression tag	UNP Q00604
E	24	GLY	-	expression tag	UNP Q00604
E	134	GLY	-	expression tag	UNP Q00604
E	135	THR	-	expression tag	UNP Q00604
E	136	GLU	-	expression tag	UNP Q00604
E	137	THR	-	expression tag	UNP Q00604

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	138	SER	-	expression tag	UNP Q00604
E	139	GLN	-	expression tag	UNP Q00604
E	140	VAL	-	expression tag	UNP Q00604
E	141	ALA	-	expression tag	UNP Q00604
E	142	PRO	-	expression tag	UNP Q00604
E	143	ALA	-	expression tag	UNP Q00604
F	22	GLY	-	expression tag	UNP Q00604
F	23	PRO	-	expression tag	UNP Q00604
F	24	GLY	-	expression tag	UNP Q00604
F	134	GLY	-	expression tag	UNP Q00604
F	135	THR	-	expression tag	UNP Q00604
F	136	GLU	-	expression tag	UNP Q00604
F	137	THR	-	expression tag	UNP Q00604
F	138	SER	-	expression tag	UNP Q00604
F	139	GLN	-	expression tag	UNP Q00604
F	140	VAL	-	expression tag	UNP Q00604
F	141	ALA	-	expression tag	UNP Q00604
F	142	PRO	-	expression tag	UNP Q00604
F	143	ALA	-	expression tag	UNP Q00604

- Molecule 2 is a protein called (GGL)EEEEEE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	H	7	59	35	7	17	0	0	0

- Molecule 3 is a protein called (GGL)EEE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	I	4	35	20	4	11	0	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	28	28	28	0	0
4	B	29	29	29	0	0
4	C	27	27	27	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	D	22	Total 22	O 22	0	0
4	E	31	Total 31	O 31	0	0
4	F	25	Total 25	O 25	0	0
4	H	2	Total 2	O 2	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Norrin




- Molecule 1: Norrin



- Molecule 1: Norrin




- Molecule 1: Norrin




- Molecule 1: Norrin







• Molecule 1: Norrin



• Molecule 2: (GGL)EEEEEE



• Molecule 3: (GGL)EEE



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.37Å 79.10Å 234.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	65.56 – 2.40 65.56 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (65.56-2.40) 99.8 (65.56-2.40)	Depositor EDS
$R_{merge}$	0.13	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.71 (at 2.40Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.217 , 0.263 0.240 , 0.238	Depositor DCC
$R_{free}$ test set	1717 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.3	Xtrriage
Anisotropy	0.214	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 57.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	5087	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	69.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GGL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.51	2/882 (0.2%)	0.56	1/1184 (0.1%)
1	B	0.27	0/798	0.48	0/1070
1	C	0.50	0/823	0.55	0/1104
1	D	0.28	0/820	0.46	0/1099
1	E	0.41	0/804	0.54	0/1078
1	F	0.37	0/798	0.57	1/1070 (0.1%)
2	H	1.03	0/44	1.28	0/44
3	I	1.01	0/23	1.19	0/23
All	All	0.42	2/4992 (0.0%)	0.54	2/6672 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	2
1	F	0	1
2	H	1	0
3	I	1	0
All	All	2	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	88	PRO	N-CD	5.19	1.55	1.47
1	A	36	PRO	N-CD	5.18	1.55	1.47

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	35	ASP	C-N-CD	6.22	141.46	128.40
1	A	35	ASP	C-N-CD	5.29	139.50	128.40

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	H	1	GGL	CA
3	I	1	GGL	CA

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	82	SER	Peptide
1	E	84	VAL	Peptide
1	F	85	LEU	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	866	0	850	9	0
1	B	782	0	771	3	0
1	C	807	0	790	15	0
1	D	804	0	789	3	0
1	E	788	0	776	14	0
1	F	782	0	771	9	0
2	H	59	0	37	1	0
3	I	35	0	22	0	0
4	A	28	0	0	1	0
4	B	29	0	0	1	0
4	C	27	0	0	1	0
4	D	22	0	0	0	0
4	E	31	0	0	1	0
4	F	25	0	0	1	0
4	H	2	0	0	0	0
All	All	5087	0	4806	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 48 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:35:ASP:O	1:F:38:ARG:NH1	1.87	1.06
1:C:110:CYS:HB2	1:C:116:LEU:HD13	1.56	0.86
1:A:87:GLN:N	1:A:87:GLN:OE1	2.30	0.65
1:E:76:GLU:OE1	1:E:90:ARG:NH1	2.29	0.65
1:C:116:LEU:HD12	1:C:116:LEU:N	2.14	0.63

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	109/122 (89%)	107 (98%)	2 (2%)	0	100	100
1	B	97/122 (80%)	95 (98%)	2 (2%)	0	100	100
1	C	101/122 (83%)	99 (98%)	2 (2%)	0	100	100
1	D	100/122 (82%)	99 (99%)	1 (1%)	0	100	100
1	E	98/122 (80%)	94 (96%)	4 (4%)	0	100	100
1	F	97/122 (80%)	92 (95%)	5 (5%)	0	100	100
All	All	602/732 (82%)	586 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	102/110 (93%)	94 (92%)	8 (8%)	10	17
1	B	92/110 (84%)	85 (92%)	7 (8%)	11	18
1	C	95/110 (86%)	86 (90%)	9 (10%)	7	11
1	D	95/110 (86%)	89 (94%)	6 (6%)	15	25
1	E	93/110 (84%)	83 (89%)	10 (11%)	5	7
1	F	92/110 (84%)	77 (84%)	15 (16%)	2	2
2	H	6/6 (100%)	5 (83%)	1 (17%)	2	2
3	I	3/3 (100%)	3 (100%)	0	100	100
All	All	578/669 (86%)	522 (90%)	56 (10%)	6	10

5 of 56 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	119	THR
2	H	7	GLU
1	E	115	ARG
1	F	127	HIS
1	F	86	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GGL	H	1	2	7,8,9	0.93	0	7,9,11	0.77	0
3	GGL	I	1	3	7,8,9	0.82	0	7,9,11	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GGL	H	1	2	1/1/2/3	6/7/8/9	-
3	GGL	I	1	3	1/1/2/3	0/7/8/9	-

There are no bond length outliers.

There are no bond angle outliers.

All (2) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	H	1	GGL	CA
3	I	1	GGL	CA

5 of 6 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	H	1	GGL	O-C-CA-N
2	H	1	GGL	C-CA-CB-CG
2	H	1	GGL	OXT-C-CA-N
2	H	1	GGL	N-CA-CB-CG
2	H	1	GGL	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	H	5
3	I	2

The worst 5 of 7 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	H	5:GLU	C	6:GLU	N	6.13
1	H	2:GLU	C	3:GLU	N	5.99
1	H	6:GLU	C	7:GLU	N	5.81
1	H	4:GLU	C	5:GLU	N	5.62
1	I	3:GLU	C	4:GLU	N	5.46



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	111/122 (90%)	0.67	17 (15%) 6 6	29, 56, 116, 141	0
1	B	99/122 (81%)	0.73	12 (12%) 10 8	32, 58, 103, 132	0
1	C	103/122 (84%)	1.14	24 (23%) 2 2	35, 66, 123, 147	0
1	D	102/122 (83%)	0.62	14 (13%) 8 7	36, 61, 96, 116	0
1	E	100/122 (81%)	1.13	20 (20%) 3 3	35, 62, 128, 144	0
1	F	99/122 (81%)	1.11	15 (15%) 6 6	40, 69, 143, 189	0
2	H	6/7 (85%)	2.47	4 (66%) 0 0	60, 72, 97, 130	0
3	I	3/4 (75%)	2.51	2 (66%) 0 0	78, 78, 78, 98	0
All	All	623/743 (83%)	0.92	108 (17%) 5 5	29, 64, 121, 189	0

The worst 5 of 108 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	141	ALA	5.7
1	A	31	ILE	4.9
1	A	84	VAL	4.7
1	F	83	THR	4.5
1	E	34	SER	4.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GGL	H	1	9/10	0.78	0.33	98,109,154,156	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	GGL	I	1	9/10	0.78	0.28	76,98,127,140	0

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.