



# wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 28, 2024 – 04:53 pm BST

PDB ID : 4BQ2  
Title : Structural analysis of an exo-beta-agarase  
Authors : Pluinage, B.; Hehemann, J.H.; Boraston, A.B.  
Deposited on : 2013-05-29  
Resolution : 1.90 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*





The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	750	
1	B	750	
1	C	750	
1	D	750	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	C	1795	-	-	X	-

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 26607 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called B-AGARASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	746	6012	3838	1007	1143	24	0	10	0
1	B	747	6027	3841	1005	1157	24	0	12	0
1	C	748	6028	3840	1011	1154	23	0	12	0
1	D	726	5850	3730	981	1116	23	1	11	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	GLY	-	expression tag	UNP Q21HC5
A	45	SER	-	expression tag	UNP Q21HC5
A	46	HIS	-	expression tag	UNP Q21HC5
B	44	GLY	-	expression tag	UNP Q21HC5
B	45	SER	-	expression tag	UNP Q21HC5
B	46	HIS	-	expression tag	UNP Q21HC5
C	44	GLY	-	expression tag	UNP Q21HC5
C	45	SER	-	expression tag	UNP Q21HC5
C	46	HIS	-	expression tag	UNP Q21HC5
D	44	GLY	-	expression tag	UNP Q21HC5
D	45	SER	-	expression tag	UNP Q21HC5
D	46	HIS	-	expression tag	UNP Q21HC5

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0
2	D	1	Total C O 6 3 3	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0
3	B	1	Total Ca 1 1	0	0
3	C	1	Total Ca 1 1	0	0
3	D	1	Total Ca 1 1	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	645	Total O 645 645	0	0
4	B	655	Total O 655 655	0	0
4	C	651	Total O 651 651	0	0
4	D	603	Total O 603 603	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: B-AGARASE



#### • Molecule 1: B-AGARASE






#### • Molecule 1: B-AGARASE





● Molecule 1: B-AGARASE



SER  
LYS



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	166.67Å 166.67Å 114.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.28 – 1.90 39.28 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.6 (39.28-1.90) 99.7 (39.28-1.90)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.44 (at 1.89Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, $R_{free}$	0.170 , 0.215 0.170 , 0.214	Depositor DCC
$R_{free}$ test set	12312 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	21.6	Xtrriage
Anisotropy	0.034	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 52.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.44$ , $\langle L^2 \rangle = 0.27$	Xtrriage
Estimated twinning fraction	0.038 for h,-k,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	26607	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 53.49 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.1852e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.62	0/6187	0.74	3/8409 (0.0%)
1	B	0.63	0/6202	0.74	3/8430 (0.0%)
1	C	0.64	0/6202	0.74	4/8431 (0.0%)
1	D	0.63	1/6017 (0.0%)	0.72	3/8173 (0.0%)
All	All	0.63	1/24608 (0.0%)	0.73	13/33443 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	658	LYS	CE-NZ	-11.95	1.19	1.49

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	471	ARG	NE-CZ-NH2	-12.94	113.83	120.30
1	A	471	ARG	NE-CZ-NH1	10.75	125.67	120.30
1	B	471	ARG	NE-CZ-NH2	-9.50	115.55	120.30
1	C	471	ARG	NE-CZ-NH2	-8.86	115.87	120.30
1	B	471	ARG	NE-CZ-NH1	8.60	124.60	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6012	0	5718	59	0
1	B	6027	0	5702	52	0
1	C	6028	0	5721	49	0
1	D	5850	0	5528	73	0
2	A	36	0	48	4	0
2	B	48	0	64	5	0
2	C	30	0	40	7	1
2	D	18	0	24	1	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	645	0	0	4	1
4	B	655	0	0	6	0
4	C	651	0	0	11	0
4	D	603	0	0	9	0
All	All	26607	0	22845	238	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 238 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:786:TYR:O	1:B:789:ARG:O	1.73	1.05
1:C:440:ASP:OD1	1:C:443[A]:ARG:NH2	1.90	1.03
1:A:752:ARG:HH12	2:A:1793:GOL:H31	1.23	1.03
1:A:623:ASP:HB3	4:A:2539:HOH:O	1.57	1.03
1:A:721:GLU:HG3	4:A:2610:HOH:O	1.60	0.99

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:1795:GOL:O1	4:A:2023:HOH:O[1_556]	1.86	0.34

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	754/750 (100%)	731 (97%)	23 (3%)	0	100	100
1	B	757/750 (101%)	734 (97%)	23 (3%)	0	100	100
1	C	758/750 (101%)	734 (97%)	24 (3%)	0	100	100
1	D	729/750 (97%)	702 (96%)	25 (3%)	2 (0%)	41	31
All	All	2998/3000 (100%)	2901 (97%)	95 (3%)	2 (0%)	51	42

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	755	ASP
1	D	184	ASN

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	637/630 (101%)	630 (99%)	7 (1%)	73	73
1	B	639/630 (101%)	631 (99%)	8 (1%)	69	68
1	C	640/630 (102%)	630 (98%)	10 (2%)	62	60
1	D	615/630 (98%)	605 (98%)	10 (2%)	62	60
All	All	2531/2520 (100%)	2496 (99%)	35 (1%)	69	65

5 of 35 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	182	ARG
1	D	183	SER
1	D	406	ARG
1	B	697	HIS
1	B	581	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 36 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	184	ASN
1	D	760	ASN
1	D	194	GLN
1	D	640	HIS
1	B	409	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 4 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	C	1794	-	5,5,5	0.37	0	5,5,5	0.60	0
2	GOL	C	1796	-	5,5,5	0.40	0	5,5,5	0.65	0
2	GOL	B	1793	-	5,5,5	0.74	0	5,5,5	0.52	0
2	GOL	D	1793	-	5,5,5	0.46	0	5,5,5	0.79	0
2	GOL	A	1796	-	5,5,5	0.28	0	5,5,5	0.37	0
2	GOL	B	1797	-	5,5,5	0.31	0	5,5,5	0.70	0
2	GOL	B	1792	-	5,5,5	0.55	0	5,5,5	0.21	0
2	GOL	C	1793	-	5,5,5	0.43	0	5,5,5	0.64	0
2	GOL	A	1792	-	5,5,5	0.86	0	5,5,5	1.03	0
2	GOL	B	1798	-	5,5,5	0.19	0	5,5,5	0.28	0
2	GOL	A	1794	-	5,5,5	0.21	0	5,5,5	0.71	0
2	GOL	D	1794	-	5,5,5	0.48	0	5,5,5	0.69	0
2	GOL	B	1791	-	5,5,5	0.23	0	5,5,5	1.20	1 (20%)
2	GOL	C	1797	-	5,5,5	0.14	0	5,5,5	0.73	0
2	GOL	B	1796	-	5,5,5	0.31	0	5,5,5	0.27	0
2	GOL	A	1793	-	5,5,5	0.61	0	5,5,5	0.74	0
2	GOL	B	1794	-	5,5,5	0.43	0	5,5,5	1.45	1 (20%)
2	GOL	A	1795	-	5,5,5	0.48	0	5,5,5	0.39	0
2	GOL	D	1792	-	5,5,5	0.20	0	5,5,5	0.53	0
2	GOL	B	1795	-	5,5,5	0.39	0	5,5,5	0.33	0
2	GOL	C	1795	-	5,5,5	0.34	0	5,5,5	1.03	0
2	GOL	A	1791	-	5,5,5	0.37	0	5,5,5	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	C	1794	-	-	0/4/4/4	-
2	GOL	C	1796	-	-	0/4/4/4	-
2	GOL	B	1793	-	-	0/4/4/4	-
2	GOL	D	1793	-	-	1/4/4/4	-
2	GOL	A	1796	-	-	0/4/4/4	-
2	GOL	B	1797	-	-	2/4/4/4	-
2	GOL	B	1792	-	-	0/4/4/4	-
2	GOL	C	1793	-	-	0/4/4/4	-
2	GOL	A	1792	-	-	2/4/4/4	-
2	GOL	B	1798	-	-	2/4/4/4	-
2	GOL	A	1794	-	-	0/4/4/4	-
2	GOL	D	1794	-	-	4/4/4/4	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	1791	-	-	2/4/4/4	-
2	GOL	C	1797	-	-	0/4/4/4	-
2	GOL	B	1796	-	-	2/4/4/4	-
2	GOL	A	1793	-	-	2/4/4/4	-
2	GOL	B	1794	-	-	2/4/4/4	-
2	GOL	A	1795	-	-	0/4/4/4	-
2	GOL	D	1792	-	-	4/4/4/4	-
2	GOL	B	1795	-	-	2/4/4/4	-
2	GOL	C	1795	-	-	2/4/4/4	-
2	GOL	A	1791	-	-	2/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1794	GOL	O1-C1-C2	-2.27	99.30	110.20
2	B	1791	GOL	C3-C2-C1	-2.14	103.37	111.70

There are no chirality outliers.

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1791	GOL	C1-C2-C3-O3
2	A	1791	GOL	O2-C2-C3-O3
2	A	1792	GOL	C1-C2-C3-O3
2	B	1791	GOL	C1-C2-C3-O3
2	B	1795	GOL	O1-C1-C2-C3

There are no ring outliers.

11 monomers are involved in 18 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	1796	GOL	1	0
2	B	1793	GOL	1	0
2	D	1793	GOL	1	0
2	B	1797	GOL	1	0
2	A	1792	GOL	1	0
2	B	1798	GOL	1	0
2	C	1797	GOL	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1793	GOL	3	0
2	B	1794	GOL	1	0
2	B	1795	GOL	1	0
2	C	1795	GOL	5	1

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	746/750 (99%)	-0.17	8 (1%) 80 82	12, 21, 37, 57	0
1	B	747/750 (99%)	-0.18	11 (1%) 73 76	12, 20, 38, 54	0
1	C	748/750 (99%)	-0.10	20 (2%) 54 57	12, 21, 41, 72	0
1	D	726/750 (96%)	-0.03	33 (4%) 33 36	12, 22, 51, 81	1 (0%)
All	All	2967/3000 (98%)	-0.12	72 (2%) 59 62	12, 21, 41, 81	1 (0%)

The worst 5 of 72 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	354	SER	6.1
1	C	167	PRO	5.2
1	C	169	SER	5.1
1	D	404	TYR	5.0
1	C	172	VAL	5.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	C	1795	6/6	0.78	0.22	39,45,47,53	0
2	GOL	B	1797	6/6	0.83	0.32	50,51,57,65	0
2	GOL	C	1797	6/6	0.87	0.21	49,52,54,54	0
2	GOL	D	1794	6/6	0.87	0.27	40,43,44,47	0
2	GOL	B	1795	6/6	0.88	0.25	36,41,42,45	0
2	GOL	A	1796	6/6	0.89	0.19	48,49,55,60	0
2	GOL	A	1791	6/6	0.90	0.12	35,36,38,41	0
2	GOL	D	1792	6/6	0.90	0.16	39,46,50,52	0
2	GOL	B	1796	6/6	0.90	0.13	44,47,50,50	0
2	GOL	B	1794	6/6	0.91	0.20	38,39,41,47	0
2	GOL	A	1793	6/6	0.92	0.18	23,28,34,40	0
2	GOL	B	1798	6/6	0.92	0.15	47,49,51,53	0
2	GOL	A	1794	6/6	0.93	0.15	32,39,43,43	0
2	GOL	C	1794	6/6	0.93	0.15	26,29,30,31	0
2	GOL	D	1793	6/6	0.93	0.11	27,32,34,38	0
2	GOL	B	1791	6/6	0.93	0.19	26,31,34,38	0
2	GOL	C	1793	6/6	0.94	0.13	24,25,26,29	0
2	GOL	C	1796	6/6	0.94	0.13	19,30,33,39	0
2	GOL	B	1793	6/6	0.95	0.11	19,27,28,33	0
2	GOL	A	1792	6/6	0.95	0.11	19,25,28,31	0
2	GOL	A	1795	6/6	0.96	0.20	20,24,27,27	0
2	GOL	B	1792	6/6	0.96	0.16	19,23,24,25	0
3	CA	B	1799	1/1	0.98	0.17	55,55,55,55	0
3	CA	C	1798	1/1	0.99	0.04	22,22,22,22	0
3	CA	A	1797	1/1	1.00	0.07	22,22,22,22	0
3	CA	D	1795	1/1	1.00	0.03	20,20,20,20	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.