



# wwPDB X-ray Structure Validation Summary Report ⓘ

Jun 18, 2024 – 10:11 AM EDT

PDB ID : 4C2B  
Title : Crystal Structure of High-Affinity von Willebrand Factor A1 domain with Disulfide Mutation in Complex with High Affinity GPIb alpha  
Authors : Blenner, M.A.; Dong, X.; Springer, T.A.  
Deposited on : 2013-08-16  
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 2.37.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.37.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	215	 78% 14% 7%
1	C	215	 75% 17% 7% 2%
1	E	215	 78% 15% 7%
1	G	215	 78% 13% 8% 4%
2	B	291	 74% 16% 9%
2	D	291	 73% 16% 10% 3%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	291	 76% 14% 9%
2	H	291	 77% 12% 10%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	SO4	H	1266	-	-	X	-

## 2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 14768 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VON WILLEBRAND FACTOR.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	199	Total 1603	C 1021	N 287	O 289	S 6	0	0	0
1	C	199	Total 1602	C 1020	N 287	O 289	S 6	0	0	0
1	E	199	Total 1603	C 1021	N 287	O 289	S 6	0	0	0
1	G	197	Total 1585	C 1009	N 283	O 287	S 6	0	0	0

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1263	MET	-	expression tag	UNP P04275
A	1472	HIS	-	expression tag	UNP P04275
A	1473	HIS	-	expression tag	UNP P04275
A	1474	HIS	-	expression tag	UNP P04275
A	1475	HIS	-	expression tag	UNP P04275
A	1476	HIS	-	expression tag	UNP P04275
A	1477	HIS	-	expression tag	UNP P04275
A	1271	CYS	TYR	engineered mutation	UNP P04275
A	1272	ARG	CYS	engineered mutation	UNP P04275
A	1381	ALA	THR	variant	UNP P04275
C	1263	MET	-	expression tag	UNP P04275
C	1472	HIS	-	expression tag	UNP P04275
C	1473	HIS	-	expression tag	UNP P04275
C	1474	HIS	-	expression tag	UNP P04275
C	1475	HIS	-	expression tag	UNP P04275
C	1476	HIS	-	expression tag	UNP P04275
C	1477	HIS	-	expression tag	UNP P04275
C	1271	CYS	TYR	engineered mutation	UNP P04275
C	1272	ARG	CYS	engineered mutation	UNP P04275
C	1381	ALA	THR	variant	UNP P04275
E	1263	MET	-	expression tag	UNP P04275

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	1472	HIS	-	expression tag	UNP P04275
E	1473	HIS	-	expression tag	UNP P04275
E	1474	HIS	-	expression tag	UNP P04275
E	1475	HIS	-	expression tag	UNP P04275
E	1476	HIS	-	expression tag	UNP P04275
E	1477	HIS	-	expression tag	UNP P04275
E	1271	CYS	TYR	engineered mutation	UNP P04275
E	1272	ARG	CYS	engineered mutation	UNP P04275
E	1381	ALA	THR	variant	UNP P04275
G	1263	MET	-	expression tag	UNP P04275
G	1472	HIS	-	expression tag	UNP P04275
G	1473	HIS	-	expression tag	UNP P04275
G	1474	HIS	-	expression tag	UNP P04275
G	1475	HIS	-	expression tag	UNP P04275
G	1476	HIS	-	expression tag	UNP P04275
G	1477	HIS	-	expression tag	UNP P04275
G	1271	CYS	TYR	engineered mutation	UNP P04275
G	1272	ARG	CYS	engineered mutation	UNP P04275
G	1381	ALA	THR	variant	UNP P04275

- Molecule 2 is a protein called PLATELET GLYCOPROTEIN IB ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	264	2069	1333	345	383	8	0	0	0
2	D	262	2055	1326	343	378	8	0	0	0
2	F	264	2069	1333	345	383	8	0	0	0
2	H	261	2048	1321	341	378	8	0	0	0

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	291	GLY	-	expression tag	UNP P07359
B	21	ARG	ASN	engineered mutation	UNP P07359
B	159	ARG	ASN	engineered mutation	UNP P07359
B	233	VAL	GLY	engineered mutation	UNP P07359
B	239	VAL	MET	engineered mutation	UNP P07359
D	291	GLY	-	expression tag	UNP P07359
D	21	ARG	ASN	engineered mutation	UNP P07359

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	159	ARG	ASN	engineered mutation	UNP P07359
D	233	VAL	GLY	engineered mutation	UNP P07359
D	239	VAL	MET	engineered mutation	UNP P07359
F	291	GLY	-	expression tag	UNP P07359
F	21	ARG	ASN	engineered mutation	UNP P07359
F	159	ARG	ASN	engineered mutation	UNP P07359
F	233	VAL	GLY	engineered mutation	UNP P07359
F	239	VAL	MET	engineered mutation	UNP P07359
H	291	GLY	-	expression tag	UNP P07359
H	21	ARG	ASN	engineered mutation	UNP P07359
H	159	ARG	ASN	engineered mutation	UNP P07359
H	233	VAL	GLY	engineered mutation	UNP P07359
H	239	VAL	MET	engineered mutation	UNP P07359

- Molecule 3 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	B	1	Total O S 5 4 1	0	0
3	C	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0
3	E	1	Total O S 5 4 1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	F	1	Total	O	S	0	0
			5	4	1		
3	G	1	Total	O	S	0	0
			5	4	1		
3	G	1	Total	O	S	0	0
			5	4	1		
3	H	1	Total	O	S	0	0
			5	4	1		
3	H	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C<sub>6</sub>H<sub>13</sub>NO<sub>4</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	G	1	Total C O 7 4 3	0	0

- Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	5	Total O 5 5	0	0
6	B	5	Total O 5 5	0	0
6	C	3	Total O 3 3	0	0
6	D	14	Total O 14 14	0	0
6	E	9	Total O 9 9	0	0
6	F	11	Total O 11 11	0	0
6	G	9	Total O 9 9	0	0
6	H	9	Total O 9 9	0	0



### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: VON WILLEBRAND FACTOR

Chain A: 




#### • Molecule 1: VON WILLEBRAND FACTOR

Chain C: 




#### • Molecule 1: VON WILLEBRAND FACTOR


Chain E: 





#### • Molecule 1: VON WILLEBRAND FACTOR

Chain G: 









• Molecule 2: PLATELET GLYCOPROTEIN IB ALPHA CHAIN





• Molecule 2: PLATELET GLYCOPROTEIN IB ALPHA CHAIN



• Molecule 2: PLATELET GLYCOPROTEIN IB ALPHA CHAIN



• Molecule 2: PLATELET GLYCOPROTEIN IB ALPHA CHAIN



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	96.53Å 103.84Å 119.93Å 90.00° 90.05° 90.00°	Depositor
Resolution (Å)	48.27 – 2.80 48.26 – 2.80	Depositor EDS
% Data completeness (in resolution range)	96.6 (48.27-2.80) 96.3 (48.26-2.80)	Depositor EDS
$R_{merge}$	0.59	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.30 (at 2.81Å)	Xtrriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.253 , 0.286 0.408 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.6	Xtrriage
Anisotropy	0.089	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 54.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.310 for h,-k,-l	Xtrriage
Reported twinning fraction	0.480 for h,-k,-l	Depositor
Outliers	5 of 56511 reflections (0.009%)	Xtrriage
$F_o, F_c$ correlation	0.85	EDS
Total number of atoms	14768	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	89.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 35.82 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.4940e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MES, SO4, PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.20	0/1630	0.34	0/2194
1	C	0.20	0/1630	0.33	0/2195
1	E	0.20	0/1630	0.34	0/2194
1	G	0.20	0/1611	0.34	0/2168
2	B	0.20	0/2115	0.37	0/2888
2	D	0.20	0/2100	0.36	0/2866
2	F	0.20	0/2115	0.37	0/2888
2	H	0.21	0/2092	0.36	0/2855
All	All	0.20	0/14923	0.35	0/20248

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1603	0	1658	15	0
1	C	1602	0	1654	23	0
1	E	1603	0	1658	18	0
1	G	1585	0	1640	17	0
2	B	2069	0	2119	29	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	2055	0	2109	26	0
2	F	2069	0	2119	24	0
2	H	2048	0	2099	22	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
3	C	5	0	0	0	0
3	D	5	0	0	0	0
3	E	5	0	0	0	0
3	F	5	0	0	0	0
3	G	10	0	0	1	0
3	H	10	0	0	2	0
4	A	12	0	12	1	0
5	G	7	0	10	1	0
6	A	5	0	0	0	0
6	B	5	0	0	0	0
6	C	3	0	0	0	0
6	D	14	0	0	0	0
6	E	9	0	0	0	0
6	F	11	0	0	0	0
6	G	9	0	0	0	0
6	H	9	0	0	1	0
All	All	14768	0	15078	168	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 168 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:1318:VAL:H	1:E:1329:ILE:HB	1.55	0.71
1:G:1274:ARG:HG3	1:G:1458:CYS:HB3	1.75	0.69
1:E:1380:ILE:HG12	1:E:1410:ILE:HB	1.78	0.66
2:D:233:VAL:HG12	2:D:234:VAL:HG23	1.78	0.65
2:D:255:PRO:HG2	2:D:258:LYS:HB2	1.78	0.65

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	197/215 (92%)	188 (95%)	9 (5%)	0	100	100
1	C	197/215 (92%)	184 (93%)	13 (7%)	0	100	100
1	E	197/215 (92%)	188 (95%)	9 (5%)	0	100	100
1	G	195/215 (91%)	185 (95%)	10 (5%)	0	100	100
2	B	262/291 (90%)	240 (92%)	20 (8%)	2 (1%)	19	49
2	D	258/291 (89%)	237 (92%)	19 (7%)	2 (1%)	19	49
2	F	262/291 (90%)	240 (92%)	21 (8%)	1 (0%)	34	66
2	H	257/291 (88%)	238 (93%)	18 (7%)	1 (0%)	34	66
All	All	1825/2024 (90%)	1700 (93%)	119 (6%)	6 (0%)	41	72

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	251	SER
2	D	234	VAL
2	F	234	VAL
2	H	234	VAL
2	B	234	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/193 (92%)	175 (99%)	2 (1%)	73	92

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	177/193 (92%)	176 (99%)	1 (1%)	86	96
1	E	177/193 (92%)	176 (99%)	1 (1%)	86	96
1	G	175/193 (91%)	175 (100%)	0	100	100
2	B	238/261 (91%)	237 (100%)	1 (0%)	91	97
2	D	236/261 (90%)	234 (99%)	2 (1%)	81	94
2	F	238/261 (91%)	236 (99%)	2 (1%)	81	94
2	H	235/261 (90%)	233 (99%)	2 (1%)	78	94
All	All	1653/1816 (91%)	1642 (99%)	11 (1%)	84	95

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	210	ASN
2	F	265	PRO
2	H	265	PRO
2	H	237	LYS
2	D	209	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	1396	ASN
2	F	87	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	SO4	F	1266	-	4,4,4	0.23	0	6,6,6	0.08	0
3	SO4	H	1266	-	4,4,4	0.24	0	6,6,6	0.07	0
3	SO4	E	2466	-	4,4,4	0.24	0	6,6,6	0.08	0
3	SO4	C	2467	-	4,4,4	0.23	0	6,6,6	0.07	0
3	SO4	A	2266	-	4,4,4	0.23	0	6,6,6	0.08	0
3	SO4	G	2467	-	4,4,4	0.23	0	6,6,6	0.10	0
3	SO4	H	1267	-	4,4,4	0.24	0	6,6,6	0.06	0
3	SO4	D	1266	-	4,4,4	0.23	0	6,6,6	0.08	0
3	SO4	G	2466	-	4,4,4	0.24	0	6,6,6	0.07	0
4	MES	A	2466	-	12,12,12	2.46	1 (8%)	15,16,16	2.22	5 (33%)
3	SO4	B	1266	-	4,4,4	0.24	0	6,6,6	0.06	0
5	PEG	G	2468	-	6,6,6	0.63	0	5,5,5	0.72	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MES	A	2466	-	-	0/6/14/14	0/1/1/1
5	PEG	G	2468	-	-	1/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	2466	MES	C8-S	-8.20	1.66	1.77

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2466	MES	C5-N4-C3	5.73	121.19	108.84

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	2466	MES	O3S-S-C8	3.55	112.95	106.00
4	A	2466	MES	C8-C7-N4	-2.29	103.68	112.36
4	A	2466	MES	C7-N4-C3	2.23	117.17	111.24
4	A	2466	MES	C6-C5-N4	-2.12	106.90	110.12

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	G	2468	PEG	C1-C2-O2-C3

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	H	1266	SO4	2	0
3	G	2466	SO4	1	0
4	A	2466	MES	1	0
5	G	2468	PEG	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	199/215 (92%)	-0.08	2 (1%) 82 77	71, 96, 125, 138	0
1	C	199/215 (92%)	-0.16	5 (2%) 57 47	48, 89, 142, 156	0
1	E	199/215 (92%)	-0.42	0 100 100	46, 76, 125, 181	0
1	G	197/215 (91%)	-0.17	8 (4%) 37 27	51, 85, 131, 213	0
2	B	264/291 (90%)	-0.01	4 (1%) 73 68	75, 100, 127, 155	0
2	D	262/291 (90%)	0.02	8 (3%) 49 39	79, 103, 144, 196	0
2	F	264/291 (90%)	-0.46	1 (0%) 92 91	33, 73, 110, 207	0
2	H	261/291 (89%)	-0.41	1 (0%) 92 91	49, 71, 101, 150	0
All	All	1845/2024 (91%)	-0.21	29 (1%) 72 66	33, 89, 130, 213	0

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	263	GLY	7.9
2	F	2	PRO	5.3
2	D	253	LYS	4.1
1	C	1461	ALA	4.0
1	G	1456	TYR	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	SO4	F	1266	5/5	0.92	0.17	144,147,147,148	0
3	SO4	D	1266	5/5	0.94	0.23	112,113,118,123	0
3	SO4	G	2467	5/5	0.94	0.18	90,92,97,104	0
3	SO4	A	2266	5/5	0.96	0.11	85,88,90,91	0
4	MES	A	2466	12/12	0.96	0.14	67,71,84,89	0
5	PEG	G	2468	7/7	0.96	0.09	33,46,58,60	0
3	SO4	H	1266	5/5	0.98	0.13	84,87,90,92	0
3	SO4	H	1267	5/5	0.98	0.15	59,62,80,84	0
3	SO4	G	2466	5/5	0.98	0.13	72,79,81,83	0
3	SO4	B	1266	5/5	0.98	0.14	66,76,80,84	0
3	SO4	E	2466	5/5	0.99	0.11	45,47,53,55	0
3	SO4	C	2467	5/5	0.99	0.09	37,57,61,73	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.