



# Full wwPDB X-ray Structure Validation Report ⓘ

May 14, 2024 – 06:11 pm BST

PDB ID : 8C48  
Title : Crystal structure of *Thermothelomyces thermophila* GH30 (double mutant EE) in complex with xylopentaose  
Authors : Dimarogona, M.; Pentari, C.; Kosinas, C.; Topakas, E.  
Deposited on : 2023-01-03  
Resolution : 1.40 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1714 (1.40-1.40)
Clashscore	141614	1812 (1.40-1.40)
Ramachandran outliers	138981	1763 (1.40-1.40)
Sidechain outliers	138945	1762 (1.40-1.40)
RSRZ outliers	127900	1674 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	482	 91% 6% 3%
1	B	482	 89% 7% 3%
2	C	9	 44% 44% 11%
3	D	2	 50% 50%
4	E	5	 80% 20%

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
5	F	3	 67% 33%
6	G	4	 75% 25%

## 2 Entry composition i

There are 9 unique types of molecules in this entry. The entry contains 7929 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GH30 family xylanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	453	3472	2164	606	687	15	0	11	0
1	B	449	3413	2131	594	673	15	0	6	0

There are 52 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	SER	-	expression tag	UNP G2Q1N4
A	2	MET	-	expression tag	UNP G2Q1N4
A	188	ALA	GLU	conflict	UNP G2Q1N4
A	278	ALA	GLU	conflict	UNP G2Q1N4
A	461	LEU	-	expression tag	UNP G2Q1N4
A	462	GLU	-	expression tag	UNP G2Q1N4
A	463	GLN	-	expression tag	UNP G2Q1N4
A	464	LYS	-	expression tag	UNP G2Q1N4
A	465	LEU	-	expression tag	UNP G2Q1N4
A	466	ILE	-	expression tag	UNP G2Q1N4
A	467	SER	-	expression tag	UNP G2Q1N4
A	468	GLU	-	expression tag	UNP G2Q1N4
A	469	GLU	-	expression tag	UNP G2Q1N4
A	470	ASP	-	expression tag	UNP G2Q1N4
A	471	LEU	-	expression tag	UNP G2Q1N4
A	472	ASN	-	expression tag	UNP G2Q1N4
A	473	SER	-	expression tag	UNP G2Q1N4
A	474	ALA	-	expression tag	UNP G2Q1N4
A	475	VAL	-	expression tag	UNP G2Q1N4
A	476	ASP	-	expression tag	UNP G2Q1N4
A	477	HIS	-	expression tag	UNP G2Q1N4
A	478	HIS	-	expression tag	UNP G2Q1N4
A	479	HIS	-	expression tag	UNP G2Q1N4
A	480	HIS	-	expression tag	UNP G2Q1N4
A	481	HIS	-	expression tag	UNP G2Q1N4

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	482	HIS	-	expression tag	UNP G2Q1N4
B	1	SER	-	expression tag	UNP G2Q1N4
B	2	MET	-	expression tag	UNP G2Q1N4
B	188	ALA	GLU	conflict	UNP G2Q1N4
B	278	ALA	GLU	conflict	UNP G2Q1N4
B	461	LEU	-	expression tag	UNP G2Q1N4
B	462	GLU	-	expression tag	UNP G2Q1N4
B	463	GLN	-	expression tag	UNP G2Q1N4
B	464	LYS	-	expression tag	UNP G2Q1N4
B	465	LEU	-	expression tag	UNP G2Q1N4
B	466	ILE	-	expression tag	UNP G2Q1N4
B	467	SER	-	expression tag	UNP G2Q1N4
B	468	GLU	-	expression tag	UNP G2Q1N4
B	469	GLU	-	expression tag	UNP G2Q1N4
B	470	ASP	-	expression tag	UNP G2Q1N4
B	471	LEU	-	expression tag	UNP G2Q1N4
B	472	ASN	-	expression tag	UNP G2Q1N4
B	473	SER	-	expression tag	UNP G2Q1N4
B	474	ALA	-	expression tag	UNP G2Q1N4
B	475	VAL	-	expression tag	UNP G2Q1N4
B	476	ASP	-	expression tag	UNP G2Q1N4
B	477	HIS	-	expression tag	UNP G2Q1N4
B	478	HIS	-	expression tag	UNP G2Q1N4
B	479	HIS	-	expression tag	UNP G2Q1N4
B	480	HIS	-	expression tag	UNP G2Q1N4
B	481	HIS	-	expression tag	UNP G2Q1N4
B	482	HIS	-	expression tag	UNP G2Q1N4

- Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	9	105	58	2	45	0	0	0


- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a

cetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace	
			Total	C	N				O
3	D	2	28	16	2	10	0	0	0

- Molecule 4 is an oligosaccharide called beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
4	E	5	46	25	21	0	0	0

- Molecule 5 is an oligosaccharide called beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
5	F	3	28	15	13	0	0	0

- Molecule 6 is an oligosaccharide called beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
			Total	C	O			
6	G	4	37	20	17	0	0	0

- Molecule 7 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	1	Total	C O	0	0
			4	2 2		
7	B	1	Total	C O	0	0
			4	2 2		

- Molecule 8 is FLUORIDE ION (three-letter code: F) (formula: F).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	1	Total	F	0	0
			1	1		

- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	473	Total	O	0	0
			473	473		
9	B	318	Total	O	0	0
			318	318		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: GH30 family xylanase

Chain A: 



- Molecule 1: GH30 family xylanase

Chain B: 



- Molecule 2: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 




- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D: 



- Molecule 4: beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose



Chain E:  80% 20%




- Molecule 5: beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose

Chain F:  67% 33%



- Molecule 6: beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose

Chain G:  75% 25%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	40.95Å 107.81Å 88.24Å 90.00° 95.47° 90.00°	Depositor
Resolution (Å)	45.98 – 1.40 40.76 – 1.40	Depositor EDS
% Data completeness (in resolution range)	98.9 (45.98-1.40) 98.9 (40.76-1.40)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.50 (at 1.40Å)	Xtrriage
Refinement program	REFMAC 5.8.0267	Depositor
R, $R_{free}$	0.128 , 0.156 0.142 , 0.137	Depositor DCC
$R_{free}$ test set	7591 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	17.5	Xtrriage
Anisotropy	0.367	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	7929	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, XYP, MAN, NAG, F, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.65	0/3563	0.73	0/4864
1	B	0.65	0/3507	0.74	0/4788
All	All	0.65	0/7070	0.73	0/9652

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3472	0	3286	13	0
1	B	3413	0	3234	12	0
2	C	105	0	88	1	0
3	D	28	0	25	0	0
4	E	46	0	0	0	0
5	F	28	0	0	0	0
6	G	37	0	0	0	0
7	B	8	0	12	3	0
8	B	1	0	0	0	0
9	A	473	0	0	6	2
9	B	318	0	0	3	0
All	All	7929	0	6645	26	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:127[A]:TRP:CZ3	9:A:818:HOH:O	2.15	0.99
1:A:127[A]:TRP:HZ3	9:A:818:HOH:O	1.48	0.95
1:B:337:ASP:HB3	1:B:342:THR:OG1	1.90	0.71
1:B:127[A]:TRP:CZ3	9:B:732:HOH:O	2.47	0.66
1:A:127[A]:TRP:HH2	9:A:649:HOH:O	1.78	0.65
1:A:22[B]:ARG:HG2	1:A:22[B]:ARG:HH11	1.61	0.64
1:A:22[B]:ARG:HH11	1:A:22[B]:ARG:CG	2.19	0.55
1:A:41:LEU:HD21	1:A:336:ILE:HG23	1.90	0.54
1:B:414:PRO:HB3	1:B:454:LEU:HB3	1.89	0.53
1:A:22[B]:ARG:HG2	1:A:22[B]:ARG:NH1	2.25	0.51
1:B:171:GLU:O	1:B:175[A]:GLU:HG2	2.12	0.49
1:B:226:THR:HA	7:B:501:EDO:H21	1.95	0.48
9:A:645:HOH:O	2:C:5:MAN:H62	2.14	0.47
1:B:127[A]:TRP:HZ3	9:B:732:HOH:O	1.89	0.47
7:B:501:EDO:C1	9:B:650:HOH:O	2.63	0.46
1:A:414:PRO:HB3	1:A:454:LEU:HB3	1.98	0.46
1:B:411:ASP:OD1	1:B:411:ASP:N	2.46	0.46
1:A:22[B]:ARG:NH2	9:A:509:HOH:O	2.50	0.45
1:B:318:TYR:HE2	7:B:502:EDO:H22	1.82	0.45
1:A:152:GLN:HG3	1:B:455:TYR:CD1	2.52	0.44
1:B:27:GLY:HA3	1:B:64:ILE:O	2.17	0.43
1:A:27:GLY:HA3	1:A:64:ILE:O	2.19	0.42
1:B:127[B]:TRP:N	1:B:127[B]:TRP:CD1	2.86	0.42
1:A:127[A]:TRP:CH2	9:A:818:HOH:O	2.54	0.42
1:B:213:TYR:HB3	1:B:214:PRO:HD3	2.03	0.41
1:A:229:CYS:HA	1:A:230:CYS:HA	1.88	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:A:503:HOH:O	9:A:531:HOH:O[1_455]	1.82	0.38
9:A:567:HOH:O	9:A:635:HOH:O[1_655]	1.97	0.23

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	462/482 (96%)	442 (96%)	20 (4%)	0	100	100
1	B	453/482 (94%)	436 (96%)	17 (4%)	0	100	100
All	All	915/964 (95%)	878 (96%)	37 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	363/379 (96%)	360 (99%)	3 (1%)	81	62
1	B	356/379 (94%)	352 (99%)	4 (1%)	73	50
All	All	719/758 (95%)	712 (99%)	7 (1%)	76	53

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	44	GLU
1	A	51	ASP
1	A	235	TRP
1	B	51	ASP
1	B	235	TRP
1	B	372	SER
1	B	432	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

23 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	C	1	1,2	14,14,15	0.63	0	17,19,21	0.72	0
2	NAG	C	2	2	14,14,15	0.77	0	17,19,21	1.07	1 (5%)
2	BMA	C	3	2	11,11,12	0.53	0	15,15,17	0.65	0
2	MAN	C	4	2	11,11,12	0.30	0	15,15,17	0.77	0
2	MAN	C	5	2	11,11,12	0.47	0	15,15,17	1.28	2 (13%)
2	MAN	C	6	2	11,11,12	0.46	0	15,15,17	0.68	1 (6%)
2	MAN	C	7	2	11,11,12	0.28	0	15,15,17	0.98	0
2	MAN	C	8	2	11,11,12	0.62	0	15,15,17	1.65	3 (20%)
2	MAN	C	9	2	11,11,12	0.37	0	15,15,17	1.26	1 (6%)
3	NAG	D	1	1,3	14,14,15	0.37	0	17,19,21	0.61	0
3	NAG	D	2	3	14,14,15	0.66	0	17,19,21	1.62	4 (23%)
4	XYP	E	1	4	10,10,10	0.20	0	14,14,14	0.46	0
4	XYP	E	2	4	9,9,10	0.27	0	10,12,14	0.93	1 (10%)
4	XYP	E	3	4	9,9,10	0.17	0	10,12,14	0.59	0
4	XYP	E	4	4	9,9,10	0.45	0	10,12,14	0.95	0
4	XYP	E	5	4	9,9,10	0.67	0	10,12,14	0.81	0
5	XYP	F	1	5	10,10,10	0.21	0	14,14,14	0.49	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	XYP	F	2	5	9,9,10	0.30	0	10,12,14	0.80	1 (10%)
5	XYP	F	3	5	9,9,10	0.31	0	10,12,14	0.49	0
6	XYP	G	1	6	10,10,10	0.38	0	14,14,14	0.60	0
6	XYP	G	2	6	9,9,10	0.46	0	10,12,14	1.26	1 (10%)
6	XYP	G	3	6	9,9,10	0.30	0	10,12,14	1.01	0
6	XYP	G	4	6	9,9,10	0.16	0	10,12,14	0.54	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	0/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	MAN	C	4	2	-	0/2/19/22	0/1/1/1
2	MAN	C	5	2	-	2/2/19/22	0/1/1/1
2	MAN	C	6	2	-	0/2/19/22	0/1/1/1
2	MAN	C	7	2	-	0/2/19/22	0/1/1/1
2	MAN	C	8	2	-	0/2/19/22	0/1/1/1
2	MAN	C	9	2	-	2/2/19/22	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
4	XYP	E	1	4	-	-	0/1/1/1
4	XYP	E	2	4	-	-	0/1/1/1
4	XYP	E	3	4	-	-	0/1/1/1
4	XYP	E	4	4	-	-	0/1/1/1
4	XYP	E	5	4	-	-	0/1/1/1
5	XYP	F	1	5	-	-	0/1/1/1
5	XYP	F	2	5	-	-	0/1/1/1
5	XYP	F	3	5	-	-	0/1/1/1
6	XYP	G	1	6	-	-	0/1/1/1
6	XYP	G	2	6	-	-	0/1/1/1
6	XYP	G	3	6	-	-	0/1/1/1
6	XYP	G	4	6	-	-	0/1/1/1

There are no bond length outliers.

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	8	MAN	O5-C5-C6	4.28	113.92	107.20
6	G	2	XYP	C1-C2-C3	3.48	113.94	109.67
3	D	2	NAG	C1-C2-N2	-3.47	104.56	110.49
2	C	9	MAN	C1-C2-C3	3.24	113.65	109.67
3	D	2	NAG	C2-N2-C7	-3.10	118.48	122.90
4	E	2	XYP	C1-C2-C3	2.78	113.09	109.67
3	D	2	NAG	C1-O5-C5	2.57	115.68	112.19
2	C	5	MAN	C1-C2-C3	2.54	112.78	109.67
2	C	8	MAN	O2-C2-C1	2.29	113.84	109.15
2	C	5	MAN	C6-C5-C4	-2.24	107.76	113.00
5	F	2	XYP	C1-C2-C3	2.23	112.41	109.67
3	D	2	NAG	O3-C3-C2	-2.21	104.89	109.47
2	C	2	NAG	C1-C2-N2	2.08	114.03	110.49
2	C	6	MAN	C1-O5-C5	2.02	114.93	112.19
2	C	8	MAN	C1-O5-C5	2.01	114.92	112.19

There are no chirality outliers.

All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	9	MAN	O5-C5-C6-O6
2	C	5	MAN	O5-C5-C6-O6
2	C	5	MAN	C4-C5-C6-O6
2	C	9	MAN	C4-C5-C6-O6
3	D	2	NAG	C4-C5-C6-O6
3	D	2	NAG	O5-C5-C6-O6


There are no ring outliers.


1 monomer is involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	5	MAN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
7	EDO	B	502	-	3,3,3	0.19	0	2,2,2	0.59	0
7	EDO	B	501	-	3,3,3	0.53	0	2,2,2	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	EDO	B	502	-	-	0/1/1/1	-
7	EDO	B	501	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	B	501	EDO	O1-C1-C2-O2

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
7	B	502	EDO	1	0
7	B	501	EDO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	453/482 (93%)	-0.62	1 (0%) 95 93	13, 18, 34, 52	0
1	B	449/482 (93%)	-0.28	15 (3%) 46 46	17, 27, 54, 76	0
All	All	902/964 (93%)	-0.45	16 (1%) 68 68	13, 21, 47, 76	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	410	ALA	4.4
1	B	407	SER	4.2
1	B	406	SER	3.6
1	B	372	SER	3.6
1	B	437	ASP	3.4
1	B	296	ALA	3.3
1	B	399	ALA	2.8
1	B	337	ASP	2.7
1	B	286	TRP	2.6
1	B	339	ASN	2.6
1	B	408	SER	2.2
1	B	409	SER	2.1
1	B	395	GLY	2.1
1	B	328	ASN	2.1
1	A	9	GLY	2.1
1	B	411	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	D	2	14/15	0.83	0.12	29,34,47,50	0
4	XYP	E	1	10/10	0.83	0.18	40,51,65,67	0
2	MAN	C	9	11/12	0.84	0.18	55,63,71,73	0
5	XYP	F	1	10/10	0.86	0.11	47,55,60,62	0
6	XYP	G	1	10/10	0.87	0.14	38,44,55,65	0
2	MAN	C	6	11/12	0.88	0.20	33,43,52,57	0
5	XYP	F	2	9/10	0.92	0.12	41,42,44,45	0
2	MAN	C	5	11/12	0.92	0.18	40,50,59,64	0
6	XYP	G	2	9/10	0.92	0.10	25,28,32,34	0
6	XYP	G	4	9/10	0.92	0.13	28,30,32,35	0
4	XYP	E	4	9/10	0.93	0.07	20,22,24,25	0
2	BMA	C	3	11/12	0.93	0.10	26,29,32,38	0
4	XYP	E	2	9/10	0.93	0.07	29,32,37,38	0
2	MAN	C	8	11/12	0.94	0.13	34,42,51,53	0
6	XYP	G	3	9/10	0.94	0.11	20,24,28,28	0
4	XYP	E	3	9/10	0.94	0.06	22,25,27,32	0
2	MAN	C	7	11/12	0.95	0.10	26,28,37,39	0
2	MAN	C	4	11/12	0.96	0.07	26,30,35,35	0
5	XYP	F	3	9/10	0.96	0.07	35,39,41,44	0
3	NAG	D	1	14/15	0.96	0.06	20,22,33,34	0
4	XYP	E	5	9/10	0.97	0.05	20,21,23,26	0
2	NAG	C	2	14/15	0.97	0.05	20,23,29,30	0
2	NAG	C	1	14/15	0.98	0.05	15,18,26,32	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around Chain C:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around Chain D:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)









**Electron density around Chain E:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around Chain F:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
7	EDO	B	502	4/4	0.95	0.23	31,31,34,37	0
7	EDO	B	501	4/4	0.98	0.13	25,27,37,40	0
8	F	B	503	1/1	0.99	0.07	31,31,31,31	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.