



Full wwPDB X-ray Structure Validation Report

Sep 26, 2023 – 02:15 PM EDT


PDB ID : 6C9Z
Title : THE CRYSTAL STRUCTURE OF THE alpha-Glucosidase (GH 31) W169Y mutant FROM RUMINOCOCCUS OBEUM ATCC 29174 in complex with voglibose
Authors : Tan, K.; Tesar, C.; Jedrzejczak, R.; Joachimiak, A.; Midwest Center for Macromolecular Research (MCMR)
Deposited on : 2018-01-29
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	666	 2% 92% 7%
1	B	666	 2% 90% 9%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11011 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glycosyl hydrolase, family 31.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	664	5350	3443	874	998	35	0	0	0
1	B	663	5362	3445	874	1008	35	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP A5ZY13
A	-1	ASN	-	expression tag	UNP A5ZY13
A	0	ALA	-	expression tag	UNP A5ZY13
A	169	TYR	TRP	engineered mutation	UNP A5ZY13
B	-2	SER	-	expression tag	UNP A5ZY13
B	-1	ASN	-	expression tag	UNP A5ZY13
B	0	ALA	-	expression tag	UNP A5ZY13
B	169	TYR	TRP	engineered mutation	UNP A5ZY13

- Molecule 2 is (1S,2S,3R,4S,5S)-5-[(1,3-dihydroxypropan-2-yl)amino]-1-(hydroxymethyl)cyclohexane-1,2,3,4-tetrol (three-letter code: VOG) (formula: C₁₀H₂₁NO₇).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	18	10	1	7	0	0
2	B	1	18	10	1	7	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	156	156	156	0	0
3	B	107	107	107	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glycosyl hydrolase, family 31



- Molecule 1: Glycosyl hydrolase, family 31



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	67.75Å 126.35Å 88.19Å 90.00° 107.82° 90.00°	Depositor
Resolution (Å)	32.25 – 2.10 32.25 – 2.10	Depositor EDS
% Data completeness (in resolution range)	96.1 (32.25-2.10) 96.3 (32.25-2.10)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.29 (at 2.10Å)	Xtrriage
Refinement program	PHENIX 1.8.1_1168	Depositor
R, R_{free}	0.192 , 0.246 0.193 , 0.246	Depositor DCC
R_{free} test set	3938 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	38.3	Xtrriage
Anisotropy	0.782	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 33.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	11011	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: VOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.37	0/5492	0.52	0/7419
1	B	0.34	0/5504	0.51	0/7436
All	All	0.36	0/10996	0.52	0/14855

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5350	0	5049	23	0
1	B	5362	0	5046	29	0
2	A	18	0	0	0	0
2	B	18	0	0	0	0
3	A	156	0	0	1	0
3	B	107	0	0	2	0
All	All	11011	0	10095	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:324:LYS:NZ	1:B:333:ASP:O	2.23	0.71
1:B:204:PHE:HB2	1:B:272:PRO:HB3	1.82	0.61
1:B:573:MET:HG2	1:B:608:ARG:HA	1.84	0.58
1:A:333:ASP:O	1:B:324:LYS:NZ	2.37	0.57
1:B:59:GLY:HA2	1:B:433:LYS:HD3	1.87	0.56
1:B:38:PHE:CZ	1:B:132:ILE:HD11	2.42	0.54
1:A:45:ASP:O	1:A:63:ARG:NH1	2.40	0.54
1:B:344:GLN:HG3	3:B:904:HOH:O	2.07	0.54
1:A:436:PRO:O	1:A:440:MET:HG3	2.08	0.52
1:A:444:MET:HE1	1:A:519:ALA:HB2	1.93	0.51
1:B:270:VAL:HG12	1:B:271:TRP:H	1.75	0.51
1:A:270:VAL:HG12	1:A:271:TRP:H	1.76	0.51
1:B:380:TYR:HA	1:B:410:MET:HG3	1.93	0.50
1:B:527:LYS:HB2	1:B:547:MET:HB2	1.95	0.49
1:A:502:ASN:O	1:A:506:ARG:HG3	2.13	0.49
1:B:259:LYS:HD3	1:B:263:GLY:O	2.13	0.49
1:A:59:GLY:HA2	1:A:433:LYS:HD3	1.95	0.48
1:B:85:TYR:CE2	1:B:419:GLY:HA3	2.49	0.47
1:A:114:GLY:HA3	3:A:909:HOH:O	2.15	0.47
1:A:644:GLU:OE2	1:A:659:ARG:NH2	2.43	0.47
1:B:459:ARG:NH2	1:B:493:ASN:HB2	2.29	0.46
1:B:504:ARG:O	1:B:508:VAL:HG23	2.15	0.46
1:A:435:LEU:HD13	1:A:526:PHE:CE2	2.51	0.46
1:A:310:GLU:N	1:A:311:PRO:HA	2.31	0.45
1:B:397:LYS:HD3	1:B:399:PHE:CZ	2.52	0.45
1:B:157:PRO:HG3	1:B:445:TYR:CD1	2.52	0.45
1:B:659:ARG:HE	1:B:661:LEU:HD21	1.81	0.44
1:A:114:GLY:HA2	1:A:117:ARG:O	2.16	0.44
1:B:90:PHE:CE2	1:B:92:ILE:HD11	2.53	0.44
1:A:344:GLN:HG3	3:B:904:HOH:O	2.16	0.44
1:B:502:ASN:O	1:B:506:ARG:HG3	2.18	0.43
1:B:459:ARG:HH21	1:B:493:ASN:HB2	1.83	0.43
1:B:105:TYR:CE2	1:B:107:SER:HB3	2.54	0.43
1:A:38:PHE:CZ	1:A:132:ILE:HD11	2.54	0.43
1:A:574:LYS:HG3	1:A:635:TYR:CE2	2.55	0.42
1:B:164:PHE:HD1	1:B:475:MET:HG2	1.85	0.42
1:B:93:VAL:O	1:B:98:THR:HA	2.20	0.42
1:A:341:TRP:CD1	1:A:341:TRP:C	2.93	0.41
1:A:197:ASP:O	1:A:199:ASP:N	2.53	0.41
1:A:190:PRO:HB2	1:A:505:TYR:CE1	2.55	0.41
1:B:291:ASP:O	1:B:294:ARG:HG3	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:569:LEU:HA	1:B:570:PRO:HD3	1.91	0.41
1:B:210:ASN:OD1	1:B:212:LYS:HB3	2.21	0.41
1:B:282:ASN:HA	1:B:283:PRO:HD3	1.86	0.41
1:A:153:ARG:HG2	1:A:154:SER:N	2.36	0.41
1:B:346:LYS:HA	1:B:346:LYS:HD2	1.90	0.41
1:B:422:LYS:HD2	1:B:455:ASP:OD2	2.20	0.41
1:A:107:SER:OG	1:A:108:LYS:N	2.52	0.41
1:A:422:LYS:HD2	1:A:455:ASP:OD2	2.20	0.41
1:A:277:PHE:CZ	1:A:311:PRO:HB2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	662/666 (99%)	626 (95%)	35 (5%)	1 (0%)	47	49
1	B	661/666 (99%)	623 (94%)	34 (5%)	4 (1%)	25	21
All	All	1323/1332 (99%)	1249 (94%)	69 (5%)	5 (0%)	34	32

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	58	ARG
1	B	70	ASN
1	B	169	TYR
1	B	270	VAL
1	A	198	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	546/575 (95%)	534 (98%)	12 (2%)	52	57
1	B	550/575 (96%)	541 (98%)	9 (2%)	62	69
All	All	1096/1150 (95%)	1075 (98%)	21 (2%)	57	63

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	GLU
1	A	25	GLU
1	A	34	GLN
1	A	63	ARG
1	A	169	TYR
1	A	294	ARG
1	A	341	TRP
1	A	380	TYR
1	A	417	TRP
1	A	465	PHE
1	A	611	LYS
1	A	662	THR
1	B	12	ASP
1	B	192	ASP
1	B	294	ARG
1	B	305	TRP
1	B	380	TYR
1	B	417	TRP
1	B	465	PHE
1	B	537	LYS
1	B	569	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	VOG	A	701	-	18,18,18	1.74	4 (22%)	21,26,26	1.02	0
2	VOG	B	701	-	18,18,18	1.71	4 (22%)	21,26,26	1.05	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	VOG	A	701	-	-	0/11/34/34	0/1/1/1
2	VOG	B	701	-	-	5/11/34/34	0/1/1/1

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	VOG	C1-C2	-3.45	1.50	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	701	VOG	C8-N	3.07	1.53	1.48
2	B	701	VOG	O1-C1	-3.06	1.39	1.44
2	A	701	VOG	O1-C1	-2.92	1.39	1.44
2	A	701	VOG	C8-N	2.85	1.52	1.48
2	B	701	VOG	C1-C2	-2.79	1.50	1.53
2	B	701	VOG	C6-C1	-2.61	1.50	1.53
2	A	701	VOG	C6-C1	-2.23	1.50	1.53

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	701	VOG	C10-C8-N	2.07	116.27	109.97

There are no chirality outliers.


All (5) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	701	VOG	C6-C1-C7-O7
2	B	701	VOG	C10-C8-C9-O9
2	B	701	VOG	O10-C10-C8-C9
2	B	701	VOG	O10-C10-C8-N
2	B	701	VOG	O1-C1-C7-O7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	664/666 (99%)	-0.07	13 (1%) 65 69	31, 45, 65, 103	2 (0%)
1	B	663/666 (99%)	0.02	16 (2%) 59 64	35, 49, 72, 97	1 (0%)
All	All	1327/1332 (99%)	-0.02	29 (2%) 62 66	31, 47, 70, 103	3 (0%)

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	470	VAL	4.5
1	B	405	SER	3.8
1	A	438	LEU	2.8
1	B	402	PHE	2.7
1	B	474	LEU	2.7
1	B	341	TRP	2.6
1	B	235	ILE	2.6
1	B	471	PHE	2.6
1	A	469	GLY	2.6
1	B	379	GLY	2.6
1	A	432	LEU	2.5
1	A	408	ILE	2.5
1	B	446	THR	2.4
1	B	86	GLY	2.4
1	A	470	VAL	2.4
1	B	378	PHE	2.4
1	A	29	TYR	2.3
1	B	309	ASN	2.3
1	A	466	LEU	2.2
1	A	26	ALA	2.2
1	B	660	VAL	2.2
1	A	468	LEU	2.2
1	A	34	GLN	2.2
1	B	406	SER	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	403	SER	2.1
1	A	24	GLU	2.0
1	B	511	LEU	2.0
1	A	446	THR	2.0
1	A	194	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	VOG	B	701	18/18	0.93	0.11	38,42,49,49	0
2	VOG	A	701	18/18	0.95	0.10	32,38,43,46	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around VOG B 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.