



## wwPDB EM Validation Summary Report ⓘ

Dec 14, 2024 – 01:20 PM EST

PDB ID : 9CAQ  
EMDB ID : EMD-45400  
Title : Cryo-EM structure of a human MCM2-7 double hexamer formed from independently loaded MCM2-7 single hexamers  
Authors : Yang, R.; Hunker, O.; Bleichert, F.  
Deposited on : 2024-06-17  
Resolution : 3.20 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev113  
Mogul : 2022.3.0, CSD as543be (2022)  
MolProbity : 4.02b-467  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.40

# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:  
*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	210492	15764
Ramachandran outliers	207382	16835
Sidechain outliers	206894	16415

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	2	904	
1	A	904	
2	3	808	
2	B	808	
3	4	863	
3	C	863	
4	5	737	
4	D	737	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
5	6	821	 <p>67% 10% 23%</p>
5	E	821	 <p>64% 13% 23%</p>
6	7	722	 <p>71% 10% 19%</p>
6	F	722	 <p>63% 15% 21%</p>
7	O	44	 <p>93% 7%</p>
8	S	44	 <p>91% 9%</p>

## 2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 120223 atoms, of which 59848 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA replication licensing factor MCM2.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
1	2	619	9773	3071	4892	871	909	30	0	0
1	A	606	9564	3005	4784	853	892	30	0	0

- Molecule 2 is a protein called DNA replication licensing factor MCM3.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
2	3	619	9762	3045	4893	856	943	25	0	0
2	B	596	9398	2927	4720	824	903	24	0	0

- Molecule 3 is a protein called DNA replication licensing factor MCM4.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
3	4	637	10194	3195	5117	905	950	27	0	0
3	C	635	10179	3190	5113	903	946	27	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
4	650	MET	LEU	variant	UNP P33991
C	650	MET	LEU	variant	UNP P33991

- Molecule 4 is a protein called DNA replication licensing factor MCM5.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
4	5	617	9720	3033	4887	860	905	35	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
4	D	614	9669	3018	4858	854	904	35	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
5	-2	SER	-	expression tag	UNP P33992
5	-1	ASN	-	expression tag	UNP P33992
5	0	ALA	-	expression tag	UNP P33992
D	-2	SER	-	expression tag	UNP P33992
D	-1	ASN	-	expression tag	UNP P33992
D	0	ALA	-	expression tag	UNP P33992

- Molecule 5 is a protein called DNA replication licensing factor MCM6.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
5	6	635	10144	3178	5084	900	956	26	0	0
5	E	632	10099	3162	5067	895	949	26	0	0

- Molecule 6 is a protein called DNA replication licensing factor MCM7.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
6	7	587	9377	2928	4714	829	874	32	0	0
6	F	571	9109	2838	4585	805	849	32	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
7	-2	SER	-	expression tag	UNP P33993
7	-1	ASN	-	expression tag	UNP P33993
7	0	ALA	-	expression tag	UNP P33993
F	-2	SER	-	expression tag	UNP P33993
F	-1	ASN	-	expression tag	UNP P33993
F	0	ALA	-	expression tag	UNP P33993

- Molecule 7 is a DNA chain called DNA (44-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace	
7	O	44	Total	C	H	N	O	P	0	0
			1409	440	506	157	262	44		

- Molecule 8 is a DNA chain called DNA (44-MER).

Mol	Chain	Residues	Atoms					AltConf	Trace	
8	S	44	Total	C	H	N	O	P	0	0
			1409	440	508	151	266	44		

- Molecule 9 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
9	2	1	Total	Zn	0
			1	1	
9	4	1	Total	Zn	0
			1	1	
9	5	1	Total	Zn	0
			1	1	
9	6	1	Total	Zn	0
			1	1	
9	7	1	Total	Zn	0
			1	1	
9	A	1	Total	Zn	0
			1	1	
9	C	1	Total	Zn	0
			1	1	
9	D	1	Total	Zn	0
			1	1	
9	E	1	Total	Zn	0
			1	1	
9	F	1	Total	Zn	0
			1	1	

- Molecule 10 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		AltConf
10	2	1	Total	Mg	0
			1	1	
10	5	1	Total	Mg	0
			1	1	
10	6	1	Total	Mg	0
			1	1	

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
10	7	1	Total	Mg	0
			1	1	
10	A	1	Total	Mg	0
			1	1	
10	D	2	Total	Mg	0
			2	2	
10	E	1	Total	Mg	0
			1	1	
10	F	1	Total	Mg	0
			1	1	

- Molecule 11 is ADENOSINE-5'-TRIPHOSPHATE (three-letter code: ATP) (formula:  $C_{10}H_{16}N_5O_{13}P_3$ ).



Mol	Chain	Residues	Atoms					AltConf	
			Total	C	H	N	O		P
11	2	1	Total	C	H	N	O	P	0
			43	10	12	5	13	3	
11	A	1	Total	C	H	N	O	P	0
			43	10	12	5	13	3	


- Molecule 12 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula:  $C_{10}H_{15}N_5O_{10}P_2$ ).




Mol	Chain	Residues	Atoms					AltConf	
			Total	C	H	N	O		P
12	3	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	
12	4	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	
12	5	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	
12	7	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	
12	B	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	
12	D	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	
12	E	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	
12	F	1	Total	C	H	N	O	P	0
			39	10	12	5	10	2	









• Molecule 2: DNA replication licensing factor MCM3







• Molecule 3: DNA replication licensing factor MCM4




• Molecule 4: DNA replication licensing factor MCM5












• Molecule 7: DNA (44-MER)



• Molecule 8: DNA (44-MER)





## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	46418	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS GLACIOS	Depositor
Voltage (kV)	200	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	51.54	Depositor
Minimum defocus (nm)	1000	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	2.533	Depositor
Minimum map value	-0.019	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.031	Depositor
Recommended contour level	0.09	Depositor
Map size ( $\text{\AA}$ )	434.0, 434.0, 434.0	wwPDB
Map dimensions	500, 500, 500	wwPDB
Map angles ( $^\circ$ )	90.0, 90.0, 90.0	wwPDB
Pixel spacing ( $\text{\AA}$ )	0.868, 0.868, 0.868	Depositor

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: ADP, ZN, MG, ATP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	2	0.36	0/4974	0.66	0/6720
1	A	0.33	0/4870	0.63	0/6580
2	3	0.38	0/4943	0.67	0/6674
2	B	0.35	0/4749	0.66	0/6414
3	4	0.37	0/5167	0.66	0/6983
3	C	0.36	0/5156	0.65	0/6967
4	5	0.39	0/4910	0.71	1/6602 (0.0%)
4	D	0.36	0/4888	0.67	0/6573
5	6	0.36	0/5143	0.66	0/6939
5	E	0.34	0/5113	0.67	0/6898
6	7	0.38	0/4735	0.66	0/6387
6	F	0.37	0/4588	0.70	1/6185 (0.0%)
7	O	0.86	1/1013 (0.1%)	1.10	0/1561
8	S	0.88	0/1009	1.14	0/1555
All	All	0.39	1/61258 (0.0%)	0.69	2/83038 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	O	38	DT	C1 <sup>1</sup> -N1	5.10	1.55	1.49

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( <sup>o</sup> )	Ideal( <sup>o</sup> )
6	F	357	LEU	CA-CB-CG	5.58	128.14	115.30
4	5	276	LEU	CA-CB-CG	5.18	127.20	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	2	4881	4892	4892	56	0
1	A	4780	4784	4784	68	0
2	3	4869	4893	4893	89	0
2	B	4678	4720	4720	93	0
3	4	5077	5117	5117	66	0
3	C	5066	5113	5112	78	0
4	5	4833	4887	4887	89	0
4	D	4811	4858	4858	78	0
5	6	5060	5084	5084	57	0
5	E	5032	5067	5067	69	0
6	7	4663	4714	4714	51	0
6	F	4524	4585	4585	96	0
7	O	903	506	506	3	0
8	S	901	508	508	4	0
9	2	1	0	0	0	0
9	4	1	0	0	0	0
9	5	1	0	0	0	0
9	6	1	0	0	0	0
9	7	1	0	0	0	0
9	A	1	0	0	0	0
9	C	1	0	0	0	0
9	D	1	0	0	0	0
9	E	1	0	0	0	0
9	F	1	0	0	0	0
10	2	1	0	0	0	0
10	5	1	0	0	0	0
10	6	1	0	0	0	0
10	7	1	0	0	0	0
10	A	1	0	0	0	0
10	D	2	0	0	0	0
10	E	1	0	0	0	0
10	F	1	0	0	0	0
11	2	31	12	12	2	0
11	A	31	12	12	2	0
12	3	27	12	12	2	0
12	4	27	12	12	0	0
12	5	27	12	12	5	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
12	7	27	12	12	3	0
12	B	27	12	12	2	0
12	D	27	12	12	5	0
12	E	27	12	12	0	0
12	F	27	12	12	8	0
All	All	60375	59848	59847	777	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 777 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:F:343:GLU:O	12:F:1002:ADP:N1	1.58	1.35
6:F:388:SER:OG	12:F:1002:ADP:O3B	1.77	0.99
6:F:343:GLU:O	12:F:1002:ADP:C2	2.33	0.81
6:F:357:LEU:HD21	6:F:485:LEU:HD11	1.64	0.79
6:F:182:TYR:OH	6:F:193:GLN:OE1	2.03	0.77

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	2	613/904 (68%)	557 (91%)	56 (9%)	0	100	100
1	A	600/904 (66%)	556 (93%)	44 (7%)	0	100	100
2	3	613/808 (76%)	559 (91%)	54 (9%)	0	100	100
2	B	592/808 (73%)	548 (93%)	44 (7%)	0	100	100
3	4	631/863 (73%)	584 (93%)	47 (7%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	629/863 (73%)	577 (92%)	52 (8%)	0	100	100
4	5	607/737 (82%)	559 (92%)	48 (8%)	0	100	100
4	D	604/737 (82%)	559 (92%)	45 (8%)	0	100	100
5	6	629/821 (77%)	591 (94%)	38 (6%)	0	100	100
5	E	626/821 (76%)	590 (94%)	36 (6%)	0	100	100
6	7	575/722 (80%)	543 (94%)	32 (6%)	0	100	100
6	F	555/722 (77%)	498 (90%)	57 (10%)	0	100	100
All	All	7274/9710 (75%)	6721 (92%)	553 (8%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	2	540/781 (69%)	535 (99%)	5 (1%)	75	89
1	A	528/781 (68%)	523 (99%)	5 (1%)	75	89
2	3	536/707 (76%)	526 (98%)	10 (2%)	52	76
2	B	516/707 (73%)	510 (99%)	6 (1%)	67	85
3	4	562/753 (75%)	555 (99%)	7 (1%)	67	85
3	C	561/753 (74%)	556 (99%)	5 (1%)	75	89
4	5	525/627 (84%)	516 (98%)	9 (2%)	56	78
4	D	524/627 (84%)	518 (99%)	6 (1%)	70	86
5	6	563/724 (78%)	559 (99%)	4 (1%)	81	92
5	E	560/724 (77%)	550 (98%)	10 (2%)	54	77
6	7	510/621 (82%)	504 (99%)	6 (1%)	67	85
6	F	495/621 (80%)	487 (98%)	8 (2%)	58	79
All	All	6420/8426 (76%)	6339 (99%)	81 (1%)	64	83

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	32	GLN
5	E	551	ARG
4	D	361	SER
5	E	224	LEU
6	F	164	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
3	4	430	HIS
3	C	521	GLN
6	F	464	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 29 ligands modelled in this entry, 19 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
12	ADP	4	902	10	24,29,29	0.84	0	29,45,45	1.28	2 (6%)
12	ADP	D	1004	10	24,29,29	0.82	0	29,45,45	1.29	2 (6%)
12	ADP	F	1002	10	24,29,29	1.00	2 (8%)	29,45,45	1.45	3 (10%)
11	ATP	A	1002	10	28,33,33	1.43	1 (3%)	34,52,52	1.44	5 (14%)
12	ADP	3	901	-	24,29,29	0.86	0	29,45,45	1.31	3 (10%)
11	ATP	2	1002	10	28,33,33	1.01	1 (3%)	34,52,52	0.99	3 (8%)
12	ADP	7	1002	10	24,29,29	0.85	0	29,45,45	1.26	2 (6%)
12	ADP	B	901	10	24,29,29	0.89	0	29,45,45	1.35	3 (10%)
12	ADP	E	1002	10	24,29,29	0.85	0	29,45,45	1.31	2 (6%)
12	ADP	5	1002	10	24,29,29	0.82	0	29,45,45	1.27	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
12	ADP	4	902	10	-	0/12/32/32	0/3/3/3
12	ADP	D	1004	10	-	0/12/32/32	0/3/3/3
12	ADP	F	1002	10	-	4/12/32/32	0/3/3/3
11	ATP	A	1002	10	-	5/18/38/38	0/3/3/3
12	ADP	3	901	-	-	3/12/32/32	0/3/3/3
11	ATP	2	1002	10	-	5/18/38/38	0/3/3/3
12	ADP	7	1002	10	-	2/12/32/32	0/3/3/3
12	ADP	B	901	10	-	5/12/32/32	0/3/3/3
12	ADP	E	1002	10	-	3/12/32/32	0/3/3/3
12	ADP	5	1002	10	-	2/12/32/32	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	A	1002	ATP	PA-O3A	-6.53	1.52	1.59
11	2	1002	ATP	PB-O3B	-3.29	1.55	1.59
12	F	1002	ADP	O4'-C1'	2.66	1.44	1.40
12	F	1002	ADP	C2-N3	2.26	1.35	1.32

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	A	1002	ATP	C4'-O4'-C1'	-5.24	105.13	109.92
12	4	902	ADP	N3-C2-N1	-4.24	122.91	128.67
12	D	1004	ADP	N3-C2-N1	-4.22	122.94	128.67
12	3	901	ADP	N3-C2-N1	-4.15	123.05	128.67
12	E	1002	ADP	N3-C2-N1	-4.12	123.08	128.67

There are no chirality outliers.

5 of 29 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	2	1002	ATP	C5'-O5'-PA-O1A
11	A	1002	ATP	C5'-O5'-PA-O1A
11	A	1002	ATP	C5'-O5'-PA-O2A
11	A	1002	ATP	C5'-O5'-PA-O3A
12	5	1002	ADP	C5'-O5'-PA-O2A



There are no ring outliers.



8 monomers are involved in 29 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
12	D	1004	ADP	5	0
12	F	1002	ADP	8	0
11	A	1002	ATP	2	0
12	3	901	ADP	2	0
11	2	1002	ATP	2	0
12	7	1002	ADP	3	0
12	B	901	ADP	2	0
12	5	1002	ADP	5	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-45400. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)

### 6.2.1 Primary map



X Index: 250




Y Index: 250




Z Index: 250


### 6.2.2 Raw map



X Index: 250



Y Index: 250




Z Index: 250

The images above show central slices of the map in three orthogonal directions.



## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 224




Y Index: 253




Z Index: 303


### 6.3.2 Raw map



X Index: 254



Y Index: 0




Z Index: 254


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.09. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 400 nm<sup>3</sup>; this corresponds to an approximate mass of 361 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)




\*Reported resolution corresponds to spatial frequency of 0.312 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.312 Å<sup>-1</sup>

## 8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.20	-	-
Author-provided FSC curve	3.13	3.84	3.25
Unmasked-calculated*	4.28	9.17	6.49


\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 4.28 differs from the reported value 3.2 by more than 10 %



## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-45400 and PDB model 9CAQ. Per-residue inclusion information can be found in section 3 on page 9.

### 9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.09 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.09).































## 9.4 Atom inclusion [i](#)



At the recommended contour level, 79% of all backbone atoms, 79% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.09) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7940	 0.3960
2	 0.7900	 0.3800
3	 0.8050	 0.3830
4	 0.8340	 0.4400
5	 0.7890	 0.3600
6	 0.8130	 0.4180
7	 0.8270	 0.4350
A	 0.7890	 0.3960
B	 0.7700	 0.3750
C	 0.7960	 0.3980
D	 0.8040	 0.4150
E	 0.7710	 0.3900
F	 0.7760	 0.3570
O	 0.8360	 0.4080
S	 0.8470	 0.4070

