



# wwPDB X-ray Structure Validation Summary Report ⓘ

May 16, 2020 – 05:23 pm BST

PDB ID : 2COV  
Title : Crystal structure of CBM31 from beta-1,3-xylanase  
Authors : Hashimoto, H.; Tamai, Y.; Okazaki, F.; Tamaru, Y.; Shimizu, T.; Araki, T.;  
Sato, M.  
Deposited on : 2005-05-18  
Resolution : 1.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1023 (1.28-1.24)
Clashscore	141614	1060 (1.28-1.24)
Ramachandran outliers	138981	1029 (1.28-1.24)
Sidechain outliers	138945	1028 (1.28-1.24)
RSRZ outliers	127900	1004 (1.28-1.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	D	104	 3% 77% 9% • 12%
1	E	104	 4% 79% 8% • 12%
1	F	104	 2% 77% 7% 16%
1	G	104	 4% 74% 8% • 15%
1	H	104	 5% 74% 8% • 16%
1	I	104	 5% 75% 10% 15%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 4937 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called beta-1,3-xylanase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	D	92	751	473	127	147	4	0	0	0
1	E	92	751	473	127	147	4	1	0	0
1	F	87	712	450	120	138	4	13	0	0
1	G	88	720	454	122	140	4	14	0	0
1	H	87	712	450	120	138	4	13	0	0
1	I	88	720	454	122	140	4	15	0	0

There are 54 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	374	MET	-	INITIATING METHIONINE	UNP Q8RS40
D	470	LEU	-	EXPRESSION TAG	UNP Q8RS40
D	471	GLU	-	EXPRESSION TAG	UNP Q8RS40
D	472	HIS	-	EXPRESSION TAG	UNP Q8RS40
D	473	HIS	-	EXPRESSION TAG	UNP Q8RS40
D	474	HIS	-	EXPRESSION TAG	UNP Q8RS40
D	475	HIS	-	EXPRESSION TAG	UNP Q8RS40
D	476	HIS	-	EXPRESSION TAG	UNP Q8RS40
D	477	HIS	-	EXPRESSION TAG	UNP Q8RS40
E	374	MET	-	INITIATING METHIONINE	UNP Q8RS40
E	470	LEU	-	EXPRESSION TAG	UNP Q8RS40
E	471	GLU	-	EXPRESSION TAG	UNP Q8RS40
E	472	HIS	-	EXPRESSION TAG	UNP Q8RS40
E	473	HIS	-	EXPRESSION TAG	UNP Q8RS40
E	474	HIS	-	EXPRESSION TAG	UNP Q8RS40
E	475	HIS	-	EXPRESSION TAG	UNP Q8RS40
E	476	HIS	-	EXPRESSION TAG	UNP Q8RS40

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	477	HIS	-	EXPRESSION TAG	UNP Q8RS40
F	374	MET	-	INITIATING METHIONINE	UNP Q8RS40
F	470	LEU	-	EXPRESSION TAG	UNP Q8RS40
F	471	GLU	-	EXPRESSION TAG	UNP Q8RS40
F	472	HIS	-	EXPRESSION TAG	UNP Q8RS40
F	473	HIS	-	EXPRESSION TAG	UNP Q8RS40
F	474	HIS	-	EXPRESSION TAG	UNP Q8RS40
F	475	HIS	-	EXPRESSION TAG	UNP Q8RS40
F	476	HIS	-	EXPRESSION TAG	UNP Q8RS40
F	477	HIS	-	EXPRESSION TAG	UNP Q8RS40
G	374	MET	-	INITIATING METHIONINE	UNP Q8RS40
G	470	LEU	-	EXPRESSION TAG	UNP Q8RS40
G	471	GLU	-	EXPRESSION TAG	UNP Q8RS40
G	472	HIS	-	EXPRESSION TAG	UNP Q8RS40
G	473	HIS	-	EXPRESSION TAG	UNP Q8RS40
G	474	HIS	-	EXPRESSION TAG	UNP Q8RS40
G	475	HIS	-	EXPRESSION TAG	UNP Q8RS40
G	476	HIS	-	EXPRESSION TAG	UNP Q8RS40
G	477	HIS	-	EXPRESSION TAG	UNP Q8RS40
H	374	MET	-	INITIATING METHIONINE	UNP Q8RS40
H	470	LEU	-	EXPRESSION TAG	UNP Q8RS40
H	471	GLU	-	EXPRESSION TAG	UNP Q8RS40
H	472	HIS	-	EXPRESSION TAG	UNP Q8RS40
H	473	HIS	-	EXPRESSION TAG	UNP Q8RS40
H	474	HIS	-	EXPRESSION TAG	UNP Q8RS40
H	475	HIS	-	EXPRESSION TAG	UNP Q8RS40
H	476	HIS	-	EXPRESSION TAG	UNP Q8RS40
H	477	HIS	-	EXPRESSION TAG	UNP Q8RS40
I	374	MET	-	INITIATING METHIONINE	UNP Q8RS40
I	470	LEU	-	EXPRESSION TAG	UNP Q8RS40
I	471	GLU	-	EXPRESSION TAG	UNP Q8RS40
I	472	HIS	-	EXPRESSION TAG	UNP Q8RS40
I	473	HIS	-	EXPRESSION TAG	UNP Q8RS40
I	474	HIS	-	EXPRESSION TAG	UNP Q8RS40
I	475	HIS	-	EXPRESSION TAG	UNP Q8RS40
I	476	HIS	-	EXPRESSION TAG	UNP Q8RS40
I	477	HIS	-	EXPRESSION TAG	UNP Q8RS40

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	115	Total O 115 115	0	0

Continued on next page...

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
2	E	110	Total 110	O 110	0	0
2	F	85	Total 85	O 85	0	0
2	G	87	Total 87	O 87	0	0
2	H	89	Total 89	O 89	0	0
2	I	85	Total 85	O 85	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: beta-1,3-xylanase




- Molecule 1: beta-1,3-xylanase




- Molecule 1: beta-1,3-xylanase





- Molecule 1: beta-1,3-xylanase



- Molecule 1: beta-1,3-xylanase



- Molecule 1: beta-1,3-xylanase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.55Å 78.29Å 111.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.25 19.93 – 1.25	Depositor EDS
% Data completeness (in resolution range)	95.4 (20.00-1.25) 95.4 (19.93-1.25)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.31 (at 1.25Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.149 , 0.175 0.155 , 0.181	Depositor DCC
$R_{free}$ test set	7695 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	13.7	Xtrriage
Anisotropy	0.230	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 50.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4937	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	D	1.25	7/770 (0.9%)	1.12	4/1044 (0.4%)
1	E	1.35	5/770 (0.6%)	1.21	8/1044 (0.8%)
1	F	1.15	3/729 (0.4%)	1.24	6/987 (0.6%)
1	G	1.26	8/737 (1.1%)	1.34	9/998 (0.9%)
1	H	1.16	6/729 (0.8%)	1.13	7/987 (0.7%)
1	I	1.02	2/737 (0.3%)	1.05	3/998 (0.3%)
All	All	1.21	31/4472 (0.7%)	1.19	37/6058 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	E	0	1
1	G	0	1
All	All	0	2

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	380	GLU	CD-OE1	14.96	1.42	1.25
1	G	452	GLN	CD-NE2	11.45	1.61	1.32
1	D	429	ARG	CG-CD	10.88	1.79	1.51
1	G	436	LEU	CG-CD1	8.93	1.84	1.51
1	E	468	CYS	CB-SG	-8.67	1.67	1.82

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	382	CYS	O-C-N	-16.23	96.73	122.70
1	G	448	ASP	CB-CG-OD1	-14.97	104.82	118.30
1	G	448	ASP	CB-CG-OD2	13.09	130.08	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	436	LEU	CB-CG-CD1	-13.04	88.84	111.00
1	F	382	CYS	CA-C-N	11.85	143.26	117.20

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	E	380	GLU	Sidechain
1	G	452	GLN	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	751	0	685	6	0
1	E	751	0	685	3	0
1	F	712	0	652	1	0
1	G	720	0	658	3	0
1	H	712	0	652	3	0
1	I	720	0	658	4	0
2	D	115	0	0	0	0
2	E	110	0	0	1	0
2	F	85	0	0	1	0
2	G	87	0	0	3	0
2	H	89	0	0	1	0
2	I	85	0	0	0	0
All	All	4937	0	3990	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 20 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:429:ARG:CG	1:D:429:ARG:CD	1.79	1.59

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:429:ARG:CD	1:D:429:ARG:CB	2.65	0.74
1:I:389:ASN:HD21	1:I:469:ASN:H	1.37	0.73
1:I:389:ASN:ND2	1:I:468:CYS:H	1.92	0.67
1:I:389:ASN:HD22	1:I:468:CYS:H	1.50	0.59

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	90/104 (86%)	90 (100%)	0	0	100	100
1	E	90/104 (86%)	90 (100%)	0	0	100	100
1	F	85/104 (82%)	85 (100%)	0	0	100	100
1	G	86/104 (83%)	85 (99%)	1 (1%)	0	100	100
1	H	85/104 (82%)	85 (100%)	0	0	100	100
1	I	86/104 (83%)	86 (100%)	0	0	100	100
All	All	522/624 (84%)	521 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	82/93 (88%)	79 (96%)	3 (4%)	34	4
1	E	82/93 (88%)	81 (99%)	1 (1%)	71	35
1	F	77/93 (83%)	76 (99%)	1 (1%)	69	32
1	G	78/93 (84%)	76 (97%)	2 (3%)	46	8
1	H	77/93 (83%)	76 (99%)	1 (1%)	69	32
1	I	78/93 (84%)	78 (100%)	0	100	100
All	All	474/558 (85%)	466 (98%)	8 (2%)	60	23

5 of 8 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	381	ASN
1	H	429	ARG
1	G	383	GLN
1	D	469	ASN
1	F	466	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	460	ASN
1	F	438	GLN
1	H	438	GLN
1	E	438	GLN
1	G	438	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	D	92/104 (88%)	0.28	3 (3%) 46 38	11, 15, 27, 37	0
1	E	92/104 (88%)	0.26	4 (4%) 35 28	11, 16, 26, 35	1 (1%)
1	F	86/104 (82%)	0.16	2 (2%) 60 50	13, 18, 26, 36	1 (1%)
1	G	88/104 (84%)	0.44	4 (4%) 33 27	13, 19, 27, 31	6 (6%)
1	H	87/104 (83%)	0.24	5 (5%) 23 18	12, 18, 27, 37	5 (5%)
1	I	88/104 (84%)	0.26	5 (5%) 23 18	14, 19, 28, 36	6 (6%)
All	All	533/624 (85%)	0.27	23 (4%) 35 28	11, 18, 28, 37	19 (3%)

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	382	CYS	5.5
1	D	378	PRO	5.0
1	E	378	PRO	4.7
1	I	382	CYS	4.6
1	G	382	CYS	4.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.