



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 31, 2023 – 11:36 AM JST

PDB ID : 5CPK
Title : Nucleosome containing methylated Sat2L DNA
Authors : Osakabe, A.; Arimura, Y.; Adachi, F.; Maehara, K.; Ohkawa, Y.; Kurumizaka, H.
Deposited on : 2015-07-21
Resolution : 2.63 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1426 (2.66-2.62)
Clashscore	141614	1472 (2.66-2.62)
Ramachandran outliers	138981	1446 (2.66-2.62)
Sidechain outliers	138945	1446 (2.66-2.62)
RSRZ outliers	127900	1408 (2.66-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	139	 3% 55% 14% 30%
1	E	139	 % 52% 17% 29%
2	B	106	 % 61% 13% 25%
2	F	106	 % 59% 19% 22%
3	C	133	 % 61% 17% 20%
3	G	133	 % 55% 23% 22%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	D	129	 49% 22% 29%
4	H	129	 50% 20% 29%
5	I	145	 % 42% 54%
6	J	145	 % 47% 50%

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 11910 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Histone H3.1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	97	800	505	155	136	4	0	0	0
1	E	98	807	508	156	139	4	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P68431
A	-2	SER	-	expression tag	UNP P68431
A	-1	HIS	-	expression tag	UNP P68431
E	-3	GLY	-	expression tag	UNP P68431
E	-2	SER	-	expression tag	UNP P68431
E	-1	HIS	-	expression tag	UNP P68431

- Molecule 2 is a protein called Histone H4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	79	627	395	121	110	1	0	0	0
2	F	83	668	422	132	113	1	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P62805
B	-2	SER	-	expression tag	UNP P62805
B	-1	HIS	-	expression tag	UNP P62805
F	-3	GLY	-	expression tag	UNP P62805
F	-2	SER	-	expression tag	UNP P62805
F	-1	HIS	-	expression tag	UNP P62805

- Molecule 3 is a protein called Histone H2A type 1-B/E.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	106	Total	C	N	O	0	0	0
			819	517	160	142			
3	G	104	Total	C	N	O	0	0	0
			804	508	157	139			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	GLY	-	expression tag	UNP P04908
C	-2	SER	-	expression tag	UNP P04908
C	-1	HIS	-	expression tag	UNP P04908
G	-3	GLY	-	expression tag	UNP P04908
G	-2	SER	-	expression tag	UNP P04908
G	-1	HIS	-	expression tag	UNP P04908

- Molecule 4 is a protein called Histone H2B type 1-J.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	92	Total	C	N	O	S	0	0	0
			720	453	129	136	2			
4	H	91	Total	C	N	O	S	0	0	0
			708	447	125	134	2			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-3	GLY	-	expression tag	UNP P06899
D	-2	SER	-	expression tag	UNP P06899
D	-1	HIS	-	expression tag	UNP P06899
H	-3	GLY	-	expression tag	UNP P06899
H	-2	SER	-	expression tag	UNP P06899
H	-1	HIS	-	expression tag	UNP P06899


- Molecule 5 is a DNA chain called DNA (145-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
5	I	145	Total	C	N	O	P	0	0	0
			3008	1439	559	865	145			

- Molecule 6 is a DNA chain called DNA (145-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
6	J	145	2949	1421	505	878	145	0	0	0

Chain C:  61% 17% 20%




• Molecule 3: Histone H2A type 1-B/E

Chain G:  55% 23% 22%




• Molecule 4: Histone H2B type 1-J

Chain D:  49% 22% 29%




• Molecule 4: Histone H2B type 1-J

Chain H:  50% 20% 29%



• Molecule 5: DNA (145-MER)

Chain I:  42% 54%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	105.20Å 109.30Å 173.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.91 – 2.63 46.71 – 2.63	Depositor EDS
% Data completeness (in resolution range)	98.7 (19.91-2.63) 98.0 (46.71-2.63)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.84 (at 2.65Å)	Xtrriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.224 , 0.284 0.226 , 0.288	Depositor DCC
R_{free} test set	2000 reflections (3.38%)	wwPDB-VP
Wilson B-factor (Å ²)	60.4	Xtrriage
Anisotropy	0.534	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 78.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.000 for k,h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11910	wwPDB-VP
Average B, all atoms (Å ²)	104.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 5CM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	1/812 (0.1%)	0.75	0/1089
1	E	0.65	0/819	0.90	4/1097 (0.4%)
2	B	0.50	0/634	0.71	0/848
2	F	0.67	0/675	0.83	1/903 (0.1%)
3	C	0.70	0/829	0.83	1/1118 (0.1%)
3	G	0.51	0/814	0.76	1/1099 (0.1%)
4	D	0.62	0/731	0.76	0/983
4	H	0.56	0/719	0.71	0/968
5	I	0.79	2/3243 (0.1%)	1.03	3/4996 (0.1%)
6	J	0.78	1/3160 (0.0%)	1.07	3/4854 (0.1%)
All	All	0.70	4/12436 (0.0%)	0.94	13/17955 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	F	0	1
4	D	0	1
All	All	0	2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	J	1	DA	OP3-P	-9.98	1.49	1.61
5	I	1	DA	OP3-P	-9.94	1.49	1.61
5	I	50	DA	C2'-C1'	5.45	1.57	1.52
1	A	96	CYS	CB-SG	-5.05	1.73	1.81

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	131	ARG	NE-CZ-NH2	8.74	124.67	120.30
1	E	131	ARG	NE-CZ-NH1	-7.54	116.53	120.30
5	I	80	DC	OP2-P-O3'	6.17	118.78	105.20
5	I	89	DG	O5'-P-OP2	-5.98	100.32	105.70
6	J	32	DC	O4'-C1'-N1	5.85	112.10	108.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	D	32	SER	Peptide
2	F	19	ARG	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	800	0	839	22	0
1	E	807	0	844	26	0
2	B	627	0	663	14	0
2	F	668	0	719	17	0
3	C	819	0	879	26	0
3	G	804	0	861	27	0
4	D	720	0	740	33	0
4	H	708	0	727	26	0
5	I	3008	0	1652	72	0
6	J	2949	0	1658	76	0
All	All	11910	0	9582	247	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 247 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:I:133:DG:H2''	5:I:134:DA:H5''	1.65	0.78

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:50:TYR:OH	4:D:95:GLN:OE1	1.99	0.78
4:D:90:THR:H	4:D:93:GLU:HG2	1.50	0.74
5:I:58:DT:H3	6:J:88:DA:H61	1.34	0.74
5:I:126:DA:H61	6:J:20:DT:H3	1.37	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	95/139 (68%)	93 (98%)	2 (2%)	0	100	100
1	E	96/139 (69%)	93 (97%)	3 (3%)	0	100	100
2	B	77/106 (73%)	75 (97%)	2 (3%)	0	100	100
2	F	81/106 (76%)	79 (98%)	2 (2%)	0	100	100
3	C	104/133 (78%)	102 (98%)	2 (2%)	0	100	100
3	G	102/133 (77%)	101 (99%)	1 (1%)	0	100	100
4	D	90/129 (70%)	88 (98%)	2 (2%)	0	100	100
4	H	89/129 (69%)	88 (99%)	1 (1%)	0	100	100
All	All	734/1014 (72%)	719 (98%)	15 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	85/113 (75%)	83 (98%)	2 (2%)	49	67
1	E	85/113 (75%)	84 (99%)	1 (1%)	71	83
2	B	64/81 (79%)	62 (97%)	2 (3%)	40	58
2	F	69/81 (85%)	68 (99%)	1 (1%)	67	80
3	C	84/102 (82%)	81 (96%)	3 (4%)	35	52
3	G	83/102 (81%)	82 (99%)	1 (1%)	71	83
4	D	79/107 (74%)	78 (99%)	1 (1%)	69	82
4	H	77/107 (72%)	75 (97%)	2 (3%)	46	65
All	All	626/806 (78%)	613 (98%)	13 (2%)	53	71

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	D	88	THR
1	E	100	LEU
4	H	113	GLU
3	G	95	LYS
4	H	54	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	HIS
1	A	108	ASN
1	E	108	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

12 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	5CM	J	13	6,5	17,21,22	1.05	1 (5%)	24,30,33	1.30	4 (16%)
6	5CM	J	85	6,5	17,21,22	0.97	1 (5%)	24,30,33	1.45	3 (12%)
6	5CM	J	52	6,5	17,21,22	0.96	2 (11%)	24,30,33	1.38	5 (20%)
5	5CM	I	60	6,5	17,21,22	1.07	2 (11%)	24,30,33	1.30	2 (8%)
5	5CM	I	109	6,5	17,21,22	1.01	2 (11%)	24,30,33	1.47	4 (16%)
5	5CM	I	93	6,5	17,21,22	1.05	1 (5%)	24,30,33	1.53	3 (12%)
5	5CM	I	132	6,5	17,21,22	1.05	1 (5%)	24,30,33	1.17	2 (8%)
5	5CM	I	83	6,5	17,21,22	1.04	2 (11%)	24,30,33	1.48	3 (12%)
6	5CM	J	26	6,5	17,21,22	1.06	1 (5%)	24,30,33	1.55	6 (25%)
6	5CM	J	36	6,5	17,21,22	1.06	2 (11%)	24,30,33	1.42	5 (20%)
6	5CM	J	62	6,5	17,21,22	1.11	2 (11%)	24,30,33	1.53	5 (20%)
5	5CM	I	119	6,5	17,21,22	1.04	2 (11%)	24,30,33	1.79	6 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	5CM	J	13	6,5	-	2/7/21/22	0/2/2/2
6	5CM	J	85	6,5	-	7/7/21/22	0/2/2/2
6	5CM	J	52	6,5	-	4/7/21/22	0/2/2/2
5	5CM	I	60	6,5	-	4/7/21/22	0/2/2/2
5	5CM	I	109	6,5	-	1/7/21/22	0/2/2/2
5	5CM	I	93	6,5	-	6/7/21/22	0/2/2/2
5	5CM	I	132	6,5	-	0/7/21/22	0/2/2/2
5	5CM	I	83	6,5	-	0/7/21/22	0/2/2/2
6	5CM	J	26	6,5	-	6/7/21/22	0/2/2/2
6	5CM	J	36	6,5	-	4/7/21/22	0/2/2/2
6	5CM	J	62	6,5	-	6/7/21/22	0/2/2/2
5	5CM	I	119	6,5	-	3/7/21/22	0/2/2/2

The worst 5 of 19 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	J	62	5CM	C6-C5	3.38	1.40	1.34
5	I	132	5CM	C6-C5	3.32	1.40	1.34
6	J	13	5CM	C6-C5	3.14	1.39	1.34
5	I	93	5CM	C6-C5	3.07	1.39	1.34
5	I	60	5CM	C6-C5	2.98	1.39	1.34

The worst 5 of 48 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	I	119	5CM	O4'-C1'-N1	5.04	116.86	107.86
6	J	62	5CM	C5-C6-N1	-4.36	118.85	123.34
5	I	93	5CM	C5-C6-N1	-4.25	118.97	123.34
5	I	109	5CM	C5-C6-N1	-4.18	119.03	123.34
6	J	85	5CM	O4'-C1'-N1	3.88	114.80	107.86

There are no chirality outliers.

5 of 43 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	J	13	5CM	O4'-C4'-C5'-O5'
6	J	36	5CM	O4'-C1'-N1-C2
6	J	36	5CM	O4'-C1'-N1-C6
5	I	93	5CM	O4'-C4'-C5'-O5'
6	J	13	5CM	C3'-C4'-C5'-O5'

There are no ring outliers.

5 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	J	85	5CM	3	0
6	J	52	5CM	1	0
5	I	60	5CM	2	0
5	I	109	5CM	1	0
6	J	26	5CM	3	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	97/139 (69%)	0.45	4 (4%) 37 33	55, 84, 119, 139	0
1	E	98/139 (70%)	0.09	1 (1%) 82 81	35, 60, 89, 124	0
2	B	79/106 (74%)	0.37	1 (1%) 77 75	59, 77, 96, 99	0
2	F	83/106 (78%)	0.20	1 (1%) 79 77	42, 55, 74, 100	0
3	C	106/133 (79%)	0.12	0 100 100	42, 61, 98, 127	0
3	G	104/133 (78%)	0.27	1 (0%) 82 81	59, 77, 107, 115	0
4	D	92/129 (71%)	0.13	0 100 100	46, 64, 90, 120	0
4	H	91/129 (70%)	0.27	0 100 100	52, 72, 97, 107	0
5	I	139/145 (95%)	-0.11	1 (0%) 87 86	81, 141, 167, 171	0
6	J	139/145 (95%)	-0.19	2 (1%) 75 73	84, 140, 162, 169	0
All	All	1028/1304 (78%)	0.13	11 (1%) 80 78	35, 77, 156, 171	0

The worst 5 of 11 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	102	GLY	3.8
2	F	19	ARG	3.3
1	A	42	ARG	3.0
1	A	39	HIS	3.0
1	A	134	ARG	2.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	5CM	J	36	20/21	0.79	0.21	143,148,160,164	0
6	5CM	J	13	20/21	0.81	0.24	159,168,174,175	0
6	5CM	J	26	20/21	0.83	0.21	138,140,143,144	0
5	5CM	I	119	20/21	0.85	0.24	121,131,136,141	0
5	5CM	I	109	20/21	0.85	0.20	142,146,161,168	0
6	5CM	J	85	20/21	0.86	0.21	146,154,162,166	0
5	5CM	I	132	20/21	0.89	0.17	129,148,151,152	0
6	5CM	J	52	20/21	0.90	0.10	127,131,136,137	0
5	5CM	I	93	20/21	0.91	0.14	116,125,131,134	0
6	5CM	J	62	20/21	0.94	0.13	75,104,113,115	0
5	5CM	I	60	20/21	0.95	0.15	106,125,142,143	0
5	5CM	I	83	20/21	0.96	0.17	80,99,106,107	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.