

# wwPDB X-ray Structure Validation Summary Report (i)

May 15, 2020 – 09:08 am BST

PDB ID : 5CPM

Title : XenA from Pseudomonas putida in complex with NADPH4.

Authors: Knaus, T.; Paul, C.E.; Levy, C.W.; Mutti, F.G.; Hollmann, F.; Scrutton, N.S.

Deposited on : 2015-07-21

Resolution : 1.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.11

115 (20

buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

### 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$\begin{array}{c} {\rm Similar \; resolution} \\ (\#{\rm Entries, \; resolution \; range(\AA)}) \end{array}$
$R_{free}$	130704	$2936 \ (1.50 - 1.50)$
Clashscore	141614	3144 (1.50-1.50)
Ramachandran outliers	138981	3066 (1.50-1.50)
Sidechain outliers	138945	3064 (1.50-1.50)
RSRZ outliers	127900	2884 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	371	92%	5% •
1	В	371	94%	



### 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6721 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Xenobiotic reductase.


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	359	Total 2827	C 1805	N 504	O 511	S 7	0	7	0
1	В	359		C 1787		O 507	S 7	0	4	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	364	ARG	-	expression tag	UNP Q9R9V9
A	365	PRO	-	expression tag	UNP Q9R9V9
A	366	HIS	-	expression tag	UNP Q9R9V9
A	367	HIS	-	expression tag	UNP Q9R9V9
A	368	HIS	-	expression tag	UNP Q9R9V9
A	369	HIS	-	expression tag	UNP Q9R9V9
A	370	HIS	-	expression tag	UNP Q9R9V9
A	371	HIS	-	expression tag	UNP Q9R9V9
В	364	ARG	-	expression tag	UNP Q9R9V9
В	365	PRO	-	expression tag	UNP Q9R9V9
В	366	HIS	-	expression tag	UNP Q9R9V9
В	367	HIS	-	expression tag	UNP Q9R9V9
В	368	HIS	-	expression tag	UNP Q9R9V9
В	369	HIS	=	expression tag	UNP Q9R9V9
В	370	HIS	-	expression tag	UNP Q9R9V9
В	371	HIS	=	expression tag	UNP Q9R9V9


• Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NDP) (formula: C<sub>21</sub>H<sub>30</sub>N<sub>7</sub>O<sub>17</sub>P<sub>3</sub>).





Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	Λ.	1	Total	С	N	О	Р	0	0
2	$\begin{array}{c c} Z & A \end{array}$	1	33	16	2	13	2	U	
9	D	1	Total	С	N	О	Р	0	0
2	Б	1	33	16	2	13	2	U	

• Molecule 3 is 1-DEOXY-1-(7,8-DIMETHYL-2,4-DIOXO-3,4-DIHYDRO-2H-BENZO[G]P TERIDIN-1-ID-10(5H)-YL)-5-O-PHOSPHONATO-D-RIBITOL (three-letter code: FNR) (formula:  $C_{17}H_{23}N_4O_9P$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	Λ	1	Total	С	N	О	Р	0	0
)	A	1	31	17	4	9	1	U	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	D	1	Total	С	N	О	Р	0	0
3	D	1	31	17	4	9	1	U	0

### • Molecule 4 is water.


Mol	Chain	Residues	${f Atoms}$	ZeroOcc	AltConf
4	A	491	Total O 491 491	0	0
4	В	470	Total O 470 470	0	0



## 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Xenobiotic reductase





## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 2 21 21	Depositor
Cell constants	57.37Å 83.99Å 157.99Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	74.16 - 1.50	Depositor
Resolution (A)	79.00 - 1.35	EDS
% Data completeness	99.5 (74.16-1.50)	Depositor
(in resolution range)	83.7 (79.00-1.35)	EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	0.75 (at 1.35Å)	Xtriage
Refinement program	PHENIX dev_1977	Depositor
P. P.	0.154 , $0.194$	Depositor
$R, R_{free}$	0.154 , $0.194$	DCC
$R_{free}$ test set	8209 reflections $(4.91\%)$	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	6.8	Xtriage
Anisotropy	1.273	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.38 , 47.0	EDS
L-test for twinning <sup>2</sup>	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	6721	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 13.74% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

## 5 Model quality (i)

#### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, FNR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain		Bond	lengths	Bond angles		
MIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	A	0.32	0/2926	0.49	0/3989	
1	В	0.32	0/2896	0.48	0/3945	
All	All	0.32	0/5822	0.48	0/7934	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

#### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2827	0	2752	15	0
1	В	2805	0	2733	7	0
2	A	33	0	19	0	0
2	В	33	0	20	0	0
3	A	31	0	22	1	0
3	В	31	0	22	1	0
4	A	491	0	0	8	4
4	В	470	0	0	3	4
All	All	6721	0	5568	22	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.



The worst 5 of 22 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned}  ext{Interatomic} \  ext{distance} & ( ext{Å}) \end{aligned}$	$egin{array}{c}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{array}$
1:B:286:ILE:HG12	4:B:502:HOH:O	1.81	0.79
1:A:315:GLN:NE2	4:A:501:HOH:O	2.11	0.77
1:B:25:CYS:H	3:B:402:FNR:H5	1.42	0.66
1:A:293:GLU:OE2	4:A:502:HOH:O	2.14	0.65
1:A:219[B]:ARG:NH2	1:A:224:GLU:OE1	2.30	0.65

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{array}{l}  ext{Interatomic} \  ext{distance} \ ( ext{Å}) \end{array}$	$egin{array}{c}  ext{Clash} \  ext{overlap } ( ext{Å}) \end{array}$
4:A:501:HOH:O	4:B:657:HOH:O[2_587]	1.87	0.33
4:A:856:HOH:O	4:B:826:HOH:O[3_447]	1.99	0.21
4:A:847:HOH:O	4:B:623:HOH:O[2_587]	2.03	0.17
4:A:795:HOH:O	4:B:913:HOH:O[3_447]	2.16	0.04

#### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	${f Analysed}$	Favoured	Allowed	Outliers	Percentiles
1	A	364/371 (98%)	350 (96%)	11 (3%)	3 (1%)	19 5
1	В	361/371 (97%)	347 (96%)	12 (3%)	2 (1%)	25 7
All	All	725/742~(98%)	697 (96%)	23 (3%)	5 (1%)	25 6

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	302	TRP
1	A	56	GLU
1	A	302[A]	TRP

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	A	302[B]	TRP
1	В	56	GLU

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	281/286 (98%)	280 (100%)	1 (0%)	91 82
1	В	278/286 (97%)	276 (99%)	2 (1%)	84 69
All	All	559/572 (98%)	556 (100%)	3 (0%)	88 78

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	${f Res}$	Type
1	A	178	HIS
1	В	178	HIS
1	В	275	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.



#### 5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Т	Chain	Res	Res Link Bond lengths				Bond angles		
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	FNR	В	402	-	31,33,33	2.73	5 (16%)	40,50,50	1.75	5 (12%)
3	FNR	A	402	-	31,33,33	2.53	5 (16%)	40,50,50	1.76	4 (10%)
2	NDP	A	401	-	32,35,52	1.53	4 (12%)	36,52,80	1.70	7 (19%)
2	NDP	В	401	-	32,35,52	1.39	3 (9%)	38,52,80	1.84	7 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	${f Torsions}$	Rings
3	FNR	В	402	-	-	1/18/18/18	0/3/3/3
3	FNR	A	402	-	-	1/18/18/18	0/3/3/3
2	NDP	A	401	-	-	11/25/61/77	0/3/3/5
2	NDP	В	401	-	_	12/25/62/77	0/3/3/5

The worst 5 of 17 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(\mathbf{\mathring{A}})$	$\mathbf{Ideal}(\mathbf{\mathring{A}})$
3	В	402	FNR	C4A-N5	10.30	1.48	1.33
3	A	402	FNR	C4A-N5	9.43	1.46	1.33
3	В	402	FNR	C5A-N5	7.04	1.47	1.35
3	A	402	FNR	C5A-N5	6.79	1.46	1.35
3	В	402	FNR	C9A-N10	6.43	1.47	1.38

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$Observed(^o)$	$\operatorname{Ideal}({}^o)$
3	A	402	FNR	C4-N3-C2	7.24	121.25	115.14
3	В	402	FNR	C4-N3-C2	6.74	120.83	115.14

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	${f Atoms}$	$\mathbf{Z}$	$\operatorname{Observed}(^o)$	$\operatorname{Ideal}({}^o)$
2	В	401	NDP	PN-O3-PA	-6.53	110.43	132.83
2	A	401	NDP	PN-O3-PA	-5.92	112.50	132.83
3	A	402	FNR	C4A-C4-N3	-4.64	117.09	123.43

There are no chirality outliers.

5 of 25 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	NDP	C5B-O5B-PA-O1A
2	В	401	NDP	C5B-O5B-PA-O2A
2	В	401	NDP	C5D-O5D-PN-O3
2	В	401	NDP	C5D-O5D-PN-O1N
2	В	401	NDP	C5D-O5D-PN-O2N


There are no ring outliers.

2 monomers are involved in 2 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	402	FNR	1	0
3	A	402	FNR	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














### 5.7 Other polymers (i)

There are no such residues in this entry.

### 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



### 6 Fit of model and data (i)

#### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\#\mathrm{RSRZ}{>}2$	$OWAB( m \AA^2)$	Q < 0.9
1	A	359/371 (96%)	-0.74	0 100 100	8, 12, 22, 40	0
1	В	359/371 (96%)	-0.69	1 (0%) 94 95	8, 13, 23, 46	0
All	All	718/742 (96%)	-0.71	1 (0%) 95 95	8, 12, 23, 46	0

#### All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	122	ASP	2.8

#### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.


### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	NDP	В	401	33/48	0.92	0.12	10,24,106,108	0
2	NDP	A	401	33/48	0.94	0.13	8,25,97,100	0
3	FNR	В	402	31/31	0.98	0.05	7,8,10,10	0
3	FNR	A	402	31/31	0.98	0.06	5,8,10,10	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.
















## 6.5 Other polymers (i)

There are no such residues in this entry.

