



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 14, 2023 – 07:50 PM EDT

PDB ID : 8CTS
Title : Room temperature crystal structure of a K⁺ selective NaK mutant (NaK2K)
Authors : Lee, B.; White, K.I.; Socolich, M.A.; Klureza, M.A.; Henning, R.; Srajer, V.; Ranganathan, R.; Hekstra, D.
Deposited on : 2022-05-16
Resolution : 1.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3398 (1.60-1.60)
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)
RSRZ outliers	127900	3321 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	96	 91% 8% .
1	B	96	 85% 14% .
2	E	2	 50% 50%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	K	A	214	-	-	-	X

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4354 atoms, of which 2252 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Potassium channel protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	H	N	O			
1	A	95	1838	608	935	134	161	0	22	0
1	B	96	1937	636	987	144	170	0	24	0

There are 14 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	19	ALA	-	expression tag	UNP C2R3K4
A	29	VAL	ILE	conflict	UNP C2R3K4
A	66	TYR	ASP	conflict	UNP C2R3K4
A	111	LEU	-	expression tag	UNP C2R3K4
A	112	VAL	-	expression tag	UNP C2R3K4
A	113	PRO	-	expression tag	UNP C2R3K4
A	114	ARG	-	expression tag	UNP C2R3K4
B	19	ALA	-	expression tag	UNP C2R3K4
B	29	VAL	ILE	conflict	UNP C2R3K4
B	66	TYR	ASP	conflict	UNP C2R3K4
B	111	LEU	-	expression tag	UNP C2R3K4
B	112	VAL	-	expression tag	UNP C2R3K4
B	113	PRO	-	expression tag	UNP C2R3K4
B	114	ARG	-	expression tag	UNP C2R3K4

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	H	O			
2	E	2	45	12	22	11	0	0	0

- Molecule 3 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C₆H₁₄O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
3	A	1	44	12	28	4	0	1
3	A	1	44	12	28	4	0	1
3	A	1	22	6	14	2	0	0
3	A	1	22	6	14	2	0	0
3	A	1	22	6	14	2	0	0
3	A	1	22	6	14	2	0	0
3	A	1	22	6	14	2	0	0
3	A	1	22	6	14	2	0	0
3	A	1	22	6	14	2	0	0
3	A	1	22	6	14	2	0	0
3	B	1	22	6	14	2	0	0
3	B	1	22	6	14	2	0	0
3	B	1	22	6	14	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	B	1	Total	C	H	O	0	0
			22	6	14	2		
3	B	1	Total	C	H	O	0	0
			22	6	14	2		
3	B	1	Total	C	H	O	0	0
			22	6	14	2		
3	B	1	Total	C	H	O	0	1
			44	12	28	4		
3	B	1	Total	C	H	O	0	0
			22	6	14	2		
3	B	1	Total	C	H	O	0	0
			22	6	14	2		
3	B	1	Total	C	H	O	0	0
			22	6	14	2		

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	6	Total	K	0	0
			6	6		
4	B	5	Total	K	0	0
			5	5		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	23	Total	O	0	0
			23	23		
5	B	16	Total	O	0	1
			16	16		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Potassium channel protein

Chain A:  91% 8%




- Molecule 1: Potassium channel protein

Chain B:  85% 14%



- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain E:  50% 50%



4 Data and refinement statistics

Property	Value	Source
Space group	I 4	Depositor
Cell constants a, b, c, α , β , γ	68.82Å 68.82Å 90.36Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.41 – 1.60 34.41 – 1.60	Depositor EDS
% Data completeness (in resolution range)	92.7 (34.41-1.60) 92.7 (34.41-1.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.16 (at 1.60Å)	Xtrriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.141 , 0.154 0.139 , 0.152	Depositor DCC
R_{free} test set	1295 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	13.3	Xtrriage
Anisotropy	0.027	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 65.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	0.095 for -h,k,-l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4354	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.87% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, GLC, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.54	0/924	0.70	0/1260
1	B	0.52	0/971	0.66	0/1314
All	All	0.53	0/1895	0.68	0/2574

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	903	935	925	10	0
1	B	950	987	979	13	0
2	E	23	22	21	1	0
3	A	88	154	154	9	0
3	B	88	154	154	10	0
4	A	6	0	0	0	0
4	B	5	0	0	0	0
5	A	23	0	0	1	0
5	B	16	0	0	1	0
All	All	2102	2252	2233	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 27 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:86[B]:ILE:HD11	3:B:202:MPD:H32	1.74	0.68
1:B:70:SER:OG	3:B:203:MPD:H52	1.98	0.62
1:A:112[B]:VAL:HG21	3:A:207:MPD:H53	1.83	0.60
1:A:105:PRO:O	3:A:209:MPD:HM3	2.02	0.59
1:B:43[A]:SER:O	3:B:209:MPD:H53	2.02	0.58

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	116/96 (121%)	115 (99%)	1 (1%)	0	100	100
1	B	118/96 (123%)	118 (100%)	0	0	100	100
All	All	234/192 (122%)	233 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	101/85 (119%)	99 (98%)	2 (2%)	55	31
1	B	107/85 (126%)	105 (98%)	2 (2%)	57	34
All	All	208/170 (122%)	204 (98%)	4 (2%)	71	34

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	90[A]	LEU
1	A	90[B]	LEU
1	B	21[A]	ASP
1	B	21[B]	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	GLC	E	1	2	12,12,12	0.51	0	17,17,17	1.86	4 (23%)
2	GLC	E	2	2	11,11,12	0.51	0	15,15,17	2.38	4 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	E	1	2	-	0/2/22/22	0/1/1/1
2	GLC	E	2	2	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	2	GLC	C1-O5-C5	6.04	120.37	112.19
2	E	2	GLC	O5-C1-C2	4.90	118.33	110.77
2	E	1	GLC	O5-C1-C2	4.21	117.79	110.28
2	E	1	GLC	C3-C4-C5	-3.57	103.87	110.24
2	E	2	GLC	C1-C2-C3	3.37	113.80	109.67

There are no chirality outliers.

All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	E	2	GLC	C4-C5-C6-O6
2	E	2	GLC	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	2	GLC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 33 ligands modelled in this entry, 11 are monoatomic - leaving 22 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	MPD	B	208	-	7,7,7	0.32	0	9,10,10	0.35	0
3	MPD	B	203	-	7,7,7	0.26	0	9,10,10	0.31	0
3	MPD	A	201[B]	-	7,7,7	0.32	0	9,10,10	0.58	0
3	MPD	A	202[A]	-	7,7,7	0.26	0	9,10,10	0.20	0
3	MPD	B	201	-	7,7,7	0.24	0	9,10,10	0.44	0
3	MPD	A	205	-	7,7,7	0.26	0	9,10,10	0.35	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MPD	B	207[A]	-	7,7,7	0.23	0	9,10,10	0.20	0
3	MPD	B	206	-	7,7,7	0.24	0	9,10,10	0.31	0
3	MPD	A	201[A]	-	7,7,7	0.29	0	9,10,10	0.31	0
3	MPD	A	204	-	7,7,7	0.27	0	9,10,10	0.29	0
3	MPD	A	206	-	7,7,7	0.28	0	9,10,10	0.28	0
3	MPD	A	209	-	7,7,7	0.29	0	9,10,10	0.33	0
3	MPD	A	203	-	7,7,7	0.30	0	9,10,10	0.29	0
3	MPD	B	210	-	7,7,7	0.22	0	9,10,10	0.58	0
3	MPD	B	202	-	7,7,7	0.19	0	9,10,10	0.75	0
3	MPD	A	202[B]	-	7,7,7	0.22	0	9,10,10	0.34	0
3	MPD	A	208	-	7,7,7	0.22	0	9,10,10	0.56	0
3	MPD	A	207	-	7,7,7	0.29	0	9,10,10	0.27	0
3	MPD	B	207[B]	-	7,7,7	0.26	0	9,10,10	0.55	0
3	MPD	B	209	-	7,7,7	0.32	0	9,10,10	0.30	0
3	MPD	B	205	-	7,7,7	0.26	0	9,10,10	0.34	0
3	MPD	B	204	-	7,7,7	0.25	0	9,10,10	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	B	208	-	-	2/5/5/5	-
3	MPD	B	203	-	-	2/5/5/5	-
3	MPD	A	201[B]	-	-	4/5/5/5	-
3	MPD	A	202[A]	-	-	2/5/5/5	-
3	MPD	B	201	-	-	2/5/5/5	-
3	MPD	A	205	-	-	1/5/5/5	-
3	MPD	B	207[A]	-	-	2/5/5/5	-
3	MPD	B	206	-	-	2/5/5/5	-
3	MPD	A	201[A]	-	-	0/5/5/5	-
3	MPD	A	204	-	-	1/5/5/5	-
3	MPD	A	206	-	-	3/5/5/5	-
3	MPD	A	209	-	-	0/5/5/5	-
3	MPD	A	203	-	-	0/5/5/5	-
3	MPD	B	210	-	-	3/5/5/5	-
3	MPD	B	202	-	-	3/5/5/5	-
3	MPD	A	202[B]	-	-	2/5/5/5	-
3	MPD	A	208	-	-	5/5/5/5	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MPD	A	207	-	-	0/5/5/5	-
3	MPD	B	207[B]	-	-	0/5/5/5	-
3	MPD	B	209	-	-	1/5/5/5	-
3	MPD	B	205	-	-	0/5/5/5	-
3	MPD	B	204	-	-	1/5/5/5	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 36 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	202[B]	MPD	C2-C3-C4-O4
3	A	208	MPD	CM-C2-C3-C4
3	A	208	MPD	C2-C3-C4-O4
3	B	203	MPD	C2-C3-C4-C5
3	B	207[A]	MPD	C2-C3-C4-O4

There are no ring outliers.

12 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	208	MPD	2	0
3	B	203	MPD	1	0
3	B	207[A]	MPD	1	0
3	A	204	MPD	3	0
3	A	206	MPD	2	0
3	A	209	MPD	1	0
3	B	202	MPD	3	0
3	A	208	MPD	1	0
3	A	207	MPD	2	0
3	B	207[B]	MPD	1	0
3	B	209	MPD	1	0
3	B	204	MPD	1	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	95/96 (98%)	-0.50	0 100 100	3, 15, 51, 69	0
1	B	96/96 (100%)	-0.69	0 100 100	6, 16, 29, 55	0
All	All	191/192 (99%)	-0.59	0 100 100	3, 16, 44, 69	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GLC	E	1	12/12	0.73	0.38	67,122,153,160	0
2	GLC	E	2	11/12	0.81	0.14	52,65,105,167	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	K	A	214	1/1	0.09	0.41	74,74,74,74	1
3	MPD	B	208	8/8	0.57	0.21	49,76,96,120	0
3	MPD	B	209	8/8	0.66	0.31	47,61,79,85	0
3	MPD	A	202[B]	8/8	0.71	0.31	43,59,74,102	22
3	MPD	A	202[A]	8/8	0.71	0.31	52,62,76,81	22
3	MPD	A	201[A]	8/8	0.76	0.29	25,51,64,67	22
3	MPD	A	201[B]	8/8	0.76	0.29	36,55,64,64	22
3	MPD	B	203	8/8	0.76	0.33	53,89,149,149	0
4	K	A	215	1/1	0.76	0.15	72,72,72,72	1
3	MPD	A	203	8/8	0.78	0.30	57,73,87,126	0
3	MPD	B	205	8/8	0.80	0.24	64,80,97,134	0
3	MPD	B	210	8/8	0.80	0.26	42,56,67,81	0
4	K	B	212	1/1	0.80	0.06	75,75,75,75	1
3	MPD	B	206	8/8	0.81	0.23	60,73,82,92	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MPD	B	207[A]	8/8	0.81	0.26	81,100,117,175	22
3	MPD	B	207[B]	8/8	0.81	0.26	67,98,113,175	22
3	MPD	A	209	8/8	0.82	0.30	85,103,138,149	0
3	MPD	A	204	8/8	0.82	0.29	56,72,111,126	0
3	MPD	A	207	8/8	0.83	0.33	50,74,105,127	0
3	MPD	A	208	8/8	0.83	0.30	44,76,124,157	0
3	MPD	B	202	8/8	0.86	0.20	47,62,78,94	0
3	MPD	A	206	8/8	0.86	0.15	47,60,80,87	0
3	MPD	B	201	8/8	0.89	0.13	40,53,94,94	0
3	MPD	B	204	8/8	0.89	0.19	44,73,97,116	0
3	MPD	A	205	8/8	0.93	0.26	55,84,112,135	0
4	K	A	213	1/1	0.99	0.05	5,5,5,5	1
4	K	A	211	1/1	1.00	0.07	3,3,3,3	1
4	K	A	212	1/1	1.00	0.07	3,3,3,3	1
4	K	B	211	1/1	1.00	0.04	12,12,12,12	1
4	K	A	210	1/1	1.00	0.09	6,6,6,6	1
4	K	B	213	1/1	1.00	0.10	7,7,7,7	1
4	K	B	214	1/1	1.00	0.07	6,6,6,6	1
4	K	B	215	1/1	1.00	0.07	8,8,8,8	1

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around K A 214:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K A 215:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K B 212:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K A 213:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K A 211:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around K A 212:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K B 211:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K A 210:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around K B 213:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around K B 214:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.