



Full wwPDB X-ray Structure Validation Report i

Aug 21, 2020 – 09:25 PM BST

PDB ID : 4CU0
Title : Structure of bovine endothelial nitric oxide synthase heme domain in complex with (R)-6-(3-amino-2-(5-(2-(6-amino-4-methylpyridin-2-yl) ethyl)pyridin-3-yl)propyl)-4-methylpyridin-2-amine
Authors : Chreifi, G.; Li, H.; Poulos, T.L.
Deposited on : 2014-03-15
Resolution : 2.08 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.13.1
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.08 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6189 (2.10-2.06)
Clashscore	141614	6738 (2.10-2.06)
Ramachandran outliers	138981	6663 (2.10-2.06)
Sidechain outliers	138945	6664 (2.10-2.06)
RSRZ outliers	127900	6057 (2.10-2.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 6932 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, ENDOTHELIAL.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	405	Total	As	C	N	O	S	0	0	0
			3223	1	2049	568	589	16			
1	B	404	Total	As	C	N	O	S	0	0	0
			3221	1	2048	568	588	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	conflict	UNP P29473
B	100	ARG	CYS	conflict	UNP P29473

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	B	1	43	34	1	4	4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: C₉H₁₅N₅O₃).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O			
3	A	1	17	9	5	3		0	0
3	B	1	17	9	5	3		0	0

- Molecule 4 is (R)-6-(3-amino-2-(5-(2-(6-amino-4-methylpyridin-2-yl)ethyl)pyridin-3-yl)propyl)-4-methylpyridin-2-amine (three-letter code: S71) (formula: C₂₂H₂₈N₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	N	0	0
			28	22	6		
4	B	1	Total	C	N	0	0
			28	22	6		

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 6 3 3	0	0
6	B	1	Total C O 6 3 3	0	0

- Molecule 7 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	B	1	Total Zn 1 1	0	0



- Molecule 8 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	161	Total O 161 161	0	0
8	B	130	Total O 130 130	0	0


3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



- Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.91 Å 106.36 Å 156.90 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.98 – 2.08 46.93 – 2.08	Depositor EDS
% Data completeness (in resolution range)	99.3 (46.98-2.08) 99.3 (46.93-2.08)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	2.05 (at 2.08 Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R , R_{free}	0.166 , 0.214 0.174 , 0.222	Depositor DCC
R_{free} test set	2927 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	40.4	Xtriage
Anisotropy	0.404	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 46.7	EDS
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6932	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ZN, H4B, CAS, ACT, S71, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.95	0/3303	0.96	6/4497 (0.1%)
1	B	0.93	0/3300	0.98	10/4491 (0.2%)
All	All	0.94	0/6603	0.97	16/8988 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	240	ARG	NE-CZ-NH2	-11.89	114.35	120.30
1	B	240	ARG	NE-CZ-NH1	11.31	125.95	120.30
1	B	185	ARG	NE-CZ-NH1	7.72	124.16	120.30
1	A	227	ARG	NE-CZ-NH2	-7.71	116.44	120.30
1	A	100	ARG	NE-CZ-NH2	-7.68	116.46	120.30
1	A	100	ARG	NE-CZ-NH1	7.22	123.91	120.30
1	B	236	ARG	NE-CZ-NH1	-7.07	116.77	120.30
1	B	242	ASP	CB-CG-OD1	7.03	124.63	118.30
1	B	367	ARG	NE-CZ-NH1	6.07	123.34	120.30
1	A	227	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	A	331	ARG	NE-CZ-NH1	5.64	123.12	120.30
1	B	72	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	B	378	LEU	CA-CB-CG	5.32	127.54	115.30
1	B	446	ASP	CB-CG-OD1	5.32	123.09	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	414	LEU	CA-CB-CG	5.20	127.26	115.30
1	B	378	LEU	CB-CG-CD1	5.11	119.68	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	91	GLN	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3223	0	3127	23	0
1	B	3221	0	3130	30	0
2	A	43	0	30	2	0
2	B	43	0	30	8	0
3	A	17	0	15	1	0
3	B	17	0	15	2	0
4	A	28	0	28	3	0
4	B	28	0	28	3	0
5	A	4	0	3	0	0
5	B	4	0	3	0	0
6	A	6	0	8	0	0
6	B	6	0	8	1	0
7	B	1	0	0	0	0
8	A	161	0	0	3	0
8	B	130	0	0	1	0
All	All	6932	0	6425	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:384:CAS:AS	1:B:384:CAS:SG	2.46	1.33
1:B:236:ARG:HD3	1:B:351:SER:HB3	1.65	0.79
2:B:500:HEM:HBC2	2:B:500:HEM:HMC1	1.70	0.73
1:B:121:PRO:HB2	1:B:123:GLU:OE1	1.91	0.70
1:B:370:CYS:SG	1:B:378:LEU:HD13	2.36	0.65
2:B:500:HEM:HBC2	2:B:500:HEM:CMC	2.26	0.64
1:B:384:CAS:AS	1:B:384:CAS:CB	3.06	0.63
1:A:154:GLU:OE1	1:A:168:ARG:NH2	2.35	0.60
1:A:72:ARG:HB3	1:B:109:ARG:HD3	1.86	0.58
1:B:218:LYS:HD3	1:B:219:TYR:N	2.18	0.58
1:A:449:TRP:HA	3:A:600:H4B:N1	2.20	0.56
1:A:246:TRP:HB2	1:A:294:LEU:HB3	1.86	0.56
1:A:105:LEU:HD22	1:B:465:GLU:HB3	1.86	0.56
1:B:233:PHE:HB3	1:B:234:PRO:CD	2.36	0.55
1:B:478:GLN:HB2	1:B:479:PRO:HD2	1.89	0.55
1:A:340:ASN:HD22	1:A:340:ASN:H	1.55	0.53
1:B:140:ILE:C	1:B:141:LYS:HG2	2.29	0.52
1:B:308:PRO:HD2	1:B:311:LEU:HD12	1.92	0.51
2:B:500:HEM:HMC1	2:B:500:HEM:CBC	2.38	0.51
2:A:500:HEM:HBA2	4:A:800:S71:H082	1.92	0.50
1:A:138:SER:O	1:A:141:LYS:HE3	2.11	0.50
2:B:500:HEM:O1A	3:B:600:H4B:N3	2.40	0.50
8:A:2011:HOH:O	1:B:99:ARG:HD3	2.12	0.50
1:A:363:GLU:OE1	4:A:800:S71:N01	2.46	0.48
1:A:236:ARG:NH1	1:A:349:GLU:OE1	2.46	0.48
1:B:384:CAS:CE2	1:B:384:CAS:SG	3.01	0.48
1:A:423:HIS:HB2	1:B:392:THR:HB	1.93	0.48
1:A:240:ARG:HD3	1:A:241:GLY:O	2.14	0.47
2:A:500:HEM:O1D	4:A:800:S71:N21	2.47	0.47
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.95	0.47
1:A:126:LEU:O	1:A:130:ARG:HG3	2.15	0.47
1:B:455:SER:HB3	1:B:458:LEU:HD22	1.96	0.47
1:A:216:HIS:C	1:A:216:HIS:CD2	2.87	0.47
1:B:233:PHE:HB3	1:B:234:PRO:HD2	1.97	0.46
1:B:186:CYS:HB2	2:B:500:HEM:ND	2.30	0.46
1:B:323:GLU:C	1:B:323:GLU:OE1	2.55	0.46
1:B:384:CAS:AS	1:B:384:CAS:HB2	2.76	0.46
1:A:257:ARG:HG3	1:A:257:ARG:HH11	1.82	0.45
1:A:378:LEU:HB2	8:A:2120:HOH:O	2.17	0.44
2:B:500:HEM:CMC	2:B:500:HEM:CBC	2.95	0.44
1:B:378:LEU:HB2	8:B:2101:HOH:O	2.18	0.44
1:B:196:GLN:HG2	1:B:219:TYR:CZ	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:800:S71:C18	6:B:880:GOL:H32	2.47	0.44
1:B:240:ARG:HD2	1:B:241:GLY:O	2.17	0.44
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.58	0.43
1:B:249:GLN:NE2	4:B:800:S71:H12	2.34	0.43
1:A:69:LYS:HE2	1:A:69:LYS:HA	2.00	0.43
1:B:126:LEU:HD21	1:B:156:GLU:HG3	2.01	0.43
1:B:236:ARG:HG3	1:B:349:GLU:O	2.18	0.43
1:A:240:ARG:CD	1:A:241:GLY:O	2.68	0.42
1:B:455:SER:O	1:B:458:LEU:HB2	2.19	0.42
1:B:449:TRP:HA	3:B:600:H4B:N1	2.34	0.42
2:B:500:HEM:HBB2	2:B:500:HEM:HHC	2.02	0.42
1:B:340:ASN:HD22	1:B:340:ASN:H	1.68	0.42
2:B:500:HEM:O1D	4:B:800:S71:N21	2.53	0.41
1:A:121:PRO:O	1:A:122:ALA:HB3	2.21	0.41
1:A:214:CYS:SG	8:A:2064:HOH:O	2.42	0.40
1:A:101:CYS:HB3	1:B:468:ASN:HB3	2.03	0.40
1:A:249:GLN:HB2	1:A:252:ARG:HG2	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	400/443 (90%)	389 (97%)	10 (2%)	1 (0%)	41 39
1	B	399/443 (90%)	385 (96%)	13 (3%)	1 (0%)	41 39
All	All	799/886 (90%)	774 (97%)	23 (3%)	2 (0%)	41 39

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	122	ALA

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	259	GLN

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	343/375 (92%)	324 (94%)	19 (6%)	21 18
1	B	343/375 (92%)	323 (94%)	20 (6%)	20 16
All	All	686/750 (92%)	647 (94%)	39 (6%)	20 17

All (39) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	69	LYS
1	A	72	ARG
1	A	91	GLN
1	A	99	ARG
1	A	100	ARG
1	A	102	LEU
1	A	109	ARG
1	A	123	GLU
1	A	125	LEU
1	A	156	GLU
1	A	190	ILE
1	A	293	LEU
1	A	294	LEU
1	A	328	LEU
1	A	340	ASN
1	A	390	ARG
1	A	391	THR
1	A	416	LYS
1	A	458	LEU
1	B	69	LYS
1	B	100	ARG
1	B	102	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	105	LEU
1	B	110	LYS
1	B	124	GLN
1	B	141	LYS
1	B	149	GLU
1	B	156	GLU
1	B	190	ILE
1	B	202	ASP
1	B	218	LYS
1	B	225	ASN
1	B	240	ARG
1	B	323	GLU
1	B	330	LEU
1	B	340	ASN
1	B	378	LEU
1	B	389	THR
1	B	458	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	166	HIS
1	A	191	GLN
1	A	340	ASN
1	A	376	ASN
1	A	413	GLN
1	A	468	ASN
1	B	191	GLN
1	B	222	ASN
1	B	225	ASN
1	B	340	ASN
1	B	376	ASN
1	B	405	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	CAS	A	384	1	5,8,9	1.17	1 (20%)	1,9,11	0.27	0
1	CAS	B	384	1	5,8,9	1.53	2 (40%)	1,9,11	0.04	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	384	1	-	0/0/7/9	-
1	CAS	B	384	1	-	0/0/7/9	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	384	CAS	AS-CE1	2.47	2.02	1.96
1	B	384	CAS	AS-CE1	2.40	2.02	1.96
1	B	384	CAS	AS-CE2	2.06	2.01	1.96

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	B	384	CAS	4	0

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	H4B	A	600	-	16,18,18	1.36	3 (18%)	11,26,26	3.29	7 (63%)
6	GOL	B	880	-	5,5,5	0.44	0	5,5,5	1.07	0
5	ACT	A	860	-	1,3,3	1.66	0	0,3,3	0.00	-
3	H4B	B	600	-	16,18,18	1.07	2 (12%)	11,26,26	2.25	3 (27%)
6	GOL	A	880	-	5,5,5	0.69	0	5,5,5	0.88	0
5	ACT	B	860	-	1,3,3	1.98	0	0,3,3	0.00	-
2	HEM	B	500	1	27,50,50	1.26	2 (7%)	17,82,82	1.99	5 (29%)
2	HEM	A	500	1	27,50,50	1.19	3 (11%)	17,82,82	1.88	5 (29%)
4	S71	B	800	-	29,30,30	1.35	4 (13%)	36,41,41	2.55	15 (41%)
4	S71	A	800	-	29,30,30	1.03	2 (6%)	36,41,41	2.38	11 (30%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H4B	A	600	-	-	0/8/17/17	0/2/2/2
6	GOL	B	880	-	-	0/4/4/4	-
3	H4B	B	600	-	-	0/8/17/17	0/2/2/2
6	GOL	A	880	-	-	0/4/4/4	-
2	HEM	B	500	1	-	0/6/54/54	-
2	HEM	A	500	1	-	0/6/54/54	-
4	S71	B	800	-	-	1/15/15/15	0/3/3/3
4	S71	A	800	-	-	2/15/15/15	0/3/3/3

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	800	S71	C26-N21	3.43	1.40	1.34
4	A	800	S71	C15-C17	-3.22	1.45	1.52
2	B	500	HEM	C3C-C2C	-3.12	1.36	1.40
3	A	600	H4B	C7-C6	-2.99	1.49	1.52
4	B	800	S71	C23-C24	2.84	1.43	1.39
2	A	500	HEM	C3B-C2B	-2.76	1.36	1.40
2	B	500	HEM	C3B-C2B	-2.54	1.36	1.40
3	B	600	H4B	C4A-C8A	-2.50	1.36	1.41
2	A	500	HEM	C4D-C3D	2.28	1.47	1.42
2	A	500	HEM	C1B-C2B	2.28	1.47	1.42
4	B	800	S71	C20-C17	-2.24	1.51	1.54
3	B	600	H4B	C4-C4A	-2.09	1.38	1.41
3	A	600	H4B	C4A-C8A	-2.08	1.37	1.41
3	A	600	H4B	C4A-N5	-2.06	1.33	1.38
4	B	800	S71	C15-C17	-2.02	1.48	1.52
4	A	800	S71	C03-C02	2.02	1.43	1.39

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	600	H4B	C4-C4A-C8A	6.76	120.57	114.57
4	A	800	S71	C14-C15-C16	6.58	121.83	117.44
4	B	800	S71	C20-C17-C15	-6.32	102.38	111.52
4	A	800	S71	C20-C17-C15	-5.91	102.98	111.52
4	B	800	S71	C08-C06-C05	5.51	128.51	121.22
3	B	600	H4B	C4-C4A-C8A	5.35	119.32	114.57
3	A	600	H4B	C4-C4A-N5	5.08	123.38	119.12
2	B	500	HEM	CBA-CAA-C2A	-4.97	103.32	112.49
4	A	800	S71	C14-C13-C12	4.79	121.33	116.71
4	B	800	S71	C22-N21-C26	4.72	121.67	118.10
4	B	800	S71	C02-N01-C06	4.41	121.44	118.10
4	B	800	S71	C24-C25-C26	-4.30	117.50	120.32
2	A	500	HEM	CBA-CAA-C2A	-4.16	104.81	112.49
4	B	800	S71	C14-C15-C16	4.07	120.15	117.44
4	A	800	S71	C22-N21-C26	4.04	121.16	118.10
4	A	800	S71	C15-C14-C13	-3.87	115.57	120.94
2	A	500	HEM	CAA-CBA-CGA	3.75	118.97	112.67
4	B	800	S71	C15-C16-N11	-3.53	118.60	124.14
3	A	600	H4B	N2-C2-N3	3.53	122.74	117.25
3	A	600	H4B	C4-N3-C2	3.38	121.31	115.93
4	A	800	S71	N22-C22-N21	3.29	121.69	116.49
4	B	800	S71	C08-C06-N01	-3.27	111.09	115.95

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	600	H4B	N2-C2-N3	3.09	122.06	117.25
4	B	800	S71	N22-C22-N21	3.07	121.34	116.49
2	B	500	HEM	CBD-CAD-C3D	-2.89	107.15	112.48
2	B	500	HEM	CAD-CBD-CGD	2.86	117.47	112.67
3	B	600	H4B	C2-N1-C8A	2.86	120.94	114.54
3	A	600	H4B	N3-C2-N1	-2.82	120.99	125.42
4	B	800	S71	C12-N11-C16	2.73	121.21	117.48
4	A	800	S71	C14-C15-C17	-2.69	115.52	120.73
2	B	500	HEM	CAA-CBA-CGA	2.52	116.91	112.67
3	A	600	H4B	C2-N1-C8A	2.45	120.02	114.54
2	A	500	HEM	CMC-C2C-C3C	2.44	129.24	124.68
4	B	800	S71	C05-C06-N01	-2.37	120.39	122.90
3	A	600	H4B	C4A-C4-N3	-2.36	117.31	124.01
4	B	800	S71	C27-C24-C25	-2.35	117.46	120.94
4	A	800	S71	C04-C05-C06	2.34	121.85	120.32
4	A	800	S71	C02-N01-C06	-2.34	116.33	118.10
4	B	800	S71	C14-C13-C12	2.32	118.95	116.71
4	B	800	S71	C23-C22-N21	-2.30	116.82	121.75
4	A	800	S71	C24-C25-C26	-2.29	118.82	120.32
2	B	500	HEM	C1D-C2D-C3D	-2.27	105.42	107.00
4	B	800	S71	C20-C26-C25	-2.24	116.28	121.04
2	A	500	HEM	C4C-C3C-C2C	-2.20	105.36	106.90
4	A	800	S71	C09-C08-C06	2.15	117.81	112.99
2	A	500	HEM	C1D-C2D-C3D	-2.09	105.54	107.00

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	800	S71	N01-C06-C08-C09
4	A	800	S71	C05-C06-C08-C09
4	B	800	S71	C15-C17-C18-N19

There are no ring outliers.

7 monomers are involved in 15 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	600	H4B	1	0
6	B	880	GOL	1	0
3	B	600	H4B	2	0
2	B	500	HEM	8	0



Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	HEM	2	0
4	B	800	S71	3	0
4	A	800	S71	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	404/443 (91%)	0.27	25 (6%) 20 24	31, 44, 74, 124	0
1	B	403/443 (90%)	0.34	37 (9%) 9 11	32, 47, 76, 114	0
All	All	807/886 (91%)	0.30	62 (7%) 13 16	31, 45, 75, 124	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	259	GLN	6.5
1	B	110	LYS	5.0
1	A	109	ARG	4.6
1	A	259	GLN	4.2
1	B	121	PRO	4.0
1	B	449	TRP	4.0
1	B	448	ALA	3.9
1	A	123	GLU	3.5
1	B	143	SER	3.4
1	A	449	TRP	3.4
1	B	260	ASP	3.3
1	B	263	VAL	3.3
1	B	122	ALA	3.3
1	B	447	TRP	3.3
1	B	141	LYS	3.2
1	B	477	TYR	3.1
1	A	448	ALA	3.0
1	B	142	ARG	3.0
1	A	91	GLN	3.0
1	B	146	GLN	3.0
1	B	123	GLU	2.9
1	A	450	ILE	2.9
1	A	451	VAL	2.7
1	B	450	ILE	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	69	LYS	2.6
1	A	99	ARG	2.6
1	B	130	ARG	2.6
1	B	144	GLY	2.6
1	B	306	VAL	2.6
1	A	67	GLY	2.6
1	B	126	LEU	2.6
1	A	447	TRP	2.5
1	B	277	ILE	2.5
1	A	258	GLN	2.5
1	B	258	GLN	2.5
1	B	147	ALA	2.5
1	B	285	ASN	2.4
1	A	108	PRO	2.4
1	A	153	GLN	2.4
1	B	281	TRP	2.4
1	A	141	LYS	2.3
1	A	257	ARG	2.3
1	B	109	ARG	2.3
1	B	91	GLN	2.3
1	A	280	GLY	2.3
1	B	103	GLY	2.3
1	A	122	ALA	2.3
1	B	445	ALA	2.3
1	A	186	CYS	2.2
1	B	184	PRO	2.2
1	A	462	PHE	2.2
1	B	323	GLU	2.2
1	B	270	VAL	2.1
1	B	257	ARG	2.1
1	B	262	SER	2.0
1	A	126	LEU	2.0
1	B	261	GLY	2.0
1	B	223	ARG	2.0
1	A	187	VAL	2.0
1	B	239	GLY	2.0
1	A	127	SER	2.0
1	A	277	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CAS	B	384	9/10	0.96	0.10	52,54,78,88	0
1	CAS	A	384	9/10	0.98	0.10	41,43,76,77	0

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.


6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	GOL	B	880	6/6	0.92	0.25	51,62,63,79	0
5	ACT	B	860	4/4	0.92	0.37	53,55,57,60	0
4	S71	B	800	28/28	0.92	0.19	35,48,62,66	0
6	GOL	A	880	6/6	0.93	0.18	52,60,61,63	0
4	S71	A	800	28/28	0.94	0.18	36,49,64,68	0
5	ACT	A	860	4/4	0.95	0.15	45,47,56,58	0
3	H4B	B	600	17/17	0.97	0.22	34,37,42,45	0
2	HEM	A	500	43/43	0.98	0.18	32,35,41,53	0
3	H4B	A	600	17/17	0.98	0.21	33,36,39,40	0
2	HEM	B	500	43/43	0.98	0.17	29,36,47,55	0
7	ZN	B	900	1/1	0.99	0.07	47,47,47,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.