



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 26, 2023 – 02:46 PM EDT

PDB ID : 6CWJ
Title : Crystal structures of cyanuric acid hydrolase from *Moorella thermoacetica* complexed with 1,3-Acetone Dicarboxylic Acid
Authors : Shi, K.; Aihara, H.
Deposited on : 2018-03-30
Resolution : 2.25 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	363	 2% 92% 7%
1	B	363	 0% 92% 8%
1	C	363	 0% 92% 8%
1	D	363	 3% 91% 9%

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 11453 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyanuric acid amidohydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	363	2708	1675	485	534	14	0	2	0
1	B	363	2691	1665	482	530	14	0	0	0
1	C	363	2700	1671	484	531	14	0	1	0
1	D	363	2700	1671	484	531	14	0	1	0

There are 64 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	HIS	-	expression tag	UNP Q2RGM7
A	103	ALA	GLN	engineered mutation	UNP Q2RGM7
A	104	ALA	GLU	engineered mutation	UNP Q2RGM7
A	107	ALA	LYS	engineered mutation	UNP Q2RGM7
A	279	ILE	LEU	engineered mutation	UNP Q2RGM7
A	280	ARG	LYS	engineered mutation	UNP Q2RGM7
A	281	SER	PHE	engineered mutation	UNP Q2RGM7
A	?	-	CYS	deletion	UNP Q2RGM7
A	?	-	CYS	deletion	UNP Q2RGM7
A	?	-	PRO	deletion	UNP Q2RGM7
A	?	-	PRO	deletion	UNP Q2RGM7
A	?	-	ALA	deletion	UNP Q2RGM7
A	283	ASP	GLU	engineered mutation	UNP Q2RGM7
A	285	MET	LEU	engineered mutation	UNP Q2RGM7
A	286	ASP	ALA	engineered mutation	UNP Q2RGM7
A	287	ARG	LYS	engineered mutation	UNP Q2RGM7
B	0	HIS	-	expression tag	UNP Q2RGM7
B	103	ALA	GLN	engineered mutation	UNP Q2RGM7
B	104	ALA	GLU	engineered mutation	UNP Q2RGM7
B	107	ALA	LYS	engineered mutation	UNP Q2RGM7
B	279	ILE	LEU	engineered mutation	UNP Q2RGM7

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	280	ARG	LYS	engineered mutation	UNP Q2RGM7
B	281	SER	PHE	engineered mutation	UNP Q2RGM7
B	?	-	CYS	deletion	UNP Q2RGM7
B	?	-	CYS	deletion	UNP Q2RGM7
B	?	-	PRO	deletion	UNP Q2RGM7
B	?	-	PRO	deletion	UNP Q2RGM7
B	?	-	ALA	deletion	UNP Q2RGM7
B	283	ASP	GLU	engineered mutation	UNP Q2RGM7
B	285	MET	LEU	engineered mutation	UNP Q2RGM7
B	286	ASP	ALA	engineered mutation	UNP Q2RGM7
B	287	ARG	LYS	engineered mutation	UNP Q2RGM7
C	0	HIS	-	expression tag	UNP Q2RGM7
C	103	ALA	GLN	engineered mutation	UNP Q2RGM7
C	104	ALA	GLU	engineered mutation	UNP Q2RGM7
C	107	ALA	LYS	engineered mutation	UNP Q2RGM7
C	279	ILE	LEU	engineered mutation	UNP Q2RGM7
C	280	ARG	LYS	engineered mutation	UNP Q2RGM7
C	281	SER	PHE	engineered mutation	UNP Q2RGM7
C	?	-	CYS	deletion	UNP Q2RGM7
C	?	-	CYS	deletion	UNP Q2RGM7
C	?	-	PRO	deletion	UNP Q2RGM7
C	?	-	PRO	deletion	UNP Q2RGM7
C	?	-	ALA	deletion	UNP Q2RGM7
C	283	ASP	GLU	engineered mutation	UNP Q2RGM7
C	285	MET	LEU	engineered mutation	UNP Q2RGM7
C	286	ASP	ALA	engineered mutation	UNP Q2RGM7
C	287	ARG	LYS	engineered mutation	UNP Q2RGM7
D	0	HIS	-	expression tag	UNP Q2RGM7
D	103	ALA	GLN	engineered mutation	UNP Q2RGM7
D	104	ALA	GLU	engineered mutation	UNP Q2RGM7
D	107	ALA	LYS	engineered mutation	UNP Q2RGM7
D	279	ILE	LEU	engineered mutation	UNP Q2RGM7
D	280	ARG	LYS	engineered mutation	UNP Q2RGM7
D	281	SER	PHE	engineered mutation	UNP Q2RGM7
D	?	-	CYS	deletion	UNP Q2RGM7
D	?	-	CYS	deletion	UNP Q2RGM7
D	?	-	PRO	deletion	UNP Q2RGM7
D	?	-	PRO	deletion	UNP Q2RGM7
D	?	-	ALA	deletion	UNP Q2RGM7
D	283	ASP	GLU	engineered mutation	UNP Q2RGM7
D	285	MET	LEU	engineered mutation	UNP Q2RGM7
D	286	ASP	ALA	engineered mutation	UNP Q2RGM7

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	287	ARG	LYS	engineered mutation	UNP Q2RGM7

- Molecule 2 is MALONATE ION (three-letter code: MLI) (formula: $C_3H_2O_4$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
2	A	1	7	3	4	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	A	1	1	1	0	0
3	B	1	1	1	0	0
3	C	1	1	1	0	0
3	D	1	1	1	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 4 2 2	0	0
4	A	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0

- Molecule 5 is 3-oxopentanedioic acid (three-letter code: 6JN) (formula: C₅H₆O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			10	5	5		
5	C	1	Total	C	O	0	0
			10	5	5		
5	D	1	Total	C	O	0	0
			10	5	5		

- Molecule 6 is 1,3-PROPANDIOL (three-letter code: PDO) (formula: C₃H₈O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			5	3	2		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	161	Total	O	0	0
			161	161		
7	B	151	Total	O	0	0
			151	151		
7	C	166	Total	O	0	0
			166	166		
7	D	114	Total	O	0	0
			114	114		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyanuric acid amidohydrolase




- Molecule 1: Cyanuric acid amidohydrolase



- Molecule 1: Cyanuric acid amidohydrolase



- Molecule 1: Cyanuric acid amidohydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	79.14Å 88.84Å 190.51Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	95.25 – 2.25 95.25 – 2.25	Depositor EDS
% Data completeness (in resolution range)	98.5 (95.25-2.25) 99.4 (95.25-2.25)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.45 (at 2.25Å)	Xtrriage
Refinement program	PHENIX dev_3082	Depositor
R, R_{free}	0.186 , 0.224 0.185 , 0.224	Depositor DCC
R_{free} test set	3179 reflections (4.97%)	wwPDB-VP
Wilson B-factor (Å ²)	29.7	Xtrriage
Anisotropy	0.199	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.42$, $\langle L^2 \rangle = 0.24$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	11453	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MLI, ACT, PDO, CA, 6JN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/2741	0.47	0/3709
1	B	0.27	0/2724	0.46	0/3687
1	C	0.27	0/2733	0.46	0/3698
1	D	0.26	0/2733	0.45	0/3698
All	All	0.27	0/10931	0.46	0/14792

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2708	0	2727	21	0
1	B	2691	0	2712	16	0
1	C	2700	0	2724	18	0
1	D	2700	0	2724	20	0
2	A	7	0	2	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	8	0	6	0	0
4	D	8	0	6	0	0
5	B	10	0	0	0	0
5	C	10	0	0	2	0
5	D	10	0	0	0	0
6	B	5	0	8	0	0
7	A	161	0	0	9	0
7	B	151	0	0	6	0
7	C	166	0	0	9	0
7	D	114	0	0	6	0
All	All	11453	0	10909	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 74 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:210:ASP:OD2	7:A:501:HOH:O	1.84	0.95
1:C:258:HIS:O	7:C:501:HOH:O	1.86	0.93
1:B:27:GLY:O	7:B:501:HOH:O	1.88	0.89
1:C:150:ASP:OD1	7:C:502:HOH:O	1.91	0.88
1:D:185:THR:OG1	7:D:501:HOH:O	1.92	0.87

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	363/363 (100%)	349 (96%)	14 (4%)	0	100 100
1	B	361/363 (99%)	347 (96%)	14 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	362/363 (100%)	346 (96%)	16 (4%)	0	100	100
1	D	362/363 (100%)	349 (96%)	13 (4%)	0	100	100
All	All	1448/1452 (100%)	1391 (96%)	57 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	288/286 (101%)	285 (99%)	3 (1%)	76	84
1	B	286/286 (100%)	280 (98%)	6 (2%)	53	62
1	C	287/286 (100%)	282 (98%)	5 (2%)	60	71
1	D	287/286 (100%)	285 (99%)	2 (1%)	84	90
All	All	1148/1144 (100%)	1132 (99%)	16 (1%)	67	76

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	239	ASP
1	C	319	THR
1	B	292	LEU
1	C	292	LEU
1	B	282	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	222	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 13 ligands modelled in this entry, 4 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	PDO	B	403	-	4,4,4	0.41	0	3,3,3	0.40	0
5	6JN	B	401	-	9,9,9	1.23	0	11,11,11	1.12	0
4	ACT	A	403	-	3,3,3	0.88	0	3,3,3	0.83	0
4	ACT	D	401	-	3,3,3	0.90	0	3,3,3	0.87	0
2	MLI	A	401	-	6,6,6	1.47	0	7,7,7	1.15	0
5	6JN	C	401	-	9,9,9	1.29	0	11,11,11	1.20	1 (9%)
4	ACT	D	404	-	3,3,3	0.90	0	3,3,3	0.85	0
5	6JN	D	402	-	9,9,9	1.22	0	11,11,11	1.05	0
4	ACT	A	404	-	3,3,3	0.92	0	3,3,3	0.89	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PDO	B	403	-	-	0/2/2/2	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	6JN	B	401	-	-	1/8/8/8	-
2	MLI	A	401	-	-	2/4/4/4	-
5	6JN	C	401	-	-	3/8/8/8	-
5	6JN	D	402	-	-	3/8/8/8	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	401	6JN	OAI-CAB-CAA	2.00	120.93	114.54

There are no chirality outliers.

5 of 9 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	C	401	6JN	CAC-CAD-CAE-OAF
5	D	402	6JN	CAA-CAC-CAD-CAE
5	D	402	6JN	OAH-CAC-CAD-CAE
5	D	402	6JN	CAC-CAA-CAB-OAI
5	C	401	6JN	CAB-CAA-CAC-OAH

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	401	6JN	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	363/363 (100%)	-0.34	6 (1%) 70 73	22, 35, 67, 111	0
1	B	363/363 (100%)	-0.38	4 (1%) 80 82	21, 34, 69, 90	1 (0%)
1	C	363/363 (100%)	-0.42	2 (0%) 89 89	22, 35, 63, 88	1 (0%)
1	D	363/363 (100%)	0.02	11 (3%) 50 53	25, 52, 89, 119	0
All	All	1452/1452 (100%)	-0.28	23 (1%) 72 74	21, 37, 78, 119	2 (0%)

The worst 5 of 23 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	106	ALA	5.4
1	A	105	PRO	4.7
1	D	0	HIS	4.5
1	A	108	PRO	4.5
1	A	104	ALA	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	ACT	D	404	4/4	0.72	0.26	75,75,75,76	0
4	ACT	A	403	4/4	0.76	0.17	63,64,65,66	0
4	ACT	D	401	4/4	0.79	0.22	46,49,49,50	0
4	ACT	A	404	4/4	0.82	0.14	60,61,61,61	0
6	PDO	B	403	5/5	0.82	0.17	51,52,55,57	0
5	6JN	D	402	10/10	0.94	0.15	41,43,45,46	0
5	6JN	C	401	10/10	0.95	0.11	31,36,46,50	0
3	CA	D	403	1/1	0.95	0.05	53,53,53,53	1
5	6JN	B	401	10/10	0.95	0.14	38,40,41,42	0
3	CA	B	402	1/1	0.96	0.07	28,28,28,28	1
3	CA	C	402	1/1	0.97	0.04	34,34,34,34	1
3	CA	A	402	1/1	0.98	0.07	29,29,29,29	1
2	MLI	A	401	7/7	0.98	0.09	30,31,34,36	0

6.5 Other polymers [i](#)

There are no such residues in this entry.