

wwPDB X-ray Structure Validation Summary Report (i)

May 16, 2020 – 08:27 pm BST

PDB ID	:	5D57
Title	:	In meso X-ray crystallography structure of diacylglycerol kinase, DgkA, at 100
		К
Authors	:	Huang, CY.; Howe, N.; Olieric, V.; Wang, M.; Caffrey, M.
Deposited on		
Resolution	:	2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
$\operatorname{CCP4}$:	$7.0.044 (\mathrm{Gargrove})$
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$		
R_{free}	130704	3140(2.80-2.80)		
Clashscore	141614	3569(2.80-2.80)		
Ramachandran outliers	138981	3498 (2.80-2.80)		
Sidechain outliers	138945	3500 (2.80-2.80)		
RSRZ outliers	127900	3078 (2.80-2.80)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	А	130	3% 7 6%			13%	11%		
			%			1370	1170		
1	В	130	65%		11%	24%	ó		
1	С	130	4% 65%		5% •	29%			
1	D	130	75%			8%	17%		
1	Е	130	4% 55%	12%	•	32%			
1	F	130	58%	9%	•	32%			



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	78M	А	203	-	-	-	Х
2	78M	А	205	-	-	Х	-
2	78M	В	202	-	-	-	Х
2	78M	С	201	-	-	-	Х



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4884 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	А	116	Total	С	Ν	Ο	\mathbf{S}	0	0	0
	А	110	886	580	149	154	3	0	0	0
1	В	99	Total	С	Ν	Ο	S	0	1	0
	D	99	756	494	123	136	3	0		0
1	С	92	Total	С	Ν	Ο	S	0	0	0
	U	92	696	455	114	124	3	0		
1	D	108	Total	С	Ν	Ο	S	0	0	0
	D	108	823	538	137	145	3	0		
1	Е	89	Total	С	Ν	Ο	S	0	0	0
		09	669	437	108	121	3	0	0	0
1	F	88	Total	С	Ν	Ο	S	0	0	0
	Ľ	00	660	432	106	119	3		0	U

• Molecule 1 is a protein called Diacylglycerol kinase.

There are 96 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	-8	GLY	-	expression tag	UNP P0ABN1
А	-7	HIS	-	expression tag	UNP P0ABN1
А	-6	HIS	-	expression tag	UNP P0ABN1
А	-5	HIS	-	expression tag	UNP P0ABN1
А	-4	HIS	-	expression tag	UNP P0ABN1
А	-3	HIS	-	expression tag	UNP P0ABN1
А	-2	HIS	-	expression tag	UNP P0ABN1
А	-1	GLU	-	expression tag	UNP P0ABN1
А	0	LEU	-	expression tag	UNP P0ABN1
А	41	CYS	ALA	engineered mutation	UNP P0ABN1
А	46	ALA	CYS	engineered mutation	UNP P0ABN1
А	53	VAL	ILE	engineered mutation	UNP P0ABN1
A	70	LEU	ILE	engineered mutation	UNP P0ABN1
А	96	LEU	MET	engineered mutation	UNP P0ABN1
А	107	ASP	VAL	engineered mutation	UNP P0ABN1
А	113	ALA	CYS	engineered mutation	UNP P0ABN1
В	-8	GLY	-	expression tag	UNP P0ABN1



5D57

Continued from previous page							
Chain	Residue	Modelled	Actual	Comment	Reference		
В	-7	HIS	-	expression tag	UNP P0ABN1		
В	-6	HIS	-	expression tag	UNP P0ABN1		
В	-5	HIS	-	expression tag	UNP P0ABN1		
В	-4	HIS	-	expression tag	UNP P0ABN1		
В	-3	HIS	-	expression tag	UNP P0ABN1		
В	-2	HIS	-	expression tag	UNP P0ABN1		
В	-1	GLU	-	expression tag	UNP P0ABN1		
В	0	LEU	-	expression tag	UNP P0ABN1		
В	41	CYS	ALA	engineered mutation	UNP P0ABN1		
В	46	ALA	CYS	engineered mutation	UNP P0ABN1		
В	53	VAL	ILE	engineered mutation	UNP P0ABN1		
В	70	LEU	ILE	engineered mutation	UNP P0ABN1		
В	96	LEU	MET	engineered mutation	UNP P0ABN1		
В	107	ASP	VAL	engineered mutation	UNP P0ABN1		
В	113	ALA	CYS	engineered mutation	UNP P0ABN1		
С	-8	GLY	-	expression tag	UNP P0ABN1		
С	-7	HIS	-	expression tag	UNP P0ABN1		
С	-6	HIS	-	expression tag	UNP P0ABN1		
С	-5	HIS	-	expression tag	UNP P0ABN1		
С	-4	HIS	-	expression tag	UNP P0ABN1		
С	-3	HIS	-	expression tag	UNP P0ABN1		
С	-2	HIS	-	expression tag	UNP P0ABN1		
С	-1	GLU	-	expression tag	UNP P0ABN1		
С	0	LEU	-	expression tag	UNP P0ABN1		
С	41	CYS	ALA	engineered mutation	UNP P0ABN1		
С	46	ALA	CYS	engineered mutation	UNP P0ABN1		
С	53	VAL	ILE	engineered mutation	UNP P0ABN1		
С	70	LEU	ILE	engineered mutation	UNP P0ABN1		
С	96	LEU	MET	engineered mutation	UNP P0ABN1		
С	107	ASP	VAL	engineered mutation	UNP P0ABN1		
С	113	ALA	CYS	engineered mutation	UNP P0ABN1		
D	-8	GLY	-	expression tag	UNP P0ABN1		
D	-7	HIS	-	expression tag	UNP P0ABN1		
D	-6	HIS	-	expression tag	UNP P0ABN1		
D	-5	HIS	-	expression tag	UNP P0ABN1		
D	-4	HIS	-	expression tag	UNP P0ABN1		
D	-3	HIS	-	expression tag	UNP P0ABN1		
D	-2	HIS	-	expression tag	UNP P0ABN1		
D	-1	GLU	-	expression tag	UNP P0ABN1		
D	0	LEU	-	expression tag	UNP P0ABN1		
D	41	CYS	ALA	engineered mutation	UNP P0ABN1		
D	46	ALA	CYS	engineered mutation	UNP P0ABN1		

Continued from previous page...




Chain	Residue	Modelled	Actual	Comment	Reference
D	53	VAL	ILE	engineered mutation	UNP P0ABN1
D	70	LEU	ILE	engineered mutation	UNP P0ABN1
D	96	LEU	MET	engineered mutation	UNP P0ABN1
D	107	ASP	VAL	engineered mutation	UNP P0ABN1
D	113	ALA	CYS	engineered mutation	UNP P0ABN1
Е	-8	GLY	-	expression tag	UNP P0ABN1
Е	-7	HIS	-	expression tag	UNP P0ABN1
Е	-6	HIS	-	expression tag	UNP P0ABN1
Е	-5	HIS	-	expression tag	UNP P0ABN1
E	-4	HIS	-	expression tag	UNP P0ABN1
E	-3	HIS	-	expression tag	UNP P0ABN1
Е	-2	HIS	-	expression tag	UNP P0ABN1
E	-1	GLU	-	expression tag	UNP P0ABN1
E	0	LEU	-	expression tag	UNP P0ABN1
E	41	CYS	ALA	engineered mutation	UNP P0ABN1
E	46	ALA	CYS	engineered mutation	UNP P0ABN1
Е	53	VAL	ILE	engineered mutation	UNP P0ABN1
E	70	LEU	ILE	engineered mutation	UNP P0ABN1
Е	96	LEU	MET	engineered mutation	UNP P0ABN1
E	107	ASP	VAL	engineered mutation	UNP P0ABN1
Е	113	ALA	CYS	engineered mutation	UNP P0ABN1
F	-8	GLY	-	expression tag	UNP P0ABN1
F	-7	HIS	-	expression tag	UNP P0ABN1
F	-6	HIS	-	expression tag	UNP P0ABN1
F	-5	HIS	-	expression tag	UNP P0ABN1
F	-4	HIS	-	expression tag	UNP P0ABN1
F	-3	HIS	-	expression tag	UNP P0ABN1
F	-2	HIS	-	expression tag	UNP P0ABN1
F	-1	GLU	-	expression tag	UNP P0ABN1
F	0	LEU	-	expression tag	UNP P0ABN1
F	41	CYS	ALA	engineered mutation	UNP P0ABN1
F	46	ALA	CYS	engineered mutation	UNP P0ABN1
F	53	VAL	ILE	engineered mutation	UNP P0ABN1
F	70	LEU	ILE	engineered mutation	UNP P0ABN1
F	96	LEU	MET	engineered mutation	UNP P0ABN1
F	107	ASP	VAL	engineered mutation	UNP P0ABN1
F	113	ALA	CYS	engineered mutation	UNP P0ABN1

Continued from previous page...

• Molecule 2 is (2S)-2,3-DIHYDROXYPROPYL(7Z)-PENTADEC-7-ENOATE (three-letter code: 78M) (formula: C₁₈H₃₄O₄).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total C O 22 18 4	0	0
2	А	1	Total C O 22 18 4	0	0
2	А	1	Total C O 22 18 4	0	0
2	А	1	Total C O 22 18 4	0	0
2	А	1	Total C O 22 18 4	0	0
2	А	1	Total C O 22 18 4	0	0
2	В	1	Total C O 22 18 4	0	0
2	В	1	Total C O 22 18 4	0	0
2	В	1	Total C O 22 18 4	0	0
2	В	1	Total C O 22 18 4	0	0
2	С	1	Total C O 22 18 4	0	0
2	С	1	Total C O 22 18 4	0	0
2	D	1	Total C O 22 18 4	0	0
2	D	1	Total C O 22 18 4	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total C O 22 18 4	0	0
2	D	1	Total C O 22 18 4	0	0


• Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	7	Total O 7 7	0	0
3	В	7	$\begin{array}{cc} {\rm Total} & {\rm O} \\ 7 & 7 \end{array}$	0	0
3	С	4	$\begin{array}{cc} \text{Total} & \text{O} \\ 4 & 4 \end{array}$	0	0
3	D	5	Total O 5 5	0	0
3	Ε	11	Total O 11 11	0	0
3	F	8	Total O 8 8	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Diacylglycerol kinase



• Molecule 1: Diacylglycerol kinase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	75.32Å 91.34Å 143.36Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.67 - 2.80	Depositor
Resolution (A)	45.67 - 2.80	EDS
% Data completeness	99.5 (45.67-2.80)	Depositor
(in resolution range)	92.4(45.67-2.80)	EDS
R _{merge}	(Not available)	Depositor
R _{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.52 (at 2.81 \text{\AA})$	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
D D.	0.221 , 0.266	Depositor
R, R_{free}	0.224 , 0.266	DCC
R_{free} test set	1251 reflections $(5.02%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	48.5	Xtriage
Anisotropy	0.232	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.30, 60.9	EDS
L-test for twinning ²	$ \langle L \rangle = 0.49, \langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4884	wwPDB-VP
Average B, all atoms $(Å^2)$	58.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.27% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: $78\mathrm{M}$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Mol Chain		Bond lengths		angles
	Ullalli	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.25	0/902	0.41	0/1229
1	В	0.23	0/771	0.36	0/1054
1	С	0.24	0/706	0.36	0/964
1	D	0.23	0/838	0.36	0/1144
1	Е	0.25	0/678	0.46	0/927
1	F	0.22	0/669	0.39	0/915
All	All	0.24	0/4564	0.39	0/6233

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	886	0	922	18	0
1	В	756	0	784	12	0
1	С	696	0	730	7	0
1	D	823	0	850	8	0
1	Ε	669	0	703	11	0
1	F	660	0	695	7	0
2	А	132	0	204	19	0



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	В	88	0	136	5	0
2	С	44	0	68	0	0
2	D	88	0	136	5	0
3	А	7	0	0	0	0
3	В	7	0	0	0	0
3	С	4	0	0	0	0
3	D	5	0	0	1	0
3	Ε	11	0	0	0	0
3	F	8	0	0	0	0
All	All	4884	0	5228	67	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 67 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:39:LEU:HD22	1:E:62:VAL:HG12	1.65	0.79
2:A:205:78M:H201	2:B:201:78M:H18	1.69	0.74
1:B:81:ARG:NH2	1:C:85:GLU:OE2	2.23	0.72
1:A:55:ARG:NH2	2:A:205:78M:O19	2.27	0.68
1:B:26:ILE:HD11	2:D:203:78M:H7	1.77	0.66

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	114/130~(88%)	112~(98%)	2(2%)	0	100	100
1	В	98/130~(75%)	97~(99%)	1 (1%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	С	90/130~(69%)	90 (100%)	0	0	100	100
1	D	106/130~(82%)	106~(100%)	0	0	100	100
1	Е	87/130~(67%)	83~(95%)	3 (3%)	1 (1%)	14	41
1	F	86/130~(66%)	85~(99%)	1 (1%)	0	100	100
All	All	581/780 (74%)	573~(99%)	7 (1%)	1 (0%)	47	78

Continued from previous page...

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	Ε	50	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percent	tiles
1	А	90/102~(88%)	90~(100%)	0	100	100
1	В	79/102~(78%)	79~(100%)	0	100	100
1	С	74/102~(72%)	73~(99%)	1 (1%)	67	90
1	D	84/102~(82%)	83~(99%)	1 (1%)	71	92
1	Ε	72/102~(71%)	68~(94%)	4~(6%)	21	51
1	F	71/102~(70%)	68~(96%)	3~(4%)	30	63
All	All	470/612 (77%)	461 (98%)	9 (2%)	57	85

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Е	117	TRP
1	F	66	MET
1	F	38	VAL
1	Е	36	VAL
1	Е	119	HIS



Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	D	87	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	Bo	ond leng	ths	В	ond ang	les
MOI	Type	Unam	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	78M	А	201	-	21,21,21	0.89	1 (4%)	$22,\!22,\!22$	0.77	1 (4%)
2	78M	В	203	-	21,21,21	1.02	1 (4%)	$22,\!22,\!22$	0.79	1 (4%)
2	78M	А	205	-	21,21,21	0.94	1 (4%)	22,22,22	0.84	1 (4%)
2	78M	D	203	-	21,21,21	0.87	1 (4%)	$22,\!22,\!22$	0.80	1 (4%)
2	78M	D	201	-	21,21,21	0.89	1 (4%)	$22,\!22,\!22$	0.81	1 (4%)
2	78M	D	204	-	21,21,21	0.85	1 (4%)	$22,\!22,\!22$	0.85	1 (4%)
2	78M	В	204	-	21,21,21	0.91	1 (4%)	$22,\!22,\!22$	0.82	1 (4%)
2	78M	А	203	-	21,21,21	0.93	1 (4%)	22,22,22	0.80	1 (4%)
2	78M	В	201	-	21,21,21	0.91	1 (4%)	22,22,22	0.70	1 (4%)



Mal	Mol Type Chain		Res	Link	Bo	ond leng	\mathbf{ths}	Bond angles		
	туре	Cham	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
2	78M	В	202	-	21,21,21	0.87	1 (4%)	$22,\!22,\!22$	0.74	1 (4%)
2	78M	А	202	-	21,21,21	0.90	1 (4%)	22,22,22	0.79	1 (4%)
2	78M	D	202	-	21,21,21	1.01	1 (4%)	22,22,22	0.88	1 (4%)
2	78M	С	202	-	21,21,21	0.89	1 (4%)	22,22,22	0.79	1 (4%)
2	78M	А	206	-	21,21,21	0.85	1 (4%)	$22,\!22,\!22$	0.81	1 (4%)
2	78M	С	201	-	21,21,21	0.89	1 (4%)	22,22,22	0.85	1 (4%)
2	78M	А	204	-	21,21,21	0.89	1 (4%)	22,22,22	0.84	1 (4%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	78M	А	201	-	-	13/21/21/21	-
2	78M	В	203	-	-	10/21/21/21	-
2	78M	А	205	-	-	10/21/21/21	-
2	78M	D	203	-	-	8/21/21/21	-
2	78M	D	201	-	-	8/21/21/21	-
2	78M	D	204	-	-	8/21/21/21	-
2	78M	В	204	-	-	11/21/21/21	-
2	78M	А	203	-	-	12/21/21/21	-
2	78M	В	201	-	-	10/21/21/21	-
2	78M	В	202	-	-	10/21/21/21	-
2	78M	А	202	-	-	11/21/21/21	-
2	78M	D	202	-	-	12/21/21/21	-
2	78M	С	202	-	-	11/21/21/21	-
2	78M	А	206	-	-	9/21/21/21	-
2	78M	С	201	-	-	11/21/21/21	-
2	78M	А	204	_	_	8/21/21/21	-

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	D	202	78M	O2-C1	3.12	1.42	1.33
2	В	203	78M	O2-C1	3.10	1.42	1.33
2	А	205	78 M	O2-C1	3.00	1.42	1.33



Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
2	А	203	78M	O2-C1	2.96	1.42	1.33
2	А	202	78M	O2-C1	2.91	1.41	1.33

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	D	202	78M	O2-C1-C2	3.27	122.18	111.91
2	А	205	78M	O2-C1-C2	2.96	121.19	111.91
2	D	204	78M	O2-C1-C2	2.88	120.96	111.91
2	С	201	78M	O2-C1-C2	2.86	120.89	111.91
2	А	204	78 M	O2-C1-C2	2.86	120.87	111.91

There are no chirality outliers.

5 of 162 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	203	78M	O2-C17-C18-O19
2	А	205	78M	C17-C18-C20-O21
2	D	203	78 M	C17-C18-C20-O21
2	D	201	78 M	C2-C1-O2-C17
2	А	203	78M	C17-C18-C20-O21

There are no ring outliers.


12 monomers are involved in 28 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	А	201	78M	2	0
2	В	203	78M	1	0
2	А	205	78M	9	0
2	D	203	78M	3	0
2	D	204	78M	1	0
2	А	203	78M	2	0
2	В	201	78M	2	0
2	В	202	78M	2	0
2	А	202	78M	2	0
2	D	202	78M	1	0
2	А	206	78M	3	0
2	А	204	78M	8	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In



addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


























5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# RSRZ {>}2$	$\mathbf{OWAB}(\mathbf{A}^2)$	Q<0.9
1	А	116/130~(89%)	-0.16	4 (3%) 45 35	27, 41, 104, 121	0
1	В	99/130~(76%)	-0.47	1 (1%) 82 77	27, 40, 103, 120	0
1	С	92/130~(70%)	-0.25	5 (5%) 25 17	25, 45, 100, 119	0
1	D	108/130~(83%)	-0.42	0 100 100	28, 42, 80, 106	0
1	Ε	89/130~(68%)	-0.08	5 (5%) 24 16	32, 59, 115, 144	0
1	F	88/130~(67%)	-0.01	4 (4%) 33 23	32, 54, 128, 146	0
All	All	592/780~(75%)	-0.24	19 (3%) 47 37	25, 46, 113, 146	0

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	36	VAL	4.9
1	С	86	TYR	4.9
1	А	89	LEU	4.8
1	F	119	HIS	3.7
1	С	84	SER	3.6

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.




6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
2	78M	А	201	22/22	0.64	0.26	$81,\!106,\!114,\!119$	0
2	78M	С	201	22/22	0.70	0.44	$76,\!92,\!107,\!111$	0
2	78M	D	201	22/22	0.71	0.34	71,94,111,115	0
2	78M	А	203	22/22	0.73	0.40	49,78,89,104	0
2	78M	В	204	22/22	0.75	0.31	$72,\!89,\!99,\!110$	0
2	78M	А	205	22/22	0.77	0.37	$64,\!84,\!98,\!99$	0
2	78M	В	202	22/22	0.79	0.41	$48,\!68,\!95,\!116$	0
2	78M	D	202	22/22	0.81	0.37	$41,\!61,\!85,\!92$	0
2	78M	В	201	22/22	0.82	0.35	62,76,95,106	0
2	78M	D	204	22/22	0.82	0.38	$53,\!73,\!80,\!82$	0
2	78M	С	202	22/22	0.84	0.22	$36,\!69,\!109,\!114$	0
2	78M	В	203	22/22	0.84	0.34	40,64,101,114	0
2	78M	D	203	22/22	0.85	0.26	$61,\!83,\!98,\!104$	0
2	78M	А	206	22/22	0.86	0.55	57,62,79,86	0
2	78M	А	202	22/22	0.88	0.24	$53,\!71,\!92,\!95$	0
2	78M	А	204	22/22	0.88	0.41	42,67,81,97	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









































6.5 Other polymers (i)

There are no such residues in this entry.

