



wwPDB EM Validation Summary Report ⓘ

Nov 4, 2023 – 10:54 AM EDT

PDB ID : 6D6U
EMDB ID : EMD-7817
Title : Human GABA-A receptor alpha1-beta2-gamma2 subtype in complex with GABA and flumazenil, conformation A
Authors : Zhu, S.; Noviello, C.M.; Teng, J.; Walsh Jr, R.M.; Kim, J.J.; Hibbs, R.E.
Deposited on : 2018-04-22
Resolution : 3.92 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	341	
1	C	341	
2	B	358	
2	D	358	
3	E	394	
4	I	213	
4	L	213	
5	J	454	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	K	454	 22% 74%
6	F	3	 67% 33%
6	H	3	 67% 33%
7	G	8	 38% 38% 25%
8	M	5	 40% 60%

2 Entry composition [i](#)

There are 12 unique types of molecules in this entry. The entry contains 17515 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-aminobutyric acid receptor subunit beta-2, Gamma-aminobutyric acid receptor subunit beta-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	333	2726	1788	439	483	16	0	0
1	C	333	2726	1788	439	483	16	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	308	SER	-	linker	UNP P47870
A	309	GLN	-	linker	UNP P47870
A	310	PRO	-	linker	UNP P47870
A	311	ALA	-	linker	UNP P47870
A	312	ARG	-	linker	UNP P47870
A	313	ALA	-	linker	UNP P47870
A	314	ALA	-	linker	UNP P47870
C	308	SER	-	linker	UNP P47870
C	309	GLN	-	linker	UNP P47870
C	310	PRO	-	linker	UNP P47870
C	311	ALA	-	linker	UNP P47870
C	312	ARG	-	linker	UNP P47870
C	313	ALA	-	linker	UNP P47870
C	314	ALA	-	linker	UNP P47870

- Molecule 2 is a protein called Gamma-aminobutyric acid receptor subunit alpha-1, Gamma-aminobutyric acid receptor subunit alpha-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	B	336	2711	1753	455	487	16	0	0
2	D	338	2730	1763	461	490	16	0	0

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	313	SER	-	linker	UNP P14867
B	314	GLN	-	linker	UNP P14867
B	315	PRO	-	linker	UNP P14867
B	316	ALA	-	linker	UNP P14867
B	317	ARG	-	linker	UNP P14867
B	318	ALA	-	linker	UNP P14867
B	319	ALA	-	linker	UNP P14867
D	313	SER	-	linker	UNP P14867
D	314	GLN	-	linker	UNP P14867
D	315	PRO	-	linker	UNP P14867
D	316	ALA	-	linker	UNP P14867
D	317	ARG	-	linker	UNP P14867
D	318	ALA	-	linker	UNP P14867
D	319	ALA	-	linker	UNP P14867

- Molecule 3 is a protein called Gamma-aminobutyric acid receptor subunit gamma-2, Gamma-aminobutyric acid receptor subunit gamma-2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	E	322	2637	1716	436	471	14	0	0

There are 43 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-36	TRP	-	expression tag	UNP P18507
E	-35	SER	-	expression tag	UNP P18507
E	-34	HIS	-	expression tag	UNP P18507
E	-33	PRO	-	expression tag	UNP P18507
E	-32	GLN	-	expression tag	UNP P18507
E	-31	PHE	-	expression tag	UNP P18507
E	-30	GLU	-	expression tag	UNP P18507
E	-29	LYS	-	expression tag	UNP P18507
E	-28	GLY	-	expression tag	UNP P18507
E	-27	GLY	-	expression tag	UNP P18507
E	-26	GLY	-	expression tag	UNP P18507
E	-25	SER	-	expression tag	UNP P18507
E	-24	GLY	-	expression tag	UNP P18507
E	-23	GLY	-	expression tag	UNP P18507
E	-22	GLY	-	expression tag	UNP P18507
E	-21	SER	-	expression tag	UNP P18507

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-20	GLY	-	expression tag	UNP P18507
E	-19	GLY	-	expression tag	UNP P18507
E	-18	SER	-	expression tag	UNP P18507
E	-17	SER	-	expression tag	UNP P18507
E	-16	ALA	-	expression tag	UNP P18507
E	-15	TRP	-	expression tag	UNP P18507
E	-14	SER	-	expression tag	UNP P18507
E	-13	HIS	-	expression tag	UNP P18507
E	-12	PRO	-	expression tag	UNP P18507
E	-11	GLN	-	expression tag	UNP P18507
E	-10	PHE	-	expression tag	UNP P18507
E	-9	GLU	-	expression tag	UNP P18507
E	-8	LYS	-	expression tag	UNP P18507
E	-7	LEU	-	expression tag	UNP P18507
E	-6	GLU	-	expression tag	UNP P18507
E	-5	VAL	-	expression tag	UNP P18507
E	-4	LEU	-	expression tag	UNP P18507
E	-3	PHE	-	expression tag	UNP P18507
E	-2	GLN	-	expression tag	UNP P18507
E	-1	GLY	-	expression tag	UNP P18507
E	0	PRO	-	expression tag	UNP P18507
E	323	SER	-	linker	UNP P18507
E	324	GLN	-	linker	UNP P18507
E	325	PRO	-	linker	UNP P18507
E	326	ALA	-	linker	UNP P18507
E	327	ARG	-	linker	UNP P18507
E	328	ALA	-	linker	UNP P18507

- Molecule 4 is a protein called Kappa Fab Light Chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	I	106	Total	C	N	O	S	0	0
			811	510	132	164	5		
4	L	106	Total	C	N	O	S	0	0
			811	510	132	164	5		

- Molecule 5 is a protein called IgG2b Fab Heavy Chain.


Mol	Chain	Residues	Atoms					AltConf	Trace
5	J	116	Total	C	N	O	S	0	0
			907	574	151	178	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	K	116	907	573	151	179	4	0	0

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
6	F	3	39	22	2	15	0	0
6	H	3	39	22	2	15	0	0

- Molecule 7 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
7	G	8	94	52	2	40	0	0

- Molecule 8 is an oligosaccharide called alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
8	M	5	61	34	2	25	0	0

- Molecule 9 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				AltConf
			Total	C	N	O	
9	A	1	14	8	1	5	0
9	C	1	14	8	1	5	0


- Molecule 10 is GAMMA-AMINO-BUTANOIC ACID (three-letter code: ABU) (formula: $C_4H_9NO_2$).



Continued from previous page...

Mol	Chain	Residues	Atoms			AltConf
			Total	C	O	
11	D	1	21	20	1	0
11	D	1	21	20	1	0
11	E	1	21	20	1	0

- Molecule 12 is ethyl 8-fluoro-5-methyl-6-oxo-5,6-dihydro-4H-imidazo[1,5-a][1,4]benzodiazepine-3-carboxylate (three-letter code: FYP) (formula: C₁₅H₁₄FN₃O₃).




Mol	Chain	Residues	Atoms					AltConf
			Total	C	F	N	O	
12	D	1	22	15	1	3	3	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gamma-aminobutyric acid receptor subunit beta-2, Gamma-aminobutyric acid receptor subunit beta-2



- Molecule 1: Gamma-aminobutyric acid receptor subunit beta-2, Gamma-aminobutyric acid receptor subunit beta-2



- Molecule 2: Gamma-aminobutyric acid receptor subunit alpha-1, Gamma-aminobutyric acid receptor subunit alpha-1



GLU PHE GLU LYS CYS VAL ASN ASN LYS ASP PRO PRO ILE THR ARG THR ILE SER LYS LYS GLY VAL VAL ARG ALA PRO GLN TYR ILE LEU MET PRO PRO ALA GLU GLN LYS SER ARG LYS ASP VAL PHE SER CYS LEU THR ASN THR CYS LEU VAL ARG HIS VAL VAL GLY PHE LEU ASN PRO GLY TYR ASP ILE

SER VAL TRP THR SER ASN GLY HIS THR GLU GLU ASN TYR LYS ASP THR ALA PRO VAL LEU ASP SER GLY SER TYR PHE ILE TYR SER LYS ASN MET PRO THR SER LYS TRP GLU LYS THR ASP PHE SER CYS ASN VAL ARG HIS VAL VAL GLY PHE LEU ASN PRO GLY TYR ASP ILE

LYS THR ILE SER ARG SER PRO GLY LYS

- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:  67% 33%

MAG1
MAG2
BMA3

- Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain H:  67% 33%

MAG1
MAG2
BMA3

- Molecule 7: alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  38% 38% 25%

MAG1
MAG2
BMA3
MAN4
MAN5
MAN6
MAN7
MAN8

- Molecule 8: alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:  40% 60%

MAG1
MAG2
BMA3
MAN4
MAN5

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	292662	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	47	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	4500	Depositor
Magnification	46730	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.253	Depositor
Minimum map value	-0.175	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.006	Depositor
Recommended contour level	0.024	Depositor
Map size (Å)	278.2, 278.2, 278.2	wwPDB
Map dimensions	260, 260, 260	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.07, 1.07, 1.07	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FYP, Y01, MAN, NAG, BMA, ABU

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/2798	0.44	0/3810
1	C	0.31	0/2798	0.45	0/3810
2	B	0.31	0/2780	0.46	0/3780
2	D	0.34	0/2799	0.47	0/3805
3	E	0.32	0/2707	0.46	0/3686
4	I	0.32	0/829	0.46	0/1123
4	L	0.32	0/829	0.45	0/1123
5	J	0.28	0/928	0.45	0/1260
5	K	0.29	0/928	0.45	0/1260
All	All	0.32	0/17396	0.46	0/23657

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2726	0	2736	48	0
1	C	2726	0	2736	48	0
2	B	2711	0	2703	53	0
2	D	2730	0	2722	57	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	2637	0	2616	58	0
4	I	811	0	784	5	0
4	L	811	0	784	4	0
5	J	907	0	877	14	0
5	K	907	0	875	9	0
6	F	39	0	34	1	0
6	H	39	0	34	2	0
7	G	94	0	79	2	0
8	M	61	0	52	1	0
9	A	14	0	13	1	0
9	C	14	0	13	0	0
10	A	7	0	0	0	0
10	C	7	0	0	0	0
11	A	21	0	28	0	0
11	B	42	0	56	6	0
11	C	84	0	112	7	0
11	D	84	0	112	9	0
11	E	21	0	28	3	0
12	D	22	0	0	0	0
All	All	17515	0	17394	275	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 275 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:294:TYR:OH	11:B:410:Y01:HBG	1.37	1.23
1:A:99:LEU:HD13	1:A:155:GLU:OE2	1.77	0.83
2:B:294:TYR:OH	11:B:410:Y01:CBG	2.25	0.82
3:E:125:THR:HG22	3:E:126:THR:HG23	1.62	0.81
2:D:88:ASN:ND2	2:D:117:LYS:O	2.13	0.81

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/341 (97%)	318 (96%)	13 (4%)	0	100	100
1	C	331/341 (97%)	320 (97%)	11 (3%)	0	100	100
2	B	334/358 (93%)	319 (96%)	15 (4%)	0	100	100
2	D	336/358 (94%)	313 (93%)	23 (7%)	0	100	100
3	E	316/394 (80%)	296 (94%)	20 (6%)	0	100	100
4	I	104/213 (49%)	97 (93%)	7 (7%)	0	100	100
4	L	104/213 (49%)	98 (94%)	6 (6%)	0	100	100
5	J	114/454 (25%)	109 (96%)	5 (4%)	0	100	100
5	K	114/454 (25%)	108 (95%)	6 (5%)	0	100	100
All	All	2084/3126 (67%)	1978 (95%)	106 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	301/309 (97%)	299 (99%)	2 (1%)	84	90
1	C	301/309 (97%)	299 (99%)	2 (1%)	84	90
2	B	298/319 (93%)	291 (98%)	7 (2%)	50	70
2	D	300/319 (94%)	295 (98%)	5 (2%)	60	78
3	E	295/355 (83%)	290 (98%)	5 (2%)	60	78

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	I	90/188 (48%)	89 (99%)	1 (1%)	73	84
4	L	90/188 (48%)	90 (100%)	0	100	100
5	J	97/407 (24%)	96 (99%)	1 (1%)	76	86
5	K	97/407 (24%)	97 (100%)	0	100	100
All	All	1869/2801 (67%)	1846 (99%)	23 (1%)	72	83

5 of 23 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	275	ASN
3	E	115	ASN
3	E	58	TYR
3	E	123	TRP
2	B	275	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 28 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	275	ASN
5	J	110	GLN
3	E	66	ASN
4	I	38	GLN
2	D	337	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

19 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	F	1	6,1	14,14,15	0.23	0	17,19,21	0.45	0
6	NAG	F	2	6	14,14,15	0.18	0	17,19,21	0.50	0
6	BMA	F	3	6	11,11,12	0.48	0	15,15,17	0.76	0
7	NAG	G	1	2,7	14,14,15	0.25	0	17,19,21	0.60	0
7	NAG	G	2	7	14,14,15	0.24	0	17,19,21	0.40	0
7	BMA	G	3	7	11,11,12	0.26	0	15,15,17	0.94	1 (6%)
7	MAN	G	4	7	11,11,12	0.83	0	15,15,17	1.47	2 (13%)
7	MAN	G	5	7	11,11,12	0.73	0	15,15,17	1.02	0
7	MAN	G	6	7	11,11,12	0.45	0	15,15,17	1.24	1 (6%)
7	MAN	G	7	7	11,11,12	0.65	0	15,15,17	1.31	2 (13%)
7	MAN	G	8	7	11,11,12	0.29	0	15,15,17	1.31	2 (13%)
6	NAG	H	1	6,1	14,14,15	0.30	0	17,19,21	0.64	0
6	NAG	H	2	6	14,14,15	0.42	0	17,19,21	1.26	1 (5%)
6	BMA	H	3	6	11,11,12	1.00	0	15,15,17	1.12	2 (13%)
8	NAG	M	1	2,8	14,14,15	0.43	0	17,19,21	0.46	0
8	NAG	M	2	8	14,14,15	0.22	0	17,19,21	0.47	0
8	BMA	M	3	8	11,11,12	0.52	0	15,15,17	0.85	0
8	MAN	M	4	8	11,11,12	0.71	0	15,15,17	0.88	1 (6%)
8	MAN	M	5	8	11,11,12	0.70	1 (9%)	15,15,17	1.13	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	F	1	6,1	-	3/6/23/26	0/1/1/1
6	NAG	F	2	6	-	2/6/23/26	0/1/1/1
6	BMA	F	3	6	-	1/2/19/22	0/1/1/1
7	NAG	G	1	2,7	-	2/6/23/26	0/1/1/1
7	NAG	G	2	7	-	2/6/23/26	0/1/1/1
7	BMA	G	3	7	-	2/2/19/22	0/1/1/1
7	MAN	G	4	7	-	2/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	MAN	G	5	7	-	0/2/19/22	0/1/1/1
7	MAN	G	6	7	-	0/2/19/22	0/1/1/1
7	MAN	G	7	7	-	0/2/19/22	0/1/1/1
7	MAN	G	8	7	-	2/2/19/22	0/1/1/1
6	NAG	H	1	6,1	-	3/6/23/26	0/1/1/1
6	NAG	H	2	6	-	5/6/23/26	0/1/1/1
6	BMA	H	3	6	-	1/2/19/22	0/1/1/1
8	NAG	M	1	2,8	-	2/6/23/26	0/1/1/1
8	NAG	M	2	8	-	2/6/23/26	0/1/1/1
8	BMA	M	3	8	-	2/2/19/22	0/1/1/1
8	MAN	M	4	8	-	2/2/19/22	0/1/1/1
8	MAN	M	5	8	-	1/2/19/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	M	5	MAN	C1-C2	2.11	1.57	1.52

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	G	4	MAN	C1-O5-C5	4.38	118.12	112.19
6	H	2	NAG	C2-N2-C7	4.28	129.00	122.90
7	G	7	MAN	C1-O5-C5	3.57	117.03	112.19
7	G	4	MAN	O2-C2-C3	-3.01	104.11	110.14
7	G	6	MAN	C1-O5-C5	-2.83	108.36	112.19

There are no chirality outliers.

5 of 34 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
6	F	2	NAG	O5-C5-C6-O6
7	G	1	NAG	O5-C5-C6-O6
8	M	1	NAG	O5-C5-C6-O6
6	F	2	NAG	C4-C5-C6-O6
8	M	3	BMA	C4-C5-C6-O6


There are no ring outliers.


6 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	H	2	NAG	1	0
6	H	1	NAG	1	0
7	G	8	MAN	2	0
7	G	6	MAN	1	0
8	M	1	NAG	1	0
6	F	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.









5.6 Ligand geometry [i](#)

17 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
10	ABU	C	405	-	6,6,6	0.85	0	6,6,6	1.20	0
11	Y01	B	410	-	24,24,38	0.82	1 (4%)	37,39,57	1.71	9 (24%)
11	Y01	C	407	-	24,24,38	0.83	1 (4%)	37,39,57	1.68	9 (24%)
11	Y01	D	406	-	24,24,38	0.82	1 (4%)	37,39,57	1.71	9 (24%)
11	Y01	D	408	-	24,24,38	0.82	1 (4%)	37,39,57	1.70	9 (24%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
11	Y01	D	409	-	24,24,38	0.82	1 (4%)	37,39,57	1.71	9 (24%)
11	Y01	A	406	-	24,24,38	0.82	1 (4%)	37,39,57	1.66	9 (24%)
12	FYP	D	410	-	22,24,24	1.23	2 (9%)	26,35,35	1.27	2 (7%)
11	Y01	C	409	-	24,24,38	0.82	1 (4%)	37,39,57	1.71	9 (24%)
11	Y01	D	407	-	24,24,38	0.82	1 (4%)	37,39,57	1.69	8 (21%)
11	Y01	E	401	-	24,24,38	0.81	1 (4%)	37,39,57	1.70	9 (24%)
9	NAG	A	404	1	14,14,15	0.36	0	17,19,21	0.50	0
11	Y01	B	409	-	24,24,38	0.81	1 (4%)	37,39,57	1.69	9 (24%)
10	ABU	A	405	-	6,6,6	0.83	0	6,6,6	1.14	0
9	NAG	C	404	1	14,14,15	0.18	0	17,19,21	0.40	0
11	Y01	C	408	-	24,24,38	0.81	1 (4%)	37,39,57	1.69	9 (24%)
11	Y01	C	406	-	24,24,38	0.82	1 (4%)	37,39,57	1.70	9 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
10	ABU	C	405	-	-	2/4/4/4	-
11	Y01	B	410	-	-	-	0/4/4/4
11	Y01	C	407	-	-	-	0/4/4/4
11	Y01	D	406	-	-	-	0/4/4/4
11	Y01	D	408	-	-	-	0/4/4/4
12	FYP	D	410	-	-	0/7/23/23	0/2/3/3
11	Y01	A	406	-	-	-	0/4/4/4
11	Y01	D	409	-	-	-	0/4/4/4
11	Y01	C	409	-	-	-	0/4/4/4
11	Y01	D	407	-	-	-	0/4/4/4
11	Y01	E	401	-	-	-	0/4/4/4
9	NAG	A	404	1	-	2/6/23/26	0/1/1/1
11	Y01	B	409	-	-	-	0/4/4/4
10	ABU	A	405	-	-	2/4/4/4	-
9	NAG	C	404	1	-	2/6/23/26	0/1/1/1
11	Y01	C	408	-	-	-	0/4/4/4
11	Y01	C	406	-	-	-	0/4/4/4

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
12	D	410	FYP	C15-N16	-3.07	1.29	1.35
12	D	410	FYP	C08-C07	-2.89	1.47	1.51
11	C	406	Y01	CBH-CBF	-2.25	1.52	1.56
11	D	406	Y01	CBH-CBF	-2.23	1.52	1.56
11	C	407	Y01	CBH-CBF	-2.23	1.52	1.56

The worst 5 of 109 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	D	409	Y01	CBI-CBG-CBD	-4.82	107.23	114.38
11	E	401	Y01	CBI-CBG-CBD	-4.80	107.27	114.38
11	B	410	Y01	CBI-CBG-CBD	-4.79	107.28	114.38
11	D	406	Y01	CBI-CBG-CBD	-4.76	107.33	114.38
11	C	409	Y01	CBI-CBG-CBD	-4.76	107.33	114.38

There are no chirality outliers.

5 of 8 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
9	C	404	NAG	C4-C5-C6-O6
9	C	404	NAG	O5-C5-C6-O6
9	A	404	NAG	O5-C5-C6-O6
9	A	404	NAG	C4-C5-C6-O6
10	C	405	ABU	OXT-C-CG-CB


There are no ring outliers.



10 monomers are involved in 26 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	B	410	Y01	6	0
11	C	407	Y01	1	0
11	D	408	Y01	1	0
11	D	409	Y01	6	0
11	C	409	Y01	1	0
11	D	407	Y01	2	0
11	E	401	Y01	3	0
9	A	404	NAG	1	0
11	C	408	Y01	4	0
11	C	406	Y01	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In



addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-7817. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 130




Y Index: 130




Z Index: 130


6.2.2 Raw map



X Index: 130



Y Index: 130




Z Index: 130


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 135




Y Index: 145




Z Index: 138


6.3.2 Raw map



X Index: 135



Y Index: 145




Z Index: 138


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.024. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 156 nm³; this corresponds to an approximate mass of 141 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [\(i\)](#)




*Reported resolution corresponds to spatial frequency of 0.255 \AA^{-1}

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.255 \AA^{-1}

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.92	-	-
Author-provided FSC curve	-	-	-
Unmasked-calculated*	4.11	4.45	4.14

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-7817 and PDB model 6D6U. Per-residue inclusion information can be found in section 3 on page 11.

9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.024 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.024).





























9.4 Atom inclusion [i](#)



At the recommended contour level, 90% of all backbone atoms, 82% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (0.024) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8240	 0.4610
A	 0.7990	 0.4420
B	 0.8030	 0.4500
C	 0.8210	 0.4610
D	 0.8480	 0.4750
E	 0.8010	 0.4530
F	 0.8210	 0.4780
G	 0.8940	 0.4740
H	 0.8720	 0.4710
I	 0.8610	 0.4740
J	 0.8620	 0.4780
K	 0.8680	 0.4810
L	 0.8520	 0.4770
M	 0.7380	 0.4580

