

wwPDB X-ray Structure Validation Summary Report (i)

Jan 30, 2021 – 02:44 PM EST

PDB ID : 3DJC

Title: CRYSTAL STRUCTURE OF PANTOTHENATE KINASE FROM LE-

GIONELLA PNEUMOPHILA

Authors: Patskovsky, Y.; Bonanno, J.B.; Romero, R.; Dickey, M.; Logan, C.; Wasser-

man, S.; Maletic, M.; Koss, J.; Sauder, J.M.; Burley, S.K.; Almo, S.C.; New

York SGX Research Center for Structural Genomics (NYSGXRC)

Deposited on : 2008-06-23

Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.16

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.16

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	266	75%	18%		-
1	В	266	78%	17%		
1	С	266	78%	17%		
1	D	266	82%	11%	• 5%	%
1	Е	266	80%	12%	• 5%	_



Continued from previous page...

Mol	Chain	Length	Quality of chain	
1	F	266	77%	14% • 6%
1	G	266	12%	11% • •
1	Н	266	13%	9% • •
1	I	266	7% 83%	12% • •
1	J	266	8%	11% • 6%
1	K	266	78%	14% • 6%
1	L	266	15% 82%	11% 6%



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 23826 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Type III pantothenate kinase.

Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	Trace
1	A	255	Total	С	N	О	S	0	5	0
1	A	255	1967	1256	336	367	8	0	9	
1	В	256	Total	С	N	О	S	0	7	0
1	Б	250	1993	1271	346	368	8	0	1	
1	С	256	Total	С	N	О	S	0	5	0
1		250	1973	1259	339	367	8	0		
1	D	254	Total	С	N	О	S	0	6	0
1	D	204	1962	1251	338	365	8	0	0	
1	Е	252	Total	С	N	О	S	0	4	0
1	15	202	1938	1236	333	361	8	0	4	
1	F	251	Total	С	N	О	S	0	4	0
1	Г	231	1931	1233	331	359	8	U	4	
1	G	256	Total	С	N	О	S	0	2	0
1	G	250	1960	1250	337	365	8	0	2	
1	Н	255	Total	С	N	О	S	0	4	0
1	11	255	1967	1255	339	365	8	0	4	
1	I	255	Total	С	N	О	S	0	6	0
1	1	255	1971	1258	338	367	8	0	0	
1	J	250	Total	С	N	О	S	0	5	0
1	J	250	1941	1238	335	360	8	0	9	
1	K	250	Total	С	N	О	S	0	6	0
1	IV.	200	1941	1237	336	360	8	0	U	
1	L	250	Total	С	N	О	S	0	4	0
1	L	200	1931	1234	331	358	8	U	4	

There are 132 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP Q5ZX22
A	0	SER	-	expression tag	UNP Q5ZX22
A	1	LEU	-	expression tag	UNP Q5ZX22
A	257	GLU	-	expression tag	UNP Q5ZX22
A	258	GLY	-	expression tag	UNP Q5ZX22



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
A	259	HIS	-	expression tag	UNP Q5ZX22
A	260	HIS	-	expression tag	UNP Q5ZX22
A	261	HIS	-	expression tag	UNP Q5ZX22
A	262	HIS	-	expression tag	UNP Q5ZX22
A	263	HIS	-	expression tag	UNP Q5ZX22
A	264	HIS	-	expression tag	UNP Q5ZX22
В	-1	MET	-	expression tag	UNP Q5ZX22
В	0	SER	-	expression tag	UNP Q5ZX22
В	1	LEU	-	expression tag	UNP Q5ZX22
В	257	GLU	-	expression tag	UNP Q5ZX22
В	258	GLY	-	expression tag	UNP Q5ZX22
В	259	HIS	-	expression tag	UNP Q5ZX22
В	260	HIS	-	expression tag	UNP Q5ZX22
В	261	HIS	-	expression tag	UNP Q5ZX22
В	262	HIS	-	expression tag	UNP Q5ZX22
В	263	HIS	-	expression tag	UNP Q5ZX22
В	264	HIS	-	expression tag	UNP Q5ZX22
С	-1	MET	-	expression tag	UNP Q5ZX22
С	0	SER	-	expression tag	UNP Q5ZX22
С	1	LEU	-	expression tag	UNP Q5ZX22
С	257	GLU	-	expression tag	UNP Q5ZX22
С	258	GLY	-	expression tag	UNP Q5ZX22
С	259	HIS	-	expression tag	UNP Q5ZX22
С	260	HIS	-	expression tag	UNP Q5ZX22
С	261	HIS	-	expression tag	UNP Q5ZX22
С	262	HIS	-	expression tag	UNP Q5ZX22
С	263	HIS	-	expression tag	UNP Q5ZX22
С	264	HIS	_	expression tag	UNP Q5ZX22
D	-1	MET	-	expression tag	UNP Q5ZX22
D	0	SER	-	expression tag	UNP Q5ZX22
D	1	LEU	-	expression tag	UNP Q5ZX22
D	257	GLU	-	expression tag	UNP Q5ZX22
D	258	GLY	-	expression tag	UNP Q5ZX22
D	259	HIS	-	expression tag	UNP Q5ZX22
D	260	HIS	-	expression tag	UNP Q5ZX22
D	261	HIS	_	expression tag	UNP Q5ZX22
D	262	HIS	-	expression tag	UNP Q5ZX22
D	263	HIS	-	expression tag	UNP Q5ZX22
D	264	HIS	_	expression tag	UNP Q5ZX22
Е	-1	MET	-	expression tag	UNP Q5ZX22
Е	0	SER	-	expression tag	UNP Q5ZX22
Е	1	LEU	-	expression tag	UNP Q5ZX22



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
E	257	GLU	_	expression tag	UNP Q5ZX22
Е	258	GLY	-	expression tag	UNP Q5ZX22
Е	259	HIS	-	expression tag	UNP Q5ZX22
Е	260	HIS	-	expression tag	UNP Q5ZX22
Е	261	HIS	-	expression tag	UNP Q5ZX22
Е	262	HIS	-	expression tag	UNP Q5ZX22
Е	263	HIS	-	expression tag	UNP Q5ZX22
Е	264	HIS	-	expression tag	UNP Q5ZX22
F	-1	MET	-	expression tag	UNP Q5ZX22
F	0	SER	-	expression tag	UNP Q5ZX22
F	1	LEU	-	expression tag	UNP Q5ZX22
F	257	GLU	-	expression tag	UNP Q5ZX22
F	258	GLY	-	expression tag	UNP Q5ZX22
F	259	HIS	-	expression tag	UNP Q5ZX22
F	260	HIS	-	expression tag	UNP Q5ZX22
F	261	HIS	-	expression tag	UNP Q5ZX22
F	262	HIS	-	expression tag	UNP Q5ZX22
F	263	HIS	-	expression tag	UNP Q5ZX22
F	264	HIS	-	expression tag	UNP Q5ZX22
G	-1	MET	-	expression tag	UNP Q5ZX22
G	0	SER	-	expression tag	UNP Q5ZX22
G	1	LEU	-	expression tag	UNP Q5ZX22
G	257	GLU	-	expression tag	UNP Q5ZX22
G	258	GLY	-	expression tag	UNP Q5ZX22
G	259	HIS	-	expression tag	UNP Q5ZX22
G	260	HIS	-	expression tag	UNP Q5ZX22
G	261	HIS	-	expression tag	UNP Q5ZX22
G	262	HIS	-	expression tag	UNP Q5ZX22
G	263	HIS	-	expression tag	UNP Q5ZX22
G	264	HIS	-	expression tag	UNP Q5ZX22
Н	-1	MET	_	expression tag	UNP Q5ZX22
Н	0	SER	-	expression tag	UNP Q5ZX22
Н	1	LEU	_	expression tag	UNP Q5ZX22
Н	257	GLU	-	expression tag	UNP Q5ZX22
Н	258	GLY	-	expression tag	UNP Q5ZX22
Н	259	HIS	-	expression tag	UNP Q5ZX22
Н	260	HIS	_	expression tag	UNP Q5ZX22
Н	261	HIS	-	expression tag	UNP Q5ZX22
Н	262	HIS	_	expression tag	UNP Q5ZX22
Н	263	HIS	-	expression tag	UNP Q5ZX22
Н	264	HIS	-	expression tag	UNP Q5ZX22
I	-1	MET	_	expression tag	UNP Q5ZX22



 $Continued\ from\ previous\ page...$


Chain	Residue	Modelled Modelled	Actual	Comment	Reference
I	0	SER	-	expression tag	UNP Q5ZX22
I	1	LEU	-	expression tag	UNP Q5ZX22
I	257	GLU	-	expression tag	UNP Q5ZX22
I	258	GLY	-	expression tag	UNP Q5ZX22
I	259	HIS	-	expression tag	UNP Q5ZX22
I	260	HIS	-	expression tag	UNP Q5ZX22
I	261	HIS	-	expression tag	UNP Q5ZX22
I	262	HIS	-	expression tag	UNP Q5ZX22
I	263	HIS	-	expression tag	UNP Q5ZX22
I	264	HIS	-	expression tag	UNP Q5ZX22
J	-1	MET	-	expression tag	UNP Q5ZX22
J	0	SER	-	expression tag	UNP Q5ZX22
J	1	LEU	_	expression tag	UNP Q5ZX22
J	257	GLU	-	expression tag	UNP Q5ZX22
J	258	GLY	-	expression tag	UNP Q5ZX22
J	259	HIS	-	expression tag	UNP Q5ZX22
J	260	HIS	-	expression tag	UNP Q5ZX22
J	261	HIS	-	expression tag	UNP Q5ZX22
J	262	HIS	-	expression tag	UNP Q5ZX22
J	263	HIS	-	expression tag	UNP Q5ZX22
J	264	HIS	-	expression tag	UNP Q5ZX22
K	-1	MET	-	expression tag	UNP Q5ZX22
K	0	SER	-	expression tag	UNP Q5ZX22
K	1	LEU	-	expression tag	UNP Q5ZX22
K	257	GLU	-	expression tag	UNP Q5ZX22
K	258	GLY	-	expression tag	UNP Q5ZX22
K	259	HIS	-	expression tag	UNP Q5ZX22
K	260	HIS	-	expression tag	UNP Q5ZX22
K	261	HIS	-	expression tag	UNP Q5ZX22
K	262	HIS	-	expression tag	UNP Q5ZX22
K	263	HIS	-	expression tag	UNP Q5ZX22
K	264	HIS	-	expression tag	UNP Q5ZX22
L	-1	MET	_	expression tag	UNP Q5ZX22
L	0	SER	-	expression tag	UNP Q5ZX22
L	1	LEU	-	expression tag	UNP Q5ZX22
L	257	GLU	_	expression tag	UNP Q5ZX22
L	258	GLY	_	expression tag	UNP Q5ZX22
L	259	HIS	-	expression tag	UNP Q5ZX22
L	260	HIS	-	expression tag	UNP Q5ZX22
L	261	HIS		expression tag	UNP Q5ZX22
L	262	HIS	-	expression tag	UNP Q5ZX22
L	263	HIS	-	expression tag	UNP Q5ZX22



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
L	264	HIS	-	expression tag	UNP Q5ZX22

 \bullet Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $\mathrm{C_3H_8O_3}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	В	1	Total C O 6 3 3	0	0
2	С	1	Total C O 6 3 3	0	0
2	С	1	Total C O 6 3 3	0	0
2	Е	1	Total C O 6 3 3	0	0
2	Е	1	Total C O 6 3 3	0	0
2	Н	1	Total C O 6 3 3	0	0
2	Н	1	Total C O 6 3 3	0	0
2	J	1	Total C O 6 3 3	0	0
2	L	1	Total C O 6 3 3	0	0



• Molecule 3 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	39	Total O 39 39	0	0
3	В	37	Total O 37 37	0	0
3	С	33	Total O 33 33	0	0
3	D	28	Total O 28 28	0	0
3	E	30	Total O 30 30	0	0
3	F	27	Total O 27 27	0	0
3	G	18	Total O 18 18	0	0
3	Н	10	Total O 10 10	0	0
3	I	27	Total O 27 27	0	0
3	J	12	Total O 12 12	0	0
3	К	15	Total O 15 15	0	0
3	L	9	Total O 9 9	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Type III pantothenate kinase









• Molecule 1: Type III pantothenate kinase




• Molecule 1: Type III pantothenate kinase






• Molecule 1: Type III pantothenate kinase







• Molecule 1: Type III pantothenate kinase




• Molecule 1: Type III pantothenate kinase



 \bullet Molecule 1: Type III pantothenate kinase



• Molecule 1: Type III pantothenate kinase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	95.39Å 134.57Å 134.40Å	Depositor
a, b, c, α , β , γ	90.00° 93.24° 90.00°	Depositor
Resolution (Å)	20.00 - 2.40	Depositor
Resolution (A)	32.01 - 2.35	EDS
% Data completeness	97.5 (20.00-2.40)	Depositor
(in resolution range)	88.4 (32.01-2.35)	EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$< I/\sigma(I) > 1$	1.38 (at 2.36Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
D D.	0.259 , 0.313	Depositor
R, R_{free}	0.260 , 0.312	DCC
R_{free} test set	4086 reflections (3.02%)	wwPDB-VP
Wilson B-factor (Å ²)	60.4	Xtriage
Anisotropy	0.031	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.31, 38.3	EDS
L-test for twinning ²	$ < L >=0.42, < L^2>=0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	23826	wwPDB-VP
Average B, all atoms $(Å^2)$	63.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.68% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Во	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.41	0/2015	0.61	$1/2728 \ (0.0\%)$
1	В	0.43	0/2047	0.67	0/2768
1	С	0.44	0/2021	0.62	0/2736
1	D	0.43	0/2011	0.63	0/2719
1	Е	0.41	0/1982	0.61	0/2682
1	F	0.40	0/1974	0.61	0/2670
1	G	0.45	0/1999	0.59	0/2707
1	Н	0.39	0/2012	0.57	0/2723
1	I	0.39	0/2022	0.59	0/2737
1	J	0.38	0/1987	0.58	0/2687
1	K	0.39	0/1989	0.59	0/2690
1	L	0.39	0/1974	0.56	0/2668
All	All	0.41	0/24033	0.60	$1/32515 \ (0.0\%)$

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	D	0	1
1	G	0	2
1	Н	0	1
All	All	0	5

There are no bond length outliers.

All (1) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}(^{o})$
1	A	154	LEU	CA-CB-CG	5.36	127.62	115.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	62	CYS	Peptide
1	D	0	SER	Peptide
1	G	231	LYS	Peptide
1	G	233	GLY	Peptide
1	Н	32	VAL	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1967	0	2026	27	0
1	В	1993	0	2063	17	0
1	С	1973	0	2034	19	0
1	D	1962	0	2024	14	0
1	Е	1938	0	1987	16	0
1	F	1931	0	1988	22	0
1	G	1960	0	2013	14	0
1	Н	1967	0	2025	8	0
1	I	1971	0	2034	14	0
1	J	1941	0	1996	11	0
1	K	1941	0	1996	15	0
1	L	1931	0	1990	10	0
2	A	12	0	16	0	0
2	В	6	0	8	0	0
2	С	12	0	16	0	0
2	Ε	12	0	16	0	0
2	Н	12	0	16	0	0
2	J	6	0	8	0	0
2	L	6	0	8	0	0
3	A	39	0	0	1	0
3	В	37	0	0	1	0
3	С	33	0	0	0	0



Continued	trom	mmoninonic	maaa
COHABABACA		DIEUIUU	DUIUE
0 0 1000100000			

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	28	0	0	0	0
3	Ε	30	0	0	0	0
3	F	27	0	0	0	0
3	G	18	0	0	0	0
3	Н	10	0	0	0	0
3	I	27	0	0	0	0
3	J	12	0	0	0	0
3	K	15	0	0	0	0
3	L	9	0	0	0	0
All	All	23826	0	24264	160	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 160 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:F:174:ILE:HG13	1:F:175:LYS:N	1.94	0.79
1:D:1:LEU:HB3	1:D:56:ILE:HD13	1.67	0.75
1:C:128:ILE:HD11	1:C:219:ILE:HD11	1.67	0.75
1:C:34[A]:THR:HG21	1:F:27:ARG:HB3	1.71	0.73
1:E:206:ARG:HB3	1:F:179:VAL:HG21	1.71	0.73

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	258/266~(97%)	249 (96%)	9 (4%)	0	100	100
1	В	261/266 (98%)	252 (97%)	9 (3%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	\mathbf{C}	259/266~(97%)	248 (96%)	9 (4%)	2 (1%)	19	29
1	D	256/266~(96%)	250 (98%)	5 (2%)	1 (0%)	34	48
1	E	252/266~(95%)	245 (97%)	7 (3%)	0	100	100
1	F	251/266 (94%)	238 (95%)	12 (5%)	1 (0%)	34	48
1	G	256/266~(96%)	244 (95%)	10 (4%)	2 (1%)	19	29
1	Н	257/266 (97%)	246 (96%)	11 (4%)	0	100	100
1	Ι	259/266 (97%)	244 (94%)	15 (6%)	0	100	100
1	J	251/266 (94%)	243 (97%)	7 (3%)	1 (0%)	34	48
1	K	250/266 (94%)	244 (98%)	4 (2%)	2 (1%)	19	29
1	L	250/266 (94%)	240 (96%)	9 (4%)	1 (0%)	34	48
All	All	3060/3192 (96%)	2943 (96%)	107 (4%)	10 (0%)	41	55

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	233	GLY
1	J	213	ASN
1	F	213	ASN
1	K	213	ASN
1	L	213	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Analysed Rotameric Outliers		Percentiles
1	A	$218/222\ (98\%)$	192 (88%)	26 (12%)	5 6
1	В	221/222~(100%)	193 (87%)	28 (13%)	4 5
1	С	218/222 (98%)	195 (89%)	23 (11%)	6 9
1	D	$217/222\ (98\%)$	194 (89%)	23 (11%)	6 9
1	E	$214/222\ (96\%)$	192 (90%)	22 (10%)	7 10
1	F	$213/222\ (96\%)$	188 (88%)	25 (12%)	5 7



Continued	trom	mmoninonic	maaa
COHABABACA		DIEUIUU	DUIUE
0 0 1000100000			

Mol	Chain	Analysed	Rotameric	Outliers	Percer	ntiles
1	G	216/222 (97%)	198 (92%)	18 (8%)	11	17
1	Н	217/222 (98%)	200 (92%)	17 (8%)	12	19
1	I	219/222 (99%)	201 (92%)	18 (8%)	11	17
1	J	214/222 (96%)	197 (92%)	17 (8%)	12	19
1	K	214/222 (96%)	191 (89%)	23 (11%)	6	9
1	L	213/222 (96%)	198 (93%)	15 (7%)	15	24
All	All	2594/2664 (97%)	2339 (90%)	255 (10%)	8	11

5 of 255 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	Е	109	ARG
1	F	203	LEU
1	K	203	LEU
1	Е	179	VAL
1	F	48	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 45 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	232	GLN
1	G	232	GLN
1	K	245	GLN
1	G	112	ASN
1	G	245	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	В	ond leng	gths	В	ond ang	gles
IVIOI	Type	Chain	nes	S LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z >2
2	GOL	Н	266	-	5,5,5	0.35	0	5,5,5	0.28	0
2	GOL	L	265	-	5,5,5	0.37	0	5,5,5	0.33	0
2	GOL	J	265	-	5,5,5	0.39	0	5,5,5	0.30	0
2	GOL	Н	265	-	5,5,5	0.38	0	5,5,5	0.24	0
2	GOL	В	265	-	5,5,5	0.37	0	5,5,5	0.31	0
2	GOL	Е	265	-	5,5,5	0.35	0	5,5,5	0.27	0
2	GOL	С	265	-	5,5,5	0.37	0	5,5,5	0.25	0
2	GOL	Е	266	-	5,5,5	0.32	0	5,5,5	0.43	0
2	GOL	A	265	-	5,5,5	0.40	0	5,5,5	0.19	0
2	GOL	A	266	-	5,5,5	0.34	0	5,5,5	0.34	0
2	GOL	С	266	-	5,5,5	0.37	0	5,5,5	0.36	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	Н	266	-	-	2/4/4/4	-
2	GOL	L	265	-	-	2/4/4/4	-
2	GOL	J	265	-	=	0/4/4/4	-
2	GOL	Н	265	-	-	2/4/4/4	-
2	GOL	В	265	-	-	2/4/4/4	-
2	GOL	Е	265	-	-	2/4/4/4	-
2	GOL	С	265	-	-	4/4/4/4	-
2	GOL	E	266	-	-	4/4/4/4	-
2	GOL	A	265	-	-	2/4/4/4	-
2	GOL	A	266	-	-	4/4/4/4	-
2	GOL	С	266	-	-	4/4/4/4	-



There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 28 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	В	265	GOL	O1-C1-C2-C3
2	С	265	GOL	C1-C2-C3-O3
2	С	265	GOL	O2-C2-C3-O3
2	Е	266	GOL	O1-C1-C2-C3
2	Е	266	GOL	C1-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\#\mathrm{RSRZ}{>}2$	$OWAB(\AA^2)$	Q < 0.9
1	A	255/266~(95%)	0.24	11 (4%) 35 33	28, 54, 79, 87	0
1	В	256/266 (96%)	0.18	5 (1%) 65 63	29, 52, 74, 82	0
1	С	256/266~(96%)	0.27	10 (3%) 39 38	34, 55, 73, 94	0
1	D	254/266~(95%)	0.24	9 (3%) 44 43	38, 54, 78, 111	0
1	E	252/266 (94%)	0.36	11 (4%) 34 33	32, 52, 78, 102	0
1	F	251/266 (94%)	0.36	14 (5%) 24 23	39, 58, 83, 100	0
1	G	256/266 (96%)	0.77	31 (12%) 4 3	45, 69, 92, 135	0
1	Н	255/266 (95%)	0.74	35 (13%) 3 2	44, 68, 96, 113	0
1	I	255/266~(95%)	0.47	19 (7%) 14 13	46, 65, 90, 104	0
1	J	250/266 (93%)	0.56	20 (8%) 12 11	47, 67, 90, 99	0
1	K	250/266 (93%)	0.43	21 (8%) 11 10	39, 63, 99, 115	0
1	L	250/266 (93%)	0.97	39 (15%) 2 1	50, 75, 102, 120	0
All	All	3040/3192 (95%)	0.46	225 (7%) 14 13	28, 61, 90, 135	0

The worst 5 of 225 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	167	LYS	14.0
1	L	94	GLY	12.2
1	I	167	LYS	10.1
1	G	169	PRO	8.4
1	G	166	ALA	8.4

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	GOL	С	265	6/6	0.72	0.19	47,55,76,79	0
2	GOL	J	265	6/6	0.77	0.26	61,84,87,91	0
2	GOL	A	266	6/6	0.82	0.16	60,68,70,71	0
2	GOL	С	266	6/6	0.87	0.14	51,62,70,74	0
2	GOL	Н	266	6/6	0.88	0.26	69,76,85,96	0
2	GOL	L	265	6/6	0.89	0.18	72,83,91,96	0
2	GOL	Е	265	6/6	0.94	0.13	39,43,46,55	0
2	GOL	Н	265	6/6	0.94	0.17	76,84,104,108	0
2	GOL	Е	266	6/6	0.95	0.11	46,67,71,86	0
2	GOL	A	265	6/6	0.95	0.19	57,74,86,98	0
2	GOL	В	265	6/6	0.97	0.11	34,58,64,66	0

6.5 Other polymers (i)

There are no such residues in this entry.

