



wwPDB X-ray Structure Validation Summary Report ⓘ

Nov 20, 2023 – 10:22 PM JST

PDB ID : 7DTG
Title : Crystal structure of lamin B1 Ig-like domain from human
Authors : Ahn, J.; Lee, J.; Ha, N.-C.
Deposited on : 2021-01-05
Resolution : 3.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1257 (3.70-3.50)
Clashscore	141614	1353 (3.70-3.50)
Ramachandran outliers	138981	1307 (3.70-3.50)
Sidechain outliers	138945	1307 (3.70-3.50)
RSRZ outliers	127900	1161 (3.70-3.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	149	74% 5% 21%
1	B	149	71% 8% 21%
1	C	149	72% 7% 21%
1	D	149	67% 12% 21%
1	E	149	68% 11% 21%
1	F	149	68% 11% 21%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 5448 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lamin-B1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	118	908	566	157	182	3	0	0	0
1	B	118	908	566	157	182	3	0	0	0
1	C	118	908	566	157	182	3	0	0	0
1	D	118	908	566	157	182	3	0	0	0
1	F	118	908	566	157	182	3	0	0	0
1	E	118	908	566	157	182	3	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	405	GLY	-	expression tag	UNP P20700
A	406	ALA	-	expression tag	UNP P20700
B	405	GLY	-	expression tag	UNP P20700
B	406	ALA	-	expression tag	UNP P20700
C	405	GLY	-	expression tag	UNP P20700
C	406	ALA	-	expression tag	UNP P20700
D	405	GLY	-	expression tag	UNP P20700
D	406	ALA	-	expression tag	UNP P20700
F	405	GLY	-	expression tag	UNP P20700
F	406	ALA	-	expression tag	UNP P20700
E	405	GLY	-	expression tag	UNP P20700
E	406	ALA	-	expression tag	UNP P20700

THR
ILE
PRO
GLU
GLU

● Molecule 1: Lamin-B1



ILE
PRO
GLU
GLU

4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	144.65Å 160.75Å 132.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.83 – 3.60 49.83 – 3.60	Depositor EDS
% Data completeness (in resolution range)	89.2 (49.83-3.60) 89.2 (49.83-3.60)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.61 (at 3.57Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, R_{free}	0.222 , 0.262 0.221 , 0.261	Depositor DCC
R_{free} test set	1635 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	25.2	Xtrriage
Anisotropy	0.250	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 14.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.37$, $\langle L^2 \rangle = 0.20$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.81	EDS
Total number of atoms	5448	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/924	0.45	0/1252
1	B	0.24	0/924	0.44	0/1252
1	C	0.24	0/924	0.44	0/1252
1	D	0.24	0/924	0.45	0/1252
1	E	0.24	0/924	0.46	0/1252
1	F	0.24	0/924	0.44	0/1252
All	All	0.24	0/5544	0.45	0/7512

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	908	0	894	5	0
1	B	908	0	894	6	0
1	C	908	0	894	6	0
1	D	908	0	894	11	0
1	E	908	0	894	10	0
1	F	908	0	894	8	0
All	All	5448	0	5364	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 41 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:437:SER:HB3	1:B:541:GLN:HG2	1.68	0.75
1:B:433:SER:HB3	1:B:545:VAL:HG23	1.76	0.68
1:D:437:SER:HB3	1:D:541:GLN:HG2	1.76	0.66
1:F:457:LYS:HB2	1:F:495:THR:HG22	1.78	0.66
1:D:457:LYS:HG3	1:D:495:THR:HG22	1.77	0.64

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	116/149 (78%)	114 (98%)	2 (2%)	0	100	100
1	B	116/149 (78%)	113 (97%)	3 (3%)	0	100	100
1	C	116/149 (78%)	113 (97%)	3 (3%)	0	100	100
1	D	116/149 (78%)	111 (96%)	5 (4%)	0	100	100
1	E	116/149 (78%)	111 (96%)	5 (4%)	0	100	100
1	F	116/149 (78%)	112 (97%)	4 (3%)	0	100	100
All	All	696/894 (78%)	674 (97%)	22 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	100/127 (79%)	99 (99%)	1 (1%)	76	88
1	B	100/127 (79%)	97 (97%)	3 (3%)	41	71
1	C	100/127 (79%)	99 (99%)	1 (1%)	76	88
1	D	100/127 (79%)	100 (100%)	0	100	100
1	E	100/127 (79%)	100 (100%)	0	100	100
1	F	100/127 (79%)	99 (99%)	1 (1%)	76	88
All	All	600/762 (79%)	594 (99%)	6 (1%)	76	88

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	547	LYS
1	C	477	ASP
1	F	434	HIS
1	B	443	CYS
1	A	477	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	118/149 (79%)	-0.51	0 100 100	7, 26, 60, 78	0
1	B	118/149 (79%)	-0.47	0 100 100	7, 30, 64, 87	0
1	C	118/149 (79%)	-0.41	0 100 100	12, 36, 61, 77	0
1	D	118/149 (79%)	-0.34	0 100 100	17, 43, 77, 101	0
1	E	118/149 (79%)	-0.14	0 100 100	22, 58, 84, 89	0
1	F	118/149 (79%)	-0.11	1 (0%) 86 75	21, 59, 84, 106	0
All	All	708/894 (79%)	-0.33	1 (0%) 95 93	7, 40, 78, 106	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	547	LYS	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.