



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 27, 2024 – 09:00 AM EDT

PDB ID : 1DU3
Title : Crystal structure of TRAIL-SDR5
Authors : Cha, S.-S.; Sung, B.-J.; Oh, B.-H.
Deposited on : 2000-01-14
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	6634 (2.20-2.20)
Ramachandran outliers	177936	6560 (2.20-2.20)
Sidechain outliers	177891	6561 (2.20-2.20)





The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	130	25% 36% 8% 31%
1	B	130	34% 39% 12% 15%
1	C	130	18% 38% 12% 31%
1	G	130	33% 40% 10% 17%
1	H	130	28% 31% 10% 31%
1	I	130	22% 34% 12% 31%
2	D	168	46% 39% 5% 10%
2	E	168	40% 39% 9% 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	168	
2	J	168	
2	K	168	
2	L	168	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ZN	E	1	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 12108 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DEATH RECEPTOR 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	90	Total 701	C 419	N 122	O 148	S 12	0	0	0
1	B	110	Total 833	C 497	N 148	O 172	S 16	0	0	0
1	C	90	Total 697	C 417	N 121	O 147	S 12	0	0	0
1	G	108	Total 833	C 498	N 150	O 169	S 16	0	0	0
1	H	90	Total 691	C 414	N 118	O 147	S 12	0	0	0
1	I	90	Total 701	C 419	N 122	O 148	S 12	0	0	0

- Molecule 2 is a protein called TNF-RELATED APOPTOSIS INDUCING LIGAND.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	D	152	Total 1252	C 797	N 217	O 234	S 4	0	0	0
2	E	152	Total 1252	C 797	N 217	O 234	S 4	0	0	0
2	F	152	Total 1246	C 794	N 214	O 234	S 4	0	0	0
2	J	152	Total 1248	C 795	N 217	O 232	S 4	0	0	0
2	K	152	Total 1252	C 797	N 217	O 234	S 4	0	0	0
2	L	152	Total 1238	C 789	N 212	O 233	S 4	0	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	1	Total 1	Zn 1	0	0
3	K	1	Total 1	Zn 1	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	26	Total 26	O 26	0	0
4	B	23	Total 23	O 23	0	0
4	C	6	Total 6	O 6	0	0
4	D	22	Total 22	O 22	0	0
4	E	14	Total 14	O 14	0	0
4	F	13	Total 13	O 13	0	0
4	G	16	Total 16	O 16	0	0
4	H	7	Total 7	O 7	0	0
4	I	11	Total 11	O 11	0	0
4	J	8	Total 8	O 8	0	0
4	K	9	Total 9	O 9	0	0
4	L	7	Total 7	O 7	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: DEATH RECEPTOR 5




• Molecule 1: DEATH RECEPTOR 5



• Molecule 1: DEATH RECEPTOR 5




• Molecule 1: DEATH RECEPTOR 5








- Molecule 2: TNF-RELATED APOPTOSIS INDUCING LIGAND

Chain F: 36% 48% 7% 10%




- Molecule 2: TNF-RELATED APOPTOSIS INDUCING LIGAND

Chain J: 45% 40% 10%




- Molecule 2: TNF-RELATED APOPTOSIS INDUCING LIGAND


Chain K: 43% 40% 5% 10%



- Molecule 2: TNF-RELATED APOPTOSIS INDUCING LIGAND

Chain L: 36% 45% 10% 10%





4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	68.63Å 124.81Å 128.37Å 90.00° 104.49° 90.00°	Depositor
Resolution (Å)	8.00 – 2.20	Depositor
% Data completeness (in resolution range)	(Not available) (8.00-2.20)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	X-PLOR 3.851	Depositor
R, R_{free}	0.291 , 0.291	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	12108	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/716	0.82	0/969
1	B	2.32	1/849 (0.1%)	1.89	2/1149 (0.2%)
1	C	0.51	0/712	0.72	0/964
1	G	0.62	0/851	0.80	0/1150
1	H	0.58	0/706	0.79	0/957
1	I	0.98	1/716 (0.1%)	1.35	3/969 (0.3%)
2	D	0.86	5/1282 (0.4%)	0.81	1/1724 (0.1%)
2	E	0.61	0/1282	0.79	0/1724
2	F	0.69	1/1276 (0.1%)	0.82	0/1717
2	J	0.62	0/1278	0.78	0/1719
2	K	0.61	0/1282	0.78	0/1724
2	L	0.64	0/1268	0.82	0/1708
All	All	0.89	8/12218 (0.1%)	0.95	6/16474 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	F	0	1

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	130	SER	C-O	-64.68	0.00	1.23
1	I	117	CYS	C-N	21.03	1.82	1.34
2	D	198	GLU	CG-CD	13.28	1.71	1.51
2	D	198	GLU	CD-OE1	9.92	1.36	1.25
2	F	191	ARG	C-N	-9.60	1.11	1.34

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	130	SER	CA-C-O	-57.19	0.00	120.10
1	I	117	CYS	O-C-N	-25.30	82.22	122.70
1	I	117	CYS	C-N-CA	17.40	165.20	121.70
1	I	117	CYS	CA-C-N	17.17	154.98	117.20
1	B	127	HIS	O-C-N	-5.34	114.15	122.70

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	F	205	GLN	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	701	0	619	46	0
1	B	833	0	735	75	1
1	C	697	0	613	94	0
1	G	833	0	755	63	0
1	H	691	0	602	55	0
1	I	701	0	618	65	0
2	D	1252	0	1197	80	0
2	E	1252	0	1200	81	0
2	F	1246	0	1188	99	0
2	J	1248	0	1196	73	0
2	K	1252	0	1200	75	0
2	L	1238	0	1172	105	0
3	E	1	0	0	2	0
3	K	1	0	0	1	0
4	A	26	0	0	0	0
4	B	23	0	0	3	2
4	C	6	0	0	2	0
4	D	22	0	0	7	0
4	E	14	0	0	2	0
4	F	13	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	G	16	0	0	0	1
4	H	7	0	0	0	3
4	I	11	0	0	1	1
4	J	8	0	0	0	0
4	K	9	0	0	1	0
4	L	7	0	0	2	0
All	All	12108	0	11095	772	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 34.

The worst 5 of 772 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:117:CYS:C	1:I:118:THR:N	1.82	1.33
1:C:75:PRO:HB3	2:L:170:ARG:HD3	1.30	1.13
1:B:58:LEU:HD23	4:B:3147:HOH:O	1.59	1.02
2:K:202:ASN:HB3	4:K:3043:HOH:O	1.60	1.01
2:K:230:CYS:SG	2:L:230:CYS:SG	2.59	1.01

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:3160:HOH:O	4:H:3078:HOH:O[1_554]	0.57	1.63
4:G:3053:HOH:O	4:I:3050:HOH:O[2_656]	1.19	1.01
4:B:3160:HOH:O	4:H:3105:HOH:O[1_554]	1.82	0.38
1:B:129:GLU:CG	4:H:3079:HOH:O[1_554]	1.85	0.35

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	86/130 (66%)	77 (90%)	6 (7%)	3 (4%)	3	1
1	B	108/130 (83%)	93 (86%)	12 (11%)	3 (3%)	4	2
1	C	86/130 (66%)	61 (71%)	23 (27%)	2 (2%)	5	3
1	G	106/130 (82%)	90 (85%)	14 (13%)	2 (2%)	6	4
1	H	86/130 (66%)	75 (87%)	9 (10%)	2 (2%)	5	3
1	I	86/130 (66%)	72 (84%)	10 (12%)	4 (5%)	2	1
2	D	148/168 (88%)	132 (89%)	13 (9%)	3 (2%)	6	4
2	E	148/168 (88%)	131 (88%)	12 (8%)	5 (3%)	3	1
2	F	148/168 (88%)	124 (84%)	20 (14%)	4 (3%)	4	2
2	J	148/168 (88%)	132 (89%)	13 (9%)	3 (2%)	6	4
2	K	148/168 (88%)	130 (88%)	13 (9%)	5 (3%)	3	1
2	L	148/168 (88%)	125 (84%)	19 (13%)	4 (3%)	4	2
All	All	1446/1788 (81%)	1242 (86%)	164 (11%)	40 (3%)	4	2

5 of 40 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	E	130	ARG
2	E	157	SER
2	E	197	LYS
2	E	280	VAL
1	H	25	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	84/117 (72%)	64 (76%)	20 (24%)	0	0
1	B	97/117 (83%)	76 (78%)	21 (22%)	1	0
1	C	83/117 (71%)	62 (75%)	21 (25%)	0	0
1	G	99/117 (85%)	78 (79%)	21 (21%)	1	1

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	82/117 (70%)	61 (74%)	21 (26%)	0	0
1	I	84/117 (72%)	60 (71%)	24 (29%)	0	0
2	D	134/149 (90%)	117 (87%)	17 (13%)	3	3
2	E	134/149 (90%)	111 (83%)	23 (17%)	1	1
2	F	133/149 (89%)	119 (90%)	14 (10%)	5	5
2	J	133/149 (89%)	113 (85%)	20 (15%)	2	2
2	K	134/149 (90%)	114 (85%)	20 (15%)	2	2
2	L	131/149 (88%)	113 (86%)	18 (14%)	3	2
All	All	1328/1596 (83%)	1088 (82%)	240 (18%)	1	1

5 of 240 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	266	ILE
2	K	280	VAL
1	H	51	SER
2	K	259	SER
2	L	251	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 39 such sidechains are listed below:

Mol	Chain	Res	Type
2	J	228	ASN
2	L	161	HIS
2	J	270	HIS
2	K	202	ASN
2	L	253	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	I	1
2	D	1
2	F	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	I	117:CYS	C	118:THR	N	1.82
1	D	191:ARG	C	192:PHE	N	1.20
1	F	191:ARG	C	192:PHE	N	1.12

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.