



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 1, 2024 – 12:26 AM JST

PDB ID : 5DUR  
Title : Influenza A virus H5 hemagglutinin globular head in complex with antibody 100F4  
Authors : Zuo, T.; Sun, J.; Wang, G.; Zhou, P.; Wang, X.; Zhang, L.  
Deposited on : 2015-09-20  
Resolution : 2.82 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.82 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4293 (2.84-2.80)
Clashscore	180529	4801 (2.84-2.80)
Ramachandran outliers	177936	4739 (2.84-2.80)
Sidechain outliers	177891	4741 (2.84-2.80)
RSRZ outliers	164620	4295 (2.84-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	231	 3% 76% 14% 8%
1	H	231	 3% 69% 19% 7%
2	D	218	 2% 80% 17%
2	L	218	 % 76% 21%
3	A	233	 % 81% 12% 6%
3	C	233	 2% 75% 15% 9%

## 2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 10114 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Heavy Chain of Antibody 100F4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	H	214	Total 1604	C 1017	N 266	O 316	S 5	0	0	0
1	B	213	Total 1598	C 1013	N 265	O 315	S 5	0	0	0

- Molecule 2 is a protein called Light Chain of Antibody 100F4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	L	212	Total 1562	C 975	N 266	O 317	S 4	0	0	0
2	D	212	Total 1562	C 975	N 266	O 317	S 4	0	0	0

- Molecule 3 is a protein called Hemagglutinin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	212	Total 1708	C 1094	N 287	O 321	S 6	0	1	0
3	A	218	Total 1750	C 1118	N 298	O 328	S 6	0	0	0

There are 18 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
C	42	ALA	-	expression tag	UNP Q1WDM0
C	43	ASP	-	expression tag	UNP Q1WDM0
C	44	PRO	-	expression tag	UNP Q1WDM0
C	269	HIS	-	expression tag	UNP Q1WDM0
C	270	HIS	-	expression tag	UNP Q1WDM0
C	271	HIS	-	expression tag	UNP Q1WDM0
C	272	HIS	-	expression tag	UNP Q1WDM0
C	273	HIS	-	expression tag	UNP Q1WDM0

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	274	HIS	-	expression tag	UNP Q1WDM0
A	42	ALA	-	expression tag	UNP Q1WDM0
A	43	ASP	-	expression tag	UNP Q1WDM0
A	44	PRO	-	expression tag	UNP Q1WDM0
A	269	HIS	-	expression tag	UNP Q1WDM0
A	270	HIS	-	expression tag	UNP Q1WDM0
A	271	HIS	-	expression tag	UNP Q1WDM0
A	272	HIS	-	expression tag	UNP Q1WDM0
A	273	HIS	-	expression tag	UNP Q1WDM0
A	274	HIS	-	expression tag	UNP Q1WDM0

- Molecule 4 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
4	A	1	14	8	1	5	0	0
4	A	1	14	8	1	5	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
5	H	47	47	47	0	0
5	L	48	48	48	0	0

Continued on next page...


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
5	C	47	Total O 47 47	0	0
5	B	43	Total O 43 43	0	0
5	D	53	Total O 53 53	0	0
5	A	64	Total O 64 64	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Heavy Chain of Antibody 100F4





- Molecule 1: Heavy Chain of Antibody 100F4



- Molecule 2: Light Chain of Antibody 100F4




- Molecule 2: Light Chain of Antibody 100F4







- Molecule 3: Hemagglutinin



- Molecule 3: Hemagglutinin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.09Å 101.22Å 206.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.50 – 2.82 40.50 – 2.82	Depositor EDS
% Data completeness (in resolution range)	99.2 (40.50-2.82) 99.3 (40.50-2.82)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.88 (at 2.81Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.208 , 0.257 0.210 , 0.255	Depositor DCC
$R_{free}$ test set	2259 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	35.8	Xtrriage
Anisotropy	0.206	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 37.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	10114	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.53% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:  
NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	B	0.54	0/1639	0.72	0/2244
1	H	0.54	0/1645	0.71	1/2251 (0.0%)
2	D	0.50	0/1602	0.61	0/2190
2	L	0.51	0/1602	0.65	0/2190
3	A	0.50	0/1798	0.65	0/2447
3	C	0.51	0/1758	0.61	0/2392
All	All	0.52	0/10044	0.66	1/13714 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	H	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	H	46	GLY	N-CA-C	-5.11	100.32	113.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	H	79	HIS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1598	0	1575	25	0
1	H	1604	0	1582	32	0
2	D	1562	0	1513	21	0
2	L	1562	0	1513	24	0
3	A	1750	0	1700	19	0
3	C	1708	0	1654	21	0
4	A	28	0	26	0	0
5	A	64	0	0	4	0
5	B	43	0	0	0	0
5	C	47	0	0	4	0
5	D	53	0	0	1	0
5	H	47	0	0	2	0
5	L	48	0	0	2	0
All	All	10114	0	9563	122	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 122 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:206:ASN:ND2	5:A:401:HOH:O	2.08	0.86
3:A:216:ARG:NH2	5:A:402:HOH:O	2.11	0.84
2:L:63:ARG:NH2	2:L:84:ASP:OD2	2.12	0.82
1:B:104:THR:HG23	3:A:113:LYS:H	1.43	0.82
1:B:52:ASN:HD21	1:B:106:THR:HG21	1.48	0.79

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	209/231 (90%)	195 (93%)	13 (6%)	1 (0%)	25	54
1	H	210/231 (91%)	195 (93%)	9 (4%)	6 (3%)	3	12
2	D	210/218 (96%)	206 (98%)	4 (2%)	0	100	100
2	L	210/218 (96%)	205 (98%)	5 (2%)	0	100	100
3	A	216/233 (93%)	205 (95%)	11 (5%)	0	100	100
3	C	209/233 (90%)	200 (96%)	8 (4%)	1 (0%)	25	54
All	All	1264/1364 (93%)	1206 (95%)	50 (4%)	8 (1%)	22	49

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	29	VAL
1	H	45	LYS
1	B	45	LYS
3	C	90	CYS
1	H	31	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	185/201 (92%)	174 (94%)	11 (6%)	16	43
1	H	186/201 (92%)	173 (93%)	13 (7%)	12	34
2	D	174/180 (97%)	167 (96%)	7 (4%)	27	58
2	L	174/180 (97%)	161 (92%)	13 (8%)	11	31
3	A	195/208 (94%)	186 (95%)	9 (5%)	23	53
3	C	190/208 (91%)	179 (94%)	11 (6%)	17	43
All	All	1104/1178 (94%)	1040 (94%)	64 (6%)	17	43

5 of 64 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	145	ARG
3	A	172	LEU
2	L	171	SER
2	L	161	VAL
3	A	188	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
2	L	40	GLN
2	L	118	ASN
3	A	220	ASN
3	A	206	ASN
1	H	113	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	A	302	3	14,14,15	0.60	1 (7%)	17,19,21	0.85	0
4	NAG	A	301	3	14,14,15	0.99	1 (7%)	17,19,21	1.01	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	302	3	-	2/6/23/26	0/1/1/1
4	NAG	A	301	3	-	0/6/23/26	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	301	NAG	O5-C1	3.38	1.49	1.43
4	A	302	NAG	O5-C1	2.10	1.47	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	301	NAG	C1-O5-C5	2.26	115.25	112.19
4	A	301	NAG	C3-C4-C5	-2.05	106.58	110.24

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	302	NAG	O5-C5-C6-O6
4	A	302	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	B	213/231 (92%)	-0.08	8 (3%) 44 36	21, 29, 51, 65	0
1	H	214/231 (92%)	-0.19	8 (3%) 45 37	21, 29, 49, 57	0
2	D	212/218 (97%)	-0.21	4 (1%) 66 59	23, 32, 38, 48	0
2	L	212/218 (97%)	-0.15	2 (0%) 81 75	24, 32, 44, 50	0
3	A	218/233 (93%)	-0.24	3 (1%) 73 66	23, 32, 47, 54	0
3	C	212/233 (90%)	-0.16	4 (1%) 66 59	20, 33, 47, 53	1 (0%)
All	All	1281/1364 (93%)	-0.17	29 (2%) 61 53	20, 31, 46, 65	1 (0%)

The worst 5 of 29 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	27	ASP	4.6
2	D	113	GLY	3.9
1	B	45	LYS	3.7
1	B	197	LEU	3.1
1	B	30	SER	3.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NAG	A	301	14/15	0.85	0.15	27,27,27,27	0
4	NAG	A	302	14/15	0.88	0.11	27,27,27,27	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.