

wwPDB X-ray Structure Validation Summary Report (i)

Sep 5, 2023 – 10:29 PM EDT

PDB ID : 4DVT

Title: Crystal structure of clade A/E 93TH057 HIV-1 gp120 core in complex with

AS-II-37

Authors: Kwon, Y.D.; Kwong, P.D.

Deposited on : 2012-02-23

Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
Metric	$(\# ext{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain							
1	A	353	69%	27%						
1	В	353	7% 61%	34%						

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NAG	A	502	-	-	-	X
2	NAG	A	509	X	-	-	-
2	NAG	A	510	-	-	-	X
2	NAG	В	504	-	-	-	X
2	NAG	В	508	-	-	-	X
2	NAG	В	510	-	-	-	X
3	0LZ	В	512	-	-	-	X



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 5837 atoms, of which 34 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


• Molecule 1 is a protein called clade A/E 93TH057 HIV-1 gp120 core.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	Δ	339	Total	С	N	О	S	0	0	0
	11	000	2654	1666	460	507	21	0	O	
1	D	339	Total	С	N	О	S	0	0	0
1	Б	339	2654	1666	460	507	21	U	0	

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	375	SER	HIS	engineered mutation	UNP A0A0M3KKW9
В	375	SER	HIS	engineered mutation	UNP A0A0M3KKW9

• Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



I	Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
	2	A	1	Total	С	N 1	O 5	0	0

Continued on next page...




 $Continued\ from\ previous\ page...$

Mol	Chain	$oxed{ \mathbf{Residues} }$		ton	ns		ZeroOcc	AltConf
			Total	C	N	О		
2	A	1	14	8	1	5	0	0
2	A	1	Total	С	N	О	0	0
	71	1	14	8	1	5	0	0
2	A	1	Total	С	N	O	0	0
		_	14	8	1	5		
2	A	1	Total	С	N	O	0	0
			14	8	1	5		
2	A	1	Total	C	N	Ō	0	0
			14	8	1	5		
2	A	1	Total	С	N	O	0	0
			14	8	1	5		
2	A	1	Total	С	N	O	0	0
			14	8	1	5		
2	A	1	Total	С	N	O	0	0
			14	8	1	5		
2	A	1	Total	С	N	O	0	0
			14	8	1	5		
2	A	1	Total	$^{\circ}$	N	O	0	0
			14	8 C	$\frac{1}{N}$	5		
2	В	1	Total			O	0	0
			14	8 C	$\frac{1}{N}$	5 O		
2	В	1	Total 14	8			0	0
			Total	<u> </u>	$\frac{1}{N}$	5 O		
2	В	1	10tai	8	1	5	0	0
			Total	$\frac{\circ}{\mathrm{C}}$	N	$\frac{0}{0}$		
2	В	1	10tar	8	1	5	0	0
			Total	$\frac{6}{C}$	N	$\frac{0}{0}$		
2	В	1	10tar	8	1	5	0	0
			Total	$\frac{\sigma}{C}$	N	$\frac{0}{0}$		
2	В	1	10tai	8	1	5	0	0
			Total	$\frac{c}{C}$	N	O		
2	В	1	10tar	8	1	5	0	0
			Total	$\frac{6}{C}$	N	$\frac{0}{0}$		
2	В	1	10tar	8	1	5	0	0
			Total	$\frac{c}{C}$	N	0		
2	В	1	14	8	1	5	0	0
			Total	$\frac{c}{C}$	N	O		
2	В	1	14	8	1	5	0	0
			Total	$\frac{c}{C}$	N	O		
2	В	1	14	8	1	5	0	0
			1.1	U	1	9		




• Molecule 3 is N-(4-chloro-3-fluorophenyl)-N'-[(1S)-1,2,3,4-tetrahydroisoquinolin-1-ylmethyl] ethanediamide (three-letter code: 0LZ) (formula: $C_{18}H_{17}ClFN_3O_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
2	Λ	1	Total	С	Cl	F	Н	N	О	0	0
3	A	1	42	18	1	1	17	3	2	0	
2	D	1	Total	С	Cl	F	Н	N	0	0	0
3	3 B	1	42	18	1	1	17	3	2	0	U

• Molecule 4 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (three-letter code: EPE) (formula: $C_8H_{18}N_2O_4S$).





\mathbf{Mol}	Chain	Residues	Atoms					ZeroOcc	AltConf	
1	Λ	1	Total	С	N	О	S	0	0	
4	4 A	1	15	8	2	4	1	0		
1	D	1	Total	С	N	О	S	0	0	
4	Ъ	1	15	8	2	4	1	0	U	

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	72	Total O 72 72	0	0
5	В	35	Total O 35 35	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: clade A/E 93TH057 HIV-1 gp120 core



• Molecule 1: clade A/E 93TH057 HIV-1 gp120 core





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	64.81Å 68.40Å 93.99Å	Donogitor
a, b, c, α , β , γ	90.00° 91.94° 90.00°	Depositor
Resolution (Å)	28.19 - 2.40	Depositor
rtesolution (A)	46.97 - 2.43	EDS
% Data completeness	87.2 (28.19-2.40)	Depositor
(in resolution range)	84.3 (46.97-2.43)	EDS
R_{merge}	0.07	Depositor
R_{sym}	0.09	Depositor
$< I/\sigma(I) > 1$	$0.08 \; (at \; 2.42 \text{Å})$	Xtriage
Refinement program	PHENIX 1.8_1069	Depositor
D D.	0.196 , 0.248	Depositor
R, R_{free}	0.202 , 0.252	DCC
R_{free} test set	1338 reflections (4.80%)	wwPDB-VP
Wilson B-factor (Å ²)	45.8	Xtriage
Anisotropy	0.289	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.31 , 51.1	EDS
L-test for twinning ²	$< L >=0.49, < L^2>=0.32$	Xtriage
Estimated twinning fraction	0.029 for h,-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5837	wwPDB-VP
Average B, all atoms (Å ²)	78.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, NAG, 0LZ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Clasia.		nd lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.24	$1/2709 \ (0.0\%)$	0.39	1/3678 (0.0%)	
1	В	0.20	0/2709	0.37	0/3678	
All	All	0.22	1/5418 (0.0%)	0.38	1/7356 (0.0%)	

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(ext{\AA})$
1	A	355	ASN	C-N	-6.39	1.19	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}(^{o})$
1	A	354	ASN	O-C-N	5.03	130.75	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2654	0	2588	89	0
1	В	2654	0	2588	108	0
2	A	154	0	143	9	0
2	В	154	0	143	6	0

Continued on next page...



I 'omtamalod	trom	mmonia	maaa
Continued	11 0116	DICUIUUS	Daue
	.,	10	1

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	25	17	17	2	0
3	В	25	17	17	1	0
4	A	15	0	17	3	0
4	В	15	0	17	3	0
5	A	72	0	0	4	0
5	В	35	0	0	4	0
All	All	5803	34	5530	197	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

The worst 5 of 197 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:A:224:VAL:HG22	1:A:491:ILE:HD11	1.45	0.96
1:A:480:ARG:NH2	5:A:619:HOH:O	1.98	0.94
1:A:65:VAL:HG11	1:A:208:ILE:HD12	1.51	0.92
1:B:289:ASN:O	5:B:631:HOH:O	1.92	0.88
1:A:343:LYS:HD3	1:A:396:ILE:HG23	1.61	0.83

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Favoured Allowed		Allowed	Outliers	Perce	entiles
1	A	333/353~(94%)	314 (94%)	18 (5%)	1 (0%)	41	55
1	В	333/353 (94%)	306 (92%)	23 (7%)	4 (1%)	13	19
All	All	666/706 (94%)	620 (93%)	41 (6%)	5 (1%)	19	29

All (5) Ramachandran outliers are listed below:



Mol	Chain	Res	Type
1	В	430	THR
1	A	124	GLY
1	В	460	ALA
1	В	393	ASN
1	В	471	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	303/311 (97%)	302 (100%)	1 (0%)	92 97
1	В	303/311 (97%)	303 (100%)	0	100 100
All	All	606/622 (97%)	605 (100%)	1 (0%)	93 97

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	348	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	465	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

26 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Trino	Chain	Dag	T inle	Во	nd leng	ths	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	В	508	1	14,14,15	0.54	0	17,19,21	0.63	0
2	NAG	A	506	1	14,14,15	0.56	0	17,19,21	0.60	0
2	NAG	В	509	1	14,14,15	0.52	0	17,19,21	0.66	0
2	NAG	A	508	1	14,14,15	0.57	0	17,19,21	0.61	0
2	NAG	A	510	1	14,14,15	0.54	0	17,19,21	0.61	0
2	NAG	A	509	1	14,14,15	0.56	0	17,19,21	0.62	0
4	EPE	В	513	_	15,15,15	0.83	1 (6%)	18,20,20	1.85	4 (22%)
2	NAG	В	505	1	14,14,15	0.53	0	17,19,21	0.61	0
2	NAG	В	503	1	14,14,15	0.50	0	17,19,21	0.81	0
2	NAG	В	502	1	14,14,15	0.52	0	17,19,21	0.70	0
2	NAG	A	501	1	14,14,15	0.55	0	17,19,21	0.59	0
2	NAG	В	506	1	14,14,15	0.56	0	17,19,21	0.61	0
2	NAG	A	502	1	14,14,15	0.54	0	17,19,21	0.64	0
2	NAG	A	511	1	14,14,15	0.52	0	17,19,21	0.71	0
2	NAG	A	504	1	14,14,15	0.53	0	17,19,21	0.64	0
3	0LZ	A	512	_	26,27,27	1.47	5 (19%)	33,37,37	1.86	5 (15%)
2	NAG	A	507	1	14,14,15	0.54	0	17,19,21	0.63	0
2	NAG	В	507	1	14,14,15	0.53	0	17,19,21	0.65	0
3	0LZ	В	512	-	26,27,27	1.49	5 (19%)	33,37,37	1.86	5 (15%)
4	EPE	A	513	-	15,15,15	0.81	1 (6%)	18,20,20	1.81	5 (27%)
2	NAG	В	511	1	14,14,15	0.53	0	17,19,21	0.70	0
2	NAG	В	504	1	14,14,15	0.51	0	17,19,21	0.67	0
2	NAG	A	503	1	14,14,15	0.53	0	17,19,21	0.68	0
2	NAG	A	505	1	14,14,15	0.53	0	17,19,21	0.65	0
2	NAG	В	510	1	14,14,15	0.54	0	17,19,21	0.67	0
2	NAG	В	501	1	14,14,15	0.53	0	17,19,21	0.64	0



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	В	508	1	-	2/6/23/26	0/1/1/1
2	NAG	A	506	1	-	0/6/23/26	0/1/1/1
2	NAG	В	509	1	-	2/6/23/26	0/1/1/1
2	NAG	A	508	1	-	2/6/23/26	0/1/1/1
2	NAG	A	510	1	-	3/6/23/26	0/1/1/1
2	NAG	A	509	1	1/1/5/7	2/6/23/26	0/1/1/1
4	EPE	В	513	-	-	6/9/19/19	0/1/1/1
2	NAG	В	505	1	-	2/6/23/26	0/1/1/1
2	NAG	В	503	1	-	0/6/23/26	0/1/1/1
2	NAG	В	502	1	-	0/6/23/26	0/1/1/1
2	NAG	A	501	1	-	0/6/23/26	0/1/1/1
2	NAG	В	506	1	-	2/6/23/26	0/1/1/1
2	NAG	A	502	1	-	2/6/23/26	0/1/1/1
2	NAG	A	511	1	-	2/6/23/26	0/1/1/1
2	NAG	A	504	1	-	0/6/23/26	0/1/1/1
3	0LZ	A	512	-	-	0/13/23/23	0/3/3/3
2	NAG	A	507	1	-	0/6/23/26	0/1/1/1
2	NAG	В	507	1	-	4/6/23/26	0/1/1/1
3	0LZ	В	512	-	-	0/13/23/23	0/3/3/3
4	EPE	A	513	-	-	6/9/19/19	0/1/1/1
2	NAG	В	511	1	-	2/6/23/26	0/1/1/1
2	NAG	В	504	1	-	0/6/23/26	0/1/1/1
2	NAG	A	503	1	-	0/6/23/26	0/1/1/1
2	NAG	A	505	1	-	0/6/23/26	0/1/1/1
2	NAG	В	510	1	-	3/6/23/26	0/1/1/1
2	NAG	В	501	1	-	2/6/23/26	0/1/1/1

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(A)	Ideal(Å)
3	В	512	0LZ	C02-N01	3.56	1.39	1.33
3	A	512	0LZ	C02-N01	3.47	1.39	1.33
3	В	512	0LZ	C03-N04	3.08	1.42	1.35
3	A	512	0LZ	C03-N04	3.04	1.42	1.35
3	В	512	0LZ	C03-C02	2.84	1.58	1.53



The worst	5	of	19	bond	angle	outliers	are	listed	below:
TITO HOLDO	$\overline{}$	O.	10	OIIG	WII SIC	Cathere	COL C	IID CCC	CIC III.

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	В	512	0LZ	C02-C03-N04	5.64	121.22	112.31
3	A	512	0LZ	C02-C03-N04	5.60	121.17	112.31
3	В	512	0LZ	C03-C02-N01	5.45	119.72	113.47
3	A	512	0LZ	C03-C02-N01	5.44	119.71	113.47
4	В	513	EPE	C5-N4-C3	4.33	118.58	108.83

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	A	509	NAG	C1

5 of 42 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	502	NAG	C8-C7-N2-C2
2	A	502	NAG	O7-C7-N2-C2
2	A	509	NAG	C8-C7-N2-C2
2	A	509	NAG	O7-C7-N2-C2
2	В	507	NAG	C1-C2-N2-C7

There are no ring outliers.


13 monomers are involved in 24 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	509	NAG	3	0
2	A	508	NAG	5	0
2	A	510	NAG	1	0
4	В	513	EPE	3	0
2	В	505	NAG	1	0
2	A	511	NAG	1	0
2	A	504	NAG	1	0
3	A	512	0LZ	2	0
2	В	507	NAG	1	0
3	В	512	0LZ	1	0
4	A	513	EPE	3	0
2	В	511	NAG	1	0
2	A	505	NAG	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will



also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	A	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	A	355:ASN	С	357:LYS	N	1.19



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# \mathrm{RSRZ}{>}2$	$OWAB(Å^2)$	Q < 0.9
1	A	339/353~(96%)	-0.01	9 (2%) 54 52	45, 63, 105, 137	0
1	В	339/353~(96%)	0.44	24 (7%) 16 14	49, 80, 129, 190	0
All	All	678/706 (96%)	0.22	33 (4%) 29 28	45, 71, 120, 190	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	350	LYS	6.1
1	В	124	GLY	5.7
1	В	492	GLU	4.8
1	В	462	ASN	4.7
1	В	44	VAL	4.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
2	NAG	В	508	14/15	0.59	0.69	132,147,157,158	0
2	NAG	A	510	14/15	0.60	0.44	117,161,177,181	0
2	NAG	A	502	14/15	0.62	0.56	120,130,137,139	0
2	NAG	В	507	14/15	0.73	0.32	69,112,127,128	0
2	NAG	A	511	14/15	0.73	0.28	84,105,110,113	0
2	NAG	В	504	14/15	0.75	0.45	100,108,117,118	0
2	NAG	В	502	14/15	0.76	0.40	115,125,147,147	0
2	NAG	В	510	14/15	0.76	0.58	178,194,202,206	0
3	0LZ	В	512	25/25	0.77	0.42	89,127,199,202	0
2	NAG	В	509	14/15	0.79	0.21	86,103,111,119	0
2	NAG	В	511	14/15	0.81	0.25	92,101,106,112	0
2	NAG	В	501	14/15	0.83	0.24	87,93,100,100	0
2	NAG	A	507	14/15	0.86	0.24	76,97,122,132	0
2	NAG	A	509	14/15	0.87	0.16	76,94,99,99	0
2	NAG	В	505	14/15	0.87	0.15	79,104,114,115	0
2	NAG	В	506	14/15	0.88	0.17	66,88,101,103	0
3	0LZ	A	512	25/25	0.89	0.23	64,98,152,156	0
2	NAG	A	506	14/15	0.89	0.25	74,89,97,99	0
2	NAG	A	501	14/15	0.91	0.15	69,83,95,97	0
2	NAG	A	504	14/15	0.91	0.14	70,84,92,94	0
2	NAG	A	508	14/15	0.91	0.16	54,59,72,86	0
2	NAG	A	505	14/15	0.92	0.19	58,69,92,99	0
2	NAG	В	503	14/15	0.94	0.14	49,54,63,64	0
4	EPE	A	513	15/15	0.95	0.21	49,62,81,97	0
2	NAG	A	503	14/15	0.96	0.13	46,62,67,68	0
4	EPE	В	513	15/15	0.96	0.18	42,70,84,99	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 0LZ B 512:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around 0LZ A 512:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

