



wwPDB X-ray Structure Validation Summary Report ⓘ

May 23, 2020 – 08:35 pm BST

PDB ID : 5E9A
Title : Crystal structure analysis of the cold-adapted beta-galactosidase from *Rahnella* sp. R3
Authors : Zhang, Y.Z.; Fan, Y.T.
Deposited on : 2015-10-14
Resolution : 2.56 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.56 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1279 (2.58-2.54)
Clashscore	141614	1327 (2.58-2.54)
Ramachandran outliers	138981	1312 (2.58-2.54)
Sidechain outliers	138945	1312 (2.58-2.54)
RSRZ outliers	127900	1269 (2.58-2.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	712	
1	B	712	
1	C	712	
1	D	712	
1	E	712	
1	F	712	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	ACT	F	702	-	-	X	-

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 32981 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-galactosidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	679	Total 5387	C 3420	N 945	O 993	S 29	0	0	0
1	B	684	Total 5424	C 3443	N 952	O 1000	S 29	0	0	0
1	C	684	Total 5424	C 3443	N 952	O 1000	S 29	0	0	0
1	D	684	Total 5424	C 3443	N 952	O 1000	S 29	0	0	0
1	E	684	Total 5432	C 3448	N 955	O 1000	S 29	0	1	0
1	F	684	Total 5432	C 3448	N 955	O 1000	S 29	0	1	0

There are 150 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-24	MET	-	initiating methionine	UNP A0A0B4U8I5
A	-23	ASN	-	expression tag	UNP A0A0B4U8I5
A	-22	HIS	-	expression tag	UNP A0A0B4U8I5
A	-21	LYS	-	expression tag	UNP A0A0B4U8I5
A	-20	VAL	-	expression tag	UNP A0A0B4U8I5
A	-19	HIS	-	expression tag	UNP A0A0B4U8I5
A	-18	HIS	-	expression tag	UNP A0A0B4U8I5
A	-17	HIS	-	expression tag	UNP A0A0B4U8I5
A	-16	HIS	-	expression tag	UNP A0A0B4U8I5
A	-15	HIS	-	expression tag	UNP A0A0B4U8I5
A	-14	HIS	-	expression tag	UNP A0A0B4U8I5
A	-13	ILE	-	expression tag	UNP A0A0B4U8I5
A	-12	GLU	-	expression tag	UNP A0A0B4U8I5
A	-11	GLY	-	expression tag	UNP A0A0B4U8I5
A	-10	ARG	-	expression tag	UNP A0A0B4U8I5
A	-9	HIS	-	expression tag	UNP A0A0B4U8I5
A	-8	MET	-	expression tag	UNP A0A0B4U8I5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	GLU	-	expression tag	UNP A0A0B4U8I5
A	-6	LEU	-	expression tag	UNP A0A0B4U8I5
A	-5	GLY	-	expression tag	UNP A0A0B4U8I5
A	-4	THR	-	expression tag	UNP A0A0B4U8I5
A	-3	LEU	-	expression tag	UNP A0A0B4U8I5
A	-2	GLU	-	expression tag	UNP A0A0B4U8I5
A	-1	GLY	-	expression tag	UNP A0A0B4U8I5
A	0	SER	-	expression tag	UNP A0A0B4U8I5
B	-24	MET	-	initiating methionine	UNP A0A0B4U8I5
B	-23	ASN	-	expression tag	UNP A0A0B4U8I5
B	-22	HIS	-	expression tag	UNP A0A0B4U8I5
B	-21	LYS	-	expression tag	UNP A0A0B4U8I5
B	-20	VAL	-	expression tag	UNP A0A0B4U8I5
B	-19	HIS	-	expression tag	UNP A0A0B4U8I5
B	-18	HIS	-	expression tag	UNP A0A0B4U8I5
B	-17	HIS	-	expression tag	UNP A0A0B4U8I5
B	-16	HIS	-	expression tag	UNP A0A0B4U8I5
B	-15	HIS	-	expression tag	UNP A0A0B4U8I5
B	-14	HIS	-	expression tag	UNP A0A0B4U8I5
B	-13	ILE	-	expression tag	UNP A0A0B4U8I5
B	-12	GLU	-	expression tag	UNP A0A0B4U8I5
B	-11	GLY	-	expression tag	UNP A0A0B4U8I5
B	-10	ARG	-	expression tag	UNP A0A0B4U8I5
B	-9	HIS	-	expression tag	UNP A0A0B4U8I5
B	-8	MET	-	expression tag	UNP A0A0B4U8I5
B	-7	GLU	-	expression tag	UNP A0A0B4U8I5
B	-6	LEU	-	expression tag	UNP A0A0B4U8I5
B	-5	GLY	-	expression tag	UNP A0A0B4U8I5
B	-4	THR	-	expression tag	UNP A0A0B4U8I5
B	-3	LEU	-	expression tag	UNP A0A0B4U8I5
B	-2	GLU	-	expression tag	UNP A0A0B4U8I5
B	-1	GLY	-	expression tag	UNP A0A0B4U8I5
B	0	SER	-	expression tag	UNP A0A0B4U8I5
C	-24	MET	-	initiating methionine	UNP A0A0B4U8I5
C	-23	ASN	-	expression tag	UNP A0A0B4U8I5
C	-22	HIS	-	expression tag	UNP A0A0B4U8I5
C	-21	LYS	-	expression tag	UNP A0A0B4U8I5
C	-20	VAL	-	expression tag	UNP A0A0B4U8I5
C	-19	HIS	-	expression tag	UNP A0A0B4U8I5
C	-18	HIS	-	expression tag	UNP A0A0B4U8I5
C	-17	HIS	-	expression tag	UNP A0A0B4U8I5
C	-16	HIS	-	expression tag	UNP A0A0B4U8I5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-15	HIS	-	expression tag	UNP A0A0B4U8I5
C	-14	HIS	-	expression tag	UNP A0A0B4U8I5
C	-13	ILE	-	expression tag	UNP A0A0B4U8I5
C	-12	GLU	-	expression tag	UNP A0A0B4U8I5
C	-11	GLY	-	expression tag	UNP A0A0B4U8I5
C	-10	ARG	-	expression tag	UNP A0A0B4U8I5
C	-9	HIS	-	expression tag	UNP A0A0B4U8I5
C	-8	MET	-	expression tag	UNP A0A0B4U8I5
C	-7	GLU	-	expression tag	UNP A0A0B4U8I5
C	-6	LEU	-	expression tag	UNP A0A0B4U8I5
C	-5	GLY	-	expression tag	UNP A0A0B4U8I5
C	-4	THR	-	expression tag	UNP A0A0B4U8I5
C	-3	LEU	-	expression tag	UNP A0A0B4U8I5
C	-2	GLU	-	expression tag	UNP A0A0B4U8I5
C	-1	GLY	-	expression tag	UNP A0A0B4U8I5
C	0	SER	-	expression tag	UNP A0A0B4U8I5
D	-24	MET	-	initiating methionine	UNP A0A0B4U8I5
D	-23	ASN	-	expression tag	UNP A0A0B4U8I5
D	-22	HIS	-	expression tag	UNP A0A0B4U8I5
D	-21	LYS	-	expression tag	UNP A0A0B4U8I5
D	-20	VAL	-	expression tag	UNP A0A0B4U8I5
D	-19	HIS	-	expression tag	UNP A0A0B4U8I5
D	-18	HIS	-	expression tag	UNP A0A0B4U8I5
D	-17	HIS	-	expression tag	UNP A0A0B4U8I5
D	-16	HIS	-	expression tag	UNP A0A0B4U8I5
D	-15	HIS	-	expression tag	UNP A0A0B4U8I5
D	-14	HIS	-	expression tag	UNP A0A0B4U8I5
D	-13	ILE	-	expression tag	UNP A0A0B4U8I5
D	-12	GLU	-	expression tag	UNP A0A0B4U8I5
D	-11	GLY	-	expression tag	UNP A0A0B4U8I5
D	-10	ARG	-	expression tag	UNP A0A0B4U8I5
D	-9	HIS	-	expression tag	UNP A0A0B4U8I5
D	-8	MET	-	expression tag	UNP A0A0B4U8I5
D	-7	GLU	-	expression tag	UNP A0A0B4U8I5
D	-6	LEU	-	expression tag	UNP A0A0B4U8I5
D	-5	GLY	-	expression tag	UNP A0A0B4U8I5
D	-4	THR	-	expression tag	UNP A0A0B4U8I5
D	-3	LEU	-	expression tag	UNP A0A0B4U8I5
D	-2	GLU	-	expression tag	UNP A0A0B4U8I5
D	-1	GLY	-	expression tag	UNP A0A0B4U8I5
D	0	SER	-	expression tag	UNP A0A0B4U8I5
E	-24	MET	-	initiating methionine	UNP A0A0B4U8I5

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-23	ASN	-	expression tag	UNP A0A0B4U8I5
E	-22	HIS	-	expression tag	UNP A0A0B4U8I5
E	-21	LYS	-	expression tag	UNP A0A0B4U8I5
E	-20	VAL	-	expression tag	UNP A0A0B4U8I5
E	-19	HIS	-	expression tag	UNP A0A0B4U8I5
E	-18	HIS	-	expression tag	UNP A0A0B4U8I5
E	-17	HIS	-	expression tag	UNP A0A0B4U8I5
E	-16	HIS	-	expression tag	UNP A0A0B4U8I5
E	-15	HIS	-	expression tag	UNP A0A0B4U8I5
E	-14	HIS	-	expression tag	UNP A0A0B4U8I5
E	-13	ILE	-	expression tag	UNP A0A0B4U8I5
E	-12	GLU	-	expression tag	UNP A0A0B4U8I5
E	-11	GLY	-	expression tag	UNP A0A0B4U8I5
E	-10	ARG	-	expression tag	UNP A0A0B4U8I5
E	-9	HIS	-	expression tag	UNP A0A0B4U8I5
E	-8	MET	-	expression tag	UNP A0A0B4U8I5
E	-7	GLU	-	expression tag	UNP A0A0B4U8I5
E	-6	LEU	-	expression tag	UNP A0A0B4U8I5
E	-5	GLY	-	expression tag	UNP A0A0B4U8I5
E	-4	THR	-	expression tag	UNP A0A0B4U8I5
E	-3	LEU	-	expression tag	UNP A0A0B4U8I5
E	-2	GLU	-	expression tag	UNP A0A0B4U8I5
E	-1	GLY	-	expression tag	UNP A0A0B4U8I5
E	0	SER	-	expression tag	UNP A0A0B4U8I5
F	-24	MET	-	initiating methionine	UNP A0A0B4U8I5
F	-23	ASN	-	expression tag	UNP A0A0B4U8I5
F	-22	HIS	-	expression tag	UNP A0A0B4U8I5
F	-21	LYS	-	expression tag	UNP A0A0B4U8I5
F	-20	VAL	-	expression tag	UNP A0A0B4U8I5
F	-19	HIS	-	expression tag	UNP A0A0B4U8I5
F	-18	HIS	-	expression tag	UNP A0A0B4U8I5
F	-17	HIS	-	expression tag	UNP A0A0B4U8I5
F	-16	HIS	-	expression tag	UNP A0A0B4U8I5
F	-15	HIS	-	expression tag	UNP A0A0B4U8I5
F	-14	HIS	-	expression tag	UNP A0A0B4U8I5
F	-13	ILE	-	expression tag	UNP A0A0B4U8I5
F	-12	GLU	-	expression tag	UNP A0A0B4U8I5
F	-11	GLY	-	expression tag	UNP A0A0B4U8I5
F	-10	ARG	-	expression tag	UNP A0A0B4U8I5
F	-9	HIS	-	expression tag	UNP A0A0B4U8I5
F	-8	MET	-	expression tag	UNP A0A0B4U8I5
F	-7	GLU	-	expression tag	UNP A0A0B4U8I5

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	-6	LEU	-	expression tag	UNP A0A0B4U8I5
F	-5	GLY	-	expression tag	UNP A0A0B4U8I5
F	-4	THR	-	expression tag	UNP A0A0B4U8I5
F	-3	LEU	-	expression tag	UNP A0A0B4U8I5
F	-2	GLU	-	expression tag	UNP A0A0B4U8I5
F	-1	GLY	-	expression tag	UNP A0A0B4U8I5
F	0	SER	-	expression tag	UNP A0A0B4U8I5

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	1	Total Zn 1 1	0	0
2	E	1	Total Zn 1 1	0	0
2	B	1	Total Zn 1 1	0	0
2	C	1	Total Zn 1 1	0	0
2	A	1	Total Zn 1 1	0	0
2	F	1	Total Zn 1 1	0	0

- Molecule 3 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	E	1	Total C O 4 2 2	0	0
3	F	1	Total C O 4 2 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	105	Total O 105 105	0	0
4	B	56	Total O 56 56	0	0
4	C	66	Total O 66 66	0	0
4	D	99	Total O 99 99	0	0
4	E	48	Total O 48 48	0	0
4	F	54	Total O 54 54	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Beta-galactosidase




- Molecule 1: Beta-galactosidase




- Molecule 1: Beta-galactosidase





- Molecule 1: Beta-galactosidase

Chain D:  88% 8%





• Molecule 1: Beta-galactosidase

Chain E:  86% 9%



• Molecule 1: Beta-galactosidase

Chain F:  88% 7%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	146.43Å 106.83Å 164.24Å 90.00° 109.00° 90.00°	Depositor
Resolution (Å)	138.45 – 2.56 138.45 – 2.56	Depositor EDS
% Data completeness (in resolution range)	98.5 (138.45-2.56) 98.5 (138.45-2.56)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 2.55Å)	Xtrriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.199 , 0.247 0.204 , 0.249	Depositor DCC
R_{free} test set	2020 reflections (1.33%)	wwPDB-VP
Wilson B-factor (Å ²)	29.6	Xtrriage
Anisotropy	0.099	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	32981	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/5540	0.85	11/7537 (0.1%)
1	B	0.65	0/5579	0.81	9/7592 (0.1%)
1	C	0.65	0/5579	0.82	13/7592 (0.2%)
1	D	0.69	0/5579	0.83	8/7592 (0.1%)
1	E	0.63	0/5590	0.81	9/7607 (0.1%)
1	F	0.63	0/5590	0.81	10/7606 (0.1%)
All	All	0.66	0/33457	0.82	60/45526 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	D	0	1
1	E	0	2
1	F	0	1
All	All	0	6

There are no bond length outliers.

The worst 5 of 60 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	92	ARG	NE-CZ-NH1	13.27	126.94	120.30
1	D	92	ARG	NE-CZ-NH1	12.79	126.69	120.30
1	A	509	PHE	C-N-CD	-10.72	97.03	120.60
1	C	92	ARG	NE-CZ-NH2	-9.66	115.47	120.30
1	F	509	PHE	C-N-CD	-9.58	99.52	120.60

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	509	PHE	Peptide
1	B	509	PHE	Peptide
1	D	670	LYS	Peptide
1	E	422	GLY	Peptide
1	E	509	PHE	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5387	0	5151	38	0
1	B	5424	0	5186	26	0
1	C	5424	0	5186	22	0
1	D	5424	0	5186	28	0
1	E	5432	0	5199	33	0
1	F	5432	0	5199	22	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
3	A	4	0	3	0	0
3	B	4	0	3	0	0
3	C	4	0	3	1	0
3	D	4	0	3	0	0
3	E	4	0	3	1	0
3	F	4	0	3	2	0
4	A	105	0	0	5	0
4	B	56	0	0	1	0
4	C	66	0	0	2	0
4	D	99	0	0	2	0
4	E	48	0	0	2	0
4	F	54	0	0	3	0
All	All	32981	0	31125	159	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 159 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:E:702:ACT:H2	4:E:825:HOH:O	1.81	0.80
1:E:661:ILE:HD11	1:E:682:THR:HB	1.66	0.78
1:F:414:ARG:HD2	4:F:808:HOH:O	1.90	0.71
1:D:92:ARG:HG2	1:D:92:ARG:HH11	1.57	0.69
1:E:230:VAL:HG21	1:E:261:TYR:CZ	2.28	0.69

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	675/712 (95%)	650 (96%)	22 (3%)	3 (0%)	34	45
1	B	682/712 (96%)	652 (96%)	26 (4%)	4 (1%)	25	33
1	C	682/712 (96%)	654 (96%)	25 (4%)	3 (0%)	34	45
1	D	682/712 (96%)	654 (96%)	25 (4%)	3 (0%)	34	45
1	E	683/712 (96%)	649 (95%)	29 (4%)	5 (1%)	22	29
1	F	683/712 (96%)	651 (95%)	28 (4%)	4 (1%)	25	33
All	All	4087/4272 (96%)	3910 (96%)	155 (4%)	22 (0%)	29	39

5 of 22 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	510	PRO
1	B	510	PRO
1	D	663	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	E	493	TYR
1	E	510	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	563/592 (95%)	555 (99%)	8 (1%)	67 78
1	B	567/592 (96%)	555 (98%)	12 (2%)	53 67
1	C	567/592 (96%)	559 (99%)	8 (1%)	67 78
1	D	567/592 (96%)	557 (98%)	10 (2%)	59 73
1	E	568/592 (96%)	554 (98%)	14 (2%)	47 61
1	F	568/592 (96%)	554 (98%)	14 (2%)	47 61
All	All	3400/3552 (96%)	3334 (98%)	66 (2%)	57 71

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	72	VAL
1	D	617	ARG
1	F	494	TRP
1	D	92	ARG
1	D	360	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	602	HIS
1	F	374	HIS
1	D	374	HIS
1	B	374	HIS
1	E	374	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACT	C	702	-	1,3,3	0.64	0	0,3,3	0.00	-
3	ACT	E	702	-	1,3,3	0.28	0	0,3,3	0.00	-
3	ACT	A	702	-	1,3,3	1.02	0	0,3,3	0.00	-
3	ACT	B	702	-	1,3,3	2.41	1 (100%)	0,3,3	0.00	-
3	ACT	D	702	-	1,3,3	0.96	0	0,3,3	0.00	-
3	ACT	F	702	-	1,3,3	0.75	0	0,3,3	0.00	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	702	ACT	CH3-C	2.41	1.51	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	702	ACT	1	0
3	E	702	ACT	1	0
3	F	702	ACT	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	679/712 (95%)	-0.19	1 (0%) 95 98	15, 25, 44, 81	0
1	B	684/712 (96%)	-0.12	4 (0%) 89 93	19, 30, 49, 82	0
1	C	684/712 (96%)	-0.05	5 (0%) 87 91	19, 32, 52, 80	0
1	D	684/712 (96%)	-0.17	2 (0%) 94 97	16, 25, 45, 84	0
1	E	684/712 (96%)	0.02	7 (1%) 82 87	21, 34, 52, 87	0
1	F	684/712 (96%)	0.04	3 (0%) 92 96	21, 35, 56, 81	0
All	All	4099/4272 (95%)	-0.08	22 (0%) 91 94	15, 30, 51, 87	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	11	ILE	4.4
1	E	7	LEU	4.0
1	E	2	THR	3.8
1	B	7	LEU	3.7
1	E	6	LEU	3.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	ACT	D	702	4/4	0.84	0.18	34,37,40,45	0
3	ACT	B	702	4/4	0.85	0.16	35,38,40,43	0
3	ACT	C	702	4/4	0.88	0.17	41,42,44,49	0
3	ACT	A	702	4/4	0.88	0.20	35,37,39,43	0
3	ACT	F	702	4/4	0.90	0.13	41,43,44,45	0
3	ACT	E	702	4/4	0.96	0.13	30,32,32,34	0
2	ZN	D	701	1/1	0.99	0.10	27,27,27,27	0
2	ZN	E	701	1/1	0.99	0.09	37,37,37,37	0
2	ZN	C	701	1/1	0.99	0.09	34,34,34,34	0
2	ZN	A	701	1/1	0.99	0.10	24,24,24,24	0
2	ZN	F	701	1/1	0.99	0.09	36,36,36,36	0
2	ZN	B	701	1/1	0.99	0.08	32,32,32,32	0

6.5 Other polymers [i](#)

There are no such residues in this entry.