



# wwPDB X-ray Structure Validation Summary Report

Feb 12, 2024 – 12:07 AM EST

PDB ID : 3EOE  
Title : Crystal Structure of Pyruvate Kinase from *Toxoplasma gondii*, 55.m00007  
Authors : Wernimont, A.K.; Lew, J.; Kozieradzki, I.; Wasney, G.; Hassani, A.; Vedadi, M.; Cossar, D.; Schapiro, M.; Bochkarev, A.; Arrowsmith, C.H.; Bountra, C.; Weigelt, J.; Edwards, A.M.; Hui, R.; Pizarro, J.; Structural Genomics Consortium (SGC)  
Deposited on : 2008-09-26  
Resolution : 2.31 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	511	 5% 79% 14% 8%
1	B	511	 2% 84% 10% 5%
1	C	511	 2% 80% 12% 7%
1	D	511	 3% 65% 9% 25%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

<b>Mol</b>	<b>Type</b>	<b>Chain</b>	<b>Res</b>	<b>Chirality</b>	<b>Geometry</b>	<b>Clashes</b>	<b>Electron density</b>
2	GOL	B	512	-	-	X	-

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 14040 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pyruvate kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	472	Total 3464	C 2179	N 603	O 654	S 28	0	1	0
1	B	483	Total 3617	C 2273	N 632	O 684	S 28	0	2	0
1	C	475	Total 3510	C 2201	N 621	O 661	S 27	3	1	0
1	D	382	Total 2792	C 1762	N 485	O 519	S 26	3	0	0

There are 72 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP Q969A2
A	2	HIS	-	expression tag	UNP Q969A2
A	3	HIS	-	expression tag	UNP Q969A2
A	4	HIS	-	expression tag	UNP Q969A2
A	5	HIS	-	expression tag	UNP Q969A2
A	6	HIS	-	expression tag	UNP Q969A2
A	7	HIS	-	expression tag	UNP Q969A2
A	8	SER	-	expression tag	UNP Q969A2
A	9	SER	-	expression tag	UNP Q969A2
A	10	GLY	-	expression tag	UNP Q969A2
A	11	ARG	-	expression tag	UNP Q969A2
A	12	GLU	-	expression tag	UNP Q969A2
A	13	ASN	-	expression tag	UNP Q969A2
A	14	LEU	-	expression tag	UNP Q969A2
A	15	TYR	-	expression tag	UNP Q969A2
A	16	PHE	-	expression tag	UNP Q969A2
A	17	GLN	-	expression tag	UNP Q969A2
A	18	GLY	-	expression tag	UNP Q969A2
B	1	MET	-	expression tag	UNP Q969A2
B	2	HIS	-	expression tag	UNP Q969A2
B	3	HIS	-	expression tag	UNP Q969A2

*Continued on next page...*

*Continued from previous page...*


Chain	Residue	Modelled	Actual	Comment	Reference
B	4	HIS	-	expression tag	UNP Q969A2
B	5	HIS	-	expression tag	UNP Q969A2
B	6	HIS	-	expression tag	UNP Q969A2
B	7	HIS	-	expression tag	UNP Q969A2
B	8	SER	-	expression tag	UNP Q969A2
B	9	SER	-	expression tag	UNP Q969A2
B	10	GLY	-	expression tag	UNP Q969A2
B	11	ARG	-	expression tag	UNP Q969A2
B	12	GLU	-	expression tag	UNP Q969A2
B	13	ASN	-	expression tag	UNP Q969A2
B	14	LEU	-	expression tag	UNP Q969A2
B	15	TYR	-	expression tag	UNP Q969A2
B	16	PHE	-	expression tag	UNP Q969A2
B	17	GLN	-	expression tag	UNP Q969A2
B	18	GLY	-	expression tag	UNP Q969A2
C	1	MET	-	expression tag	UNP Q969A2
C	2	HIS	-	expression tag	UNP Q969A2
C	3	HIS	-	expression tag	UNP Q969A2
C	4	HIS	-	expression tag	UNP Q969A2
C	5	HIS	-	expression tag	UNP Q969A2
C	6	HIS	-	expression tag	UNP Q969A2
C	7	HIS	-	expression tag	UNP Q969A2
C	8	SER	-	expression tag	UNP Q969A2
C	9	SER	-	expression tag	UNP Q969A2
C	10	GLY	-	expression tag	UNP Q969A2
C	11	ARG	-	expression tag	UNP Q969A2
C	12	GLU	-	expression tag	UNP Q969A2
C	13	ASN	-	expression tag	UNP Q969A2
C	14	LEU	-	expression tag	UNP Q969A2
C	15	TYR	-	expression tag	UNP Q969A2
C	16	PHE	-	expression tag	UNP Q969A2
C	17	GLN	-	expression tag	UNP Q969A2
C	18	GLY	-	expression tag	UNP Q969A2
D	1	MET	-	expression tag	UNP Q969A2
D	2	HIS	-	expression tag	UNP Q969A2
D	3	HIS	-	expression tag	UNP Q969A2
D	4	HIS	-	expression tag	UNP Q969A2
D	5	HIS	-	expression tag	UNP Q969A2
D	6	HIS	-	expression tag	UNP Q969A2
D	7	HIS	-	expression tag	UNP Q969A2
D	8	SER	-	expression tag	UNP Q969A2
D	9	SER	-	expression tag	UNP Q969A2

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	10	GLY	-	expression tag	UNP Q969A2
D	11	ARG	-	expression tag	UNP Q969A2
D	12	GLU	-	expression tag	UNP Q969A2
D	13	ASN	-	expression tag	UNP Q969A2
D	14	LEU	-	expression tag	UNP Q969A2
D	15	TYR	-	expression tag	UNP Q969A2
D	16	PHE	-	expression tag	UNP Q969A2
D	17	GLN	-	expression tag	UNP Q969A2
D	18	GLY	-	expression tag	UNP Q969A2

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	C	1	Total C O 6 3 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	166	Total O 166 166	0	0
3	B	195	Total O 195 195	0	0

Continued on next page...


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	C	190	Total 190	O 190	0	0
3	D	88	Total 88	O 88	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Pyruvate kinase




- Molecule 1: Pyruvate kinase




- Molecule 1: Pyruvate kinase







● Molecule 1: Pyruvate kinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.04Å 130.68Å 113.85Å 90.00° 117.40° 90.00°	Depositor
Resolution (Å)	42.68 – 2.31 43.56 – 2.31	Depositor EDS
% Data completeness (in resolution range)	90.9 (42.68-2.31) 90.9 (43.56-2.31)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	0.05	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.53 (at 2.32Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.206 , 0.253 0.201 , 0.246	Depositor DCC
$R_{free}$ test set	5071 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	47.3	Xtrriage
Anisotropy	0.065	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 41.1	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.017 for h,-k,-h-l	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	14040	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/3514	0.56	0/4776
1	B	0.39	0/3668	0.55	0/4982
1	C	0.39	0/3557	0.57	1/4833 (0.0%)
1	D	0.41	1/2826 (0.0%)	0.54	0/3836
All	All	0.39	1/13565 (0.0%)	0.56	1/18427 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	82	GLN	CG-CD	9.57	1.73	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	89	LYS	CB-CG-CD	5.27	125.31	111.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3464	0	3431	37	0
1	B	3617	0	3647	37	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	3510	0	3492	31	0
1	D	2792	0	2759	29	0
2	B	12	0	16	9	0
2	C	6	0	8	0	0
3	A	166	0	0	0	0
3	B	195	0	0	2	0
3	C	190	0	0	3	0
3	D	88	0	0	1	0
All	All	14040	0	13353	124	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 124 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:445:ILE:HG23	1:B:456:THR:HG21	1.35	1.07
1:B:66:ARG:HH12	2:B:512:GOL:H32	1.12	1.05
1:B:43:THR:HG21	2:B:512:GOL:H11	1.49	0.95
1:B:347:SER:HB3	2:B:512:GOL:H31	1.50	0.93
1:B:152:VAL:CG1	1:B:169:VAL:HG21	2.03	0.88

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	461/511 (90%)	449 (97%)	11 (2%)	1 (0%)	47 58
1	B	477/511 (93%)	471 (99%)	6 (1%)	0	100 100
1	C	466/511 (91%)	454 (97%)	11 (2%)	1 (0%)	47 58

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	D	364/511 (71%)	357 (98%)	7 (2%)	0	100	100
All	All	1768/2044 (86%)	1731 (98%)	35 (2%)	2 (0%)	51	63

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	313	THR
1	C	313	THR

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	355/428 (83%)	342 (96%)	13 (4%)	34	47
1	B	385/428 (90%)	370 (96%)	15 (4%)	32	45
1	C	362/428 (85%)	344 (95%)	18 (5%)	24	34
1	D	283/428 (66%)	274 (97%)	9 (3%)	39	53
All	All	1385/1712 (81%)	1330 (96%)	55 (4%)	31	44

5 of 55 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	89	LYS
1	C	208	LYS
1	D	503	ASN
1	D	414	LEU
1	C	108	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	215	ASN
1	D	36	HIS

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	82	GLN
1	C	449	GLN
1	C	36	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	B	513	-	5,5,5	0.35	0	5,5,5	0.37	0
2	GOL	C	512	-	5,5,5	0.35	0	5,5,5	0.21	0
2	GOL	B	512	-	5,5,5	0.34	0	5,5,5	0.43	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	513	-	-	3/4/4/4	-
2	GOL	C	512	-	-	0/4/4/4	-
2	GOL	B	512	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	513	GOL	O1-C1-C2-C3
2	B	512	GOL	C1-C2-C3-O3
2	B	513	GOL	O1-C1-C2-O2
2	B	512	GOL	O2-C2-C3-O3
2	B	513	GOL	C1-C2-C3-O3

There are no ring outliers.

1 monomer is involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	512	GOL	9	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	472/511 (92%)	0.20	24 (5%) 28 35	30, 35, 41, 47	12 (2%)
1	B	483/511 (94%)	0.09	12 (2%) 57 64	30, 35, 39, 44	9 (1%)
1	C	475/511 (92%)	0.01	10 (2%) 63 70	30, 35, 40, 42	13 (2%)
1	D	382/511 (74%)	0.09	13 (3%) 45 52	30, 35, 41, 53	15 (3%)
All	All	1812/2044 (88%)	0.10	59 (3%) 46 53	30, 35, 40, 53	49 (2%)

The worst 5 of 59 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	385	LEU	7.5
1	A	49	TRP	4.8
1	D	377	PRO	4.4
1	D	380	TYR	4.0
1	C	409	ASN	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GOL	C	512	6/6	0.73	0.33	80,81,81,81	0
2	GOL	B	512	6/6	0.85	0.19	40,41,43,43	0
2	GOL	B	513	6/6	0.89	0.26	66,67,67,67	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.