



wwPDB X-ray Structure Validation Summary Report ⓘ

May 21, 2020 – 08:12 pm BST

PDB ID : 6EPG
Title : Structure of the epsilon_1 / zeta_1 antitoxin / toxin system from *Neisseria gonorrhoeae*.
Authors : Rocker, A.; Meinhart, A.
Deposited on : 2017-10-11
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.







Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	61	
1	C	61	
1	E	61	
1	G	61	
2	B	401	
2	D	401	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	401	 95% 
2	H	401	 95% 

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 14291 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Epsilon_1 antitoxin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	54	442	275	83	80	1	3	0	0	0
1	C	56	458	285	85	84	1	3	0	0	0
1	E	54	442	275	83	80	1	3	0	0	0
1	G	54	442	275	83	80	1	3	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MSE	-	initiating methionine	UNP D5K9E3
C	1	MSE	-	initiating methionine	UNP D5K9E3
E	1	MSE	-	initiating methionine	UNP D5K9E3
G	1	MSE	-	initiating methionine	UNP D5K9E3


- Molecule 2 is a protein called Zeta_1 toxin.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
2	B	390	3045	1896	531	607	3	8	0	0	0
2	D	394	3075	1914	538	612	3	8	0	0	0
2	F	395	3083	1919	539	613	3	9	0	0	0
2	H	395	3084	1921	539	612	3	9	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	1	MSE	-	initiating methionine	UNP D5K9G7
B	115	ALA	LYS	engineered mutation	UNP D5K9G7
D	1	MSE	-	initiating methionine	UNP D5K9G7
D	115	ALA	LYS	engineered mutation	UNP D5K9G7
F	1	MSE	-	initiating methionine	UNP D5K9G7
F	115	ALA	LYS	engineered mutation	UNP D5K9G7
H	1	MSE	-	initiating methionine	UNP D5K9G7
H	115	ALA	LYS	engineered mutation	UNP D5K9G7

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		
3	D	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total 5	O 4	S 1	0	0
3	D	1	Total 5	O 4	S 1	0	0
3	D	1	Total 5	O 4	S 1	0	0
3	F	1	Total 5	O 4	S 1	0	0
3	F	1	Total 5	O 4	S 1	0	0
3	F	1	Total 5	O 4	S 1	0	0
3	F	1	Total 5	O 4	S 1	0	0
3	F	1	Total 5	O 4	S 1	0	0
3	F	1	Total 5	O 4	S 1	0	0
3	F	1	Total 5	O 4	S 1	0	0
3	H	1	Total 5	O 4	S 1	0	0
3	H	1	Total 5	O 4	S 1	0	0
3	H	1	Total 5	O 4	S 1	0	0
3	H	1	Total 5	O 4	S 1	0	0
3	H	1	Total 5	O 4	S 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	O 1	0	0
4	B	21	Total 21	O 21	0	0
4	C	1	Total 1	O 1	0	0
4	D	23	Total 23	O 23	0	0
4	F	38	Total 38	O 38	0	0


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	26	Total	O	0	0
			26	26		


- Molecule 2: Zeta_1 toxin

Chain D:  93% 5%



- Molecule 2: Zeta_1 toxin

Chain F:  95%



- Molecule 2: Zeta_1 toxin

Chain H:  95%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	79.76Å 149.58Å 125.09Å 90.00° 94.82° 90.00°	Depositor
Resolution (Å)	48.91 – 2.40 48.89 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.91-2.40) 99.7 (48.89-2.40)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.38 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.226 , 0.270 0.233 , 0.273	Depositor DCC
R_{free} test set	5687 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	45.3	Xtrriage
Anisotropy	0.362	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 28.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	14291	wwPDB-VP
Average B, all atoms (Å ²)	51.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.95 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.5363e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/444	0.76	0/586
1	C	0.54	0/461	0.75	0/610
1	E	0.52	0/444	0.76	0/586
1	G	0.59	0/444	0.80	0/586
2	B	0.53	0/3074	0.70	0/4117
2	D	0.52	0/3105	0.71	1/4158 (0.0%)
2	F	0.53	0/3113	0.70	0/4168
2	H	0.51	0/3113	0.70	0/4165
All	All	0.52	0/14198	0.71	1/18976 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	382	ASN	N-CA-C	-5.76	95.45	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	442	0	457	1	0
1	C	458	0	470	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	442	0	457	2	0
1	G	442	0	457	1	0
2	B	3045	0	3036	14	0
2	D	3075	0	3072	11	0
2	F	3083	0	3084	9	0
2	H	3084	0	3090	7	0
3	B	30	0	0	0	0
3	D	25	0	0	0	0
3	F	30	0	0	0	0
3	H	25	0	0	0	0
4	A	1	0	0	0	0
4	B	21	0	0	0	0
4	C	1	0	0	0	0
4	D	23	0	0	0	0
4	F	38	0	0	0	0
4	H	26	0	0	0	0
All	All	14291	0	14123	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 41 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:294:VAL:HG13	2:B:316:MSE:HE1	1.69	0.74
1:A:58:ASN:O	2:B:28:ARG:NH2	2.21	0.74
2:D:285:GLN:HG2	2:D:298:THR:HG22	1.80	0.63
2:B:271:MSE:HE2	2:B:276:VAL:HG23	1.80	0.63
2:F:363:ASN:HD22	2:F:384:GLY:HA2	1.62	0.63

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	52/61 (85%)	51 (98%)	1 (2%)	0	100	100
1	C	54/61 (88%)	51 (94%)	3 (6%)	0	100	100
1	E	52/61 (85%)	52 (100%)	0	0	100	100
1	G	52/61 (85%)	51 (98%)	1 (2%)	0	100	100
2	B	386/401 (96%)	381 (99%)	5 (1%)	0	100	100
2	D	392/401 (98%)	387 (99%)	5 (1%)	0	100	100
2	F	393/401 (98%)	386 (98%)	7 (2%)	0	100	100
2	H	391/401 (98%)	383 (98%)	8 (2%)	0	100	100
All	All	1772/1848 (96%)	1742 (98%)	30 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	46/49 (94%)	46 (100%)	0	100	100
1	C	48/49 (98%)	48 (100%)	0	100	100
1	E	46/49 (94%)	46 (100%)	0	100	100
1	G	46/49 (94%)	46 (100%)	0	100	100
2	B	332/333 (100%)	327 (98%)	5 (2%)	65	80
2	D	335/333 (101%)	332 (99%)	3 (1%)	78	90
2	F	336/333 (101%)	333 (99%)	3 (1%)	78	90
2	H	336/333 (101%)	331 (98%)	5 (2%)	65	80
All	All	1525/1528 (100%)	1509 (99%)	16 (1%)	76	88

5 of 16 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	D	203	ARG
2	F	141	SER
2	H	124	MSE
2	D	120	LYS
2	H	167	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	325	GLN
2	D	100	GLN
2	D	148	GLN
2	F	148	GLN
2	F	363	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	D	501	-	4,4,4	0.21	0	6,6,6	0.72	0
3	SO4	B	502	-	4,4,4	0.40	0	6,6,6	0.25	0
3	SO4	H	501	-	4,4,4	0.34	0	6,6,6	0.48	0
3	SO4	D	505	-	4,4,4	0.34	0	6,6,6	0.36	0
3	SO4	D	502	-	4,4,4	0.40	0	6,6,6	0.36	0
3	SO4	F	506	-	4,4,4	0.46	0	6,6,6	0.49	0
3	SO4	B	503	-	4,4,4	0.26	0	6,6,6	0.31	0
3	SO4	D	503	-	4,4,4	0.30	0	6,6,6	0.31	0
3	SO4	B	504	-	4,4,4	0.42	0	6,6,6	0.15	0
3	SO4	F	503	-	4,4,4	0.52	0	6,6,6	0.16	0
3	SO4	D	504	-	4,4,4	0.43	0	6,6,6	0.21	0
3	SO4	B	505	-	4,4,4	0.37	0	6,6,6	0.23	0
3	SO4	B	501	-	4,4,4	0.37	0	6,6,6	0.78	0
3	SO4	B	506	-	4,4,4	0.39	0	6,6,6	0.41	0
3	SO4	F	501	-	4,4,4	0.19	0	6,6,6	0.51	0
3	SO4	H	502	-	4,4,4	0.43	0	6,6,6	0.34	0
3	SO4	H	505	-	4,4,4	0.34	0	6,6,6	0.49	0
3	SO4	F	505	-	4,4,4	0.39	0	6,6,6	0.14	0
3	SO4	F	502	-	4,4,4	0.33	0	6,6,6	0.28	0
3	SO4	H	503	-	4,4,4	0.39	0	6,6,6	0.52	0
3	SO4	H	504	-	4,4,4	0.41	0	6,6,6	0.18	0
3	SO4	F	504	-	4,4,4	0.31	0	6,6,6	0.26	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	51/61 (83%)	0.11	1 (1%) 65 63	40, 62, 90, 110	0
1	C	53/61 (86%)	0.25	3 (5%) 23 22	41, 64, 101, 110	0
1	E	51/61 (83%)	-0.06	0 100 100	35, 55, 78, 92	0
1	G	51/61 (83%)	0.33	4 (7%) 13 11	35, 63, 96, 105	0
2	B	382/401 (95%)	-0.11	1 (0%) 94 93	32, 48, 71, 87	0
2	D	386/401 (96%)	-0.03	4 (1%) 82 80	31, 47, 71, 101	0
2	F	386/401 (96%)	-0.11	1 (0%) 94 93	28, 44, 69, 117	0
2	H	386/401 (96%)	-0.10	0 100 100	29, 45, 69, 101	0
All	All	1746/1848 (94%)	-0.06	14 (0%) 86 84	28, 47, 77, 117	0

The worst 5 of 14 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	384	GLY	5.5
1	G	30	ALA	4.0
2	D	382	ASN	3.8
1	A	7	GLN	3.7
1	G	7	GLN	3.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SO4	H	504	5/5	0.91	0.18	84,87,88,95	0
3	SO4	H	503	5/5	0.93	0.17	66,71,82,84	0
3	SO4	B	505	5/5	0.94	0.19	87,87,92,94	0
3	SO4	D	504	5/5	0.95	0.18	78,84,86,87	0
3	SO4	D	502	5/5	0.95	0.14	69,69,71,72	0
3	SO4	H	502	5/5	0.95	0.11	58,59,64,65	0
3	SO4	B	504	5/5	0.95	0.16	72,75,76,78	0
3	SO4	F	503	5/5	0.95	0.14	65,66,69,70	0
3	SO4	F	504	5/5	0.95	0.11	67,75,76,79	0
3	SO4	F	505	5/5	0.96	0.14	62,68,72,75	0
3	SO4	B	503	5/5	0.96	0.13	61,64,70,71	0
3	SO4	B	506	5/5	0.96	0.13	53,53,56,58	0
3	SO4	B	502	5/5	0.96	0.09	68,71,72,77	0
3	SO4	F	506	5/5	0.97	0.11	51,52,58,60	0
3	SO4	F	502	5/5	0.97	0.16	62,65,69,72	0
3	SO4	D	505	5/5	0.97	0.14	50,54,56,60	0
3	SO4	D	503	5/5	0.97	0.11	66,72,80,85	0
3	SO4	H	505	5/5	0.97	0.11	55,55,57,58	0
3	SO4	F	501	5/5	0.99	0.15	41,43,44,46	0
3	SO4	D	501	5/5	0.99	0.14	42,42,46,48	0
3	SO4	B	501	5/5	0.99	0.13	40,40,40,43	0
3	SO4	H	501	5/5	0.99	0.13	43,44,46,47	0

6.5 Other polymers [i](#)

There are no such residues in this entry.