



wwPDB X-ray Structure Validation Summary Report

Feb 18, 2024 – 12:26 PM EST

PDB ID : 4ESV
Title : A New Twist on the Translocation Mechanism of Helicases from the Structure of DnaB with its Substrates
Authors : Itsathitphaisarn, O.; Wing, R.A.; Eliason, W.K.; Wang, J.; Steitz, T.A.
Deposited on : 2012-04-23
Resolution : 3.20 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1133 (3.20-3.20)
Clashscore	141614	1253 (3.20-3.20)
Ramachandran outliers	138981	1234 (3.20-3.20)
Sidechain outliers	138945	1233 (3.20-3.20)
RSRZ outliers	127900	1095 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	V	14	
2	W	13	
3	A	454	
3	B	454	
3	C	454	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain	
3	D	454		62% 27% 6% • 5%
3	E	454		61% 25% 5% • 8%
3	F	454		62% 24% 6% • 7%
3	G	454		66% 23% • 7%
3	H	454		65% 23% 5% 6%
3	I	454		61% 27% • 8%
3	J	454		62% 25% • 8%
3	K	454		58% 30% • 7%
3	L	454		65% 24% • 8%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	MES	C	505	-	-	-	X
4	MES	W	101	-	-	-	X
5	CA	G	503	-	-	-	X
6	GDP	I	501	-	-	X	-
7	ALF	A	504	-	-	X	-

2 Entry composition [i](#)

There are 8 unique types of molecules in this entry. The entry contains 40296 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*T P*TP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	V	14	280	140	28	98	14	0	0	0

- Molecule 2 is a DNA chain called 5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*T P*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	W	13	260	130	26	91	13	0	0	0

- Molecule 3 is a protein called Replicative helicase.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	418	3255	2036	570	636	13	8	1	0
3	B	430	3350	2093	591	652	14	0	0	0
3	C	434	3367	2104	591	658	14	0	0	0
3	D	431	3333	2084	583	652	14	0	0	0
3	E	418	3243	2028	565	637	13	0	0	0
3	F	421	3274	2049	569	643	13	0	0	0
3	G	421	3259	2037	568	641	13	0	0	0
3	H	425	3283	2052	572	645	14	0	0	0
3	I	418	3245	2034	561	637	13	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	J	416	Total	C	N	O	S	0	0	0
			3237	2025	563	636	13			
3	K	421	Total	C	N	O	S	0	0	0
			3267	2044	568	642	13			
3	L	419	Total	C	N	O	S	0	0	0
			3246	2031	565	637	13			

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	W	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	E	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	I	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total	Ca	0	0
			5	5		
5	B	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total Ca 1 1	0	0
5	D	2	Total Ca 2 2	0	0
5	E	1	Total Ca 1 1	0	0
5	F	1	Total Ca 1 1	0	0
5	G	1	Total Ca 1 1	0	0
5	I	1	Total Ca 1 1	0	0
5	L	1	Total Ca 1 1	0	0

- Molecule 6 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C N O P 28 10 5 11 2	0	0
6	C	1	Total C N O P 28 10 5 11 2	0	0
6	D	1	Total C N O P 28 10 5 11 2	0	0
6	E	1	Total C N O P 28 10 5 11 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
6	F	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
6	G	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
6	I	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
6	J	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
6	K	1	Total	C	N	O	P	0	0
			28	10	5	11	2		
6	L	1	Total	C	N	O	P	0	0
			28	10	5	11	2		

- Molecule 7 is TETRAFLUOROALUMINATE ION (three-letter code: ALF) (formula: AlF_4).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	1	Total	Al F	0	0
			5	1 4		
7	C	1	Total	Al F	0	0
			5	1 4		
7	C	1	Total	Al F	0	0
			5	1 4		
7	D	1	Total	Al F	0	0
			5	1 4		
7	E	1	Total	Al F	0	0
			5	1 4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	G	1	Total 5	Al 1	F 4	0	0
7	I	1	Total 5	Al 1	F 4	0	0
7	J	1	Total 5	Al 1	F 4	0	0
7	K	1	Total 5	Al 1	F 4	0	0
7	L	1	Total 5	Al 1	F 4	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total 1	O 1	0	0
8	B	1	Total 1	O 1	0	0
8	D	1	Total 1	O 1	0	0
8	E	1	Total 1	O 1	0	0
8	K	1	Total 1	O 1	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.




- Molecule 1: 5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*T)-3'



- Molecule 2: 5'-D(P*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*TP*T)-3'




- Molecule 3: Replicative helicase




- Molecule 3: Replicative helicase







• Molecule 3: Replicative helicase




• Molecule 3: Replicative helicase







• Molecule 3: Replicative helicase




• Molecule 3: Replicative helicase




• Molecule 3: Replicative helicase




• Molecule 3: Replicative helicase




• Molecule 3: Replicative helicase







• Molecule 3: Replicative helicase




• Molecule 3: Replicative helicase





● Molecule 3: Replicative helicase



4 Data and refinement statistics

Property	Value	Source
Space group	P 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	149.12Å 180.32Å 279.13Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.92 – 3.20 47.92 – 3.20	Depositor EDS
% Data completeness (in resolution range)	87.8 (47.92-3.20) 87.8 (47.92-3.20)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.32 (at 3.19Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.242 , 0.289 0.239 , 0.281	Depositor DCC
R_{free} test set	5442 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	94.6	Xtrriage
Anisotropy	0.042	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.26 , 83.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	40296	wwPDB-VP
Average B, all atoms (Å ²)	140.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.00% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MES, CA, ALF, GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	V	1.30	2/307 (0.7%)	2.43	29/472 (6.1%)
2	W	1.61	2/285 (0.7%)	2.37	18/438 (4.1%)
3	A	0.61	0/3295	0.77	3/4450 (0.1%)
3	B	0.61	1/3390 (0.0%)	0.82	2/4576 (0.0%)
3	C	0.58	0/3408	0.75	2/4607 (0.0%)
3	D	0.52	0/3372	0.72	0/4557
3	E	0.54	0/3279	0.73	1/4428 (0.0%)
3	F	0.54	0/3312	0.74	2/4475 (0.0%)
3	G	0.40	0/3296	0.61	1/4454 (0.0%)
3	H	0.44	0/3319	0.64	3/4482 (0.1%)
3	I	0.43	0/3283	0.62	1/4437 (0.0%)
3	J	0.43	0/3273	0.65	2/4420 (0.0%)
3	K	0.46	0/3305	0.68	3/4467 (0.1%)
3	L	0.41	0/3284	0.62	0/4437
All	All	0.53	5/40408 (0.0%)	0.76	67/54700 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	A	0	4
3	B	0	5
3	C	0	1
3	D	0	2
3	E	0	1
3	F	0	1
3	I	0	1
3	J	0	1
3	K	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
All	All	0	17

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	421	ASN	CB-CG	5.99	1.64	1.51
2	W	6	DT	C1'-N1	5.61	1.56	1.49
1	V	12	DT	C1'-N1	5.56	1.56	1.49
2	W	11	DT	N1-C2	5.10	1.42	1.38
1	V	8	DT	C3'-O3'	5.03	1.50	1.44

The worst 5 of 67 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	V	8	DT	O4'-C1'-N1	12.56	116.79	108.00
2	W	7	DT	O4'-C1'-N1	10.63	115.44	108.00
2	W	4	DT	C4-C5-C7	10.36	125.22	119.00
2	W	11	DT	O4'-C4'-C3'	-9.32	100.41	106.00
2	W	10	DT	C1'-O4'-C4'	-9.01	101.09	110.10

There are no chirality outliers.

5 of 17 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
3	A	166	LEU	Peptide
3	A	167	VAL	Peptide
3	A	184	THR	Peptide
3	A	320	ASP	Peptide
3	B	151	SER	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	V	280	0	169	11	0
2	W	260	0	157	24	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	3255	0	3296	128	0
3	B	3350	0	3382	177	0
3	C	3367	0	3384	118	0
3	D	3333	0	3337	119	0
3	E	3243	0	3262	110	0
3	F	3274	0	3305	105	0
3	G	3259	0	3274	87	0
3	H	3283	0	3282	110	0
3	I	3245	0	3268	118	0
3	J	3237	0	3265	108	0
3	K	3267	0	3287	115	0
3	L	3246	0	3263	89	0
4	C	12	0	12	0	0
4	E	12	0	12	0	0
4	I	12	0	12	0	0
4	W	12	0	12	4	0
5	A	5	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	2	0	0	0	0
5	E	1	0	0	0	0
5	F	1	0	0	0	0
5	G	1	0	0	0	0
5	I	1	0	0	0	0
5	L	1	0	0	0	0
6	A	28	0	12	4	0
6	C	28	0	12	4	0
6	D	28	0	12	3	0
6	E	28	0	12	4	0
6	F	28	0	12	4	0
6	G	28	0	12	5	0
6	I	28	0	12	10	0
6	J	28	0	12	6	0
6	K	28	0	12	3	0
6	L	28	0	12	5	0
7	A	5	0	0	2	0
7	C	10	0	0	0	0
7	D	5	0	0	0	0
7	E	5	0	0	0	0
7	G	5	0	0	0	0
7	I	5	0	0	0	0
7	J	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	K	5	0	0	1	0
7	L	5	0	0	0	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	D	1	0	0	0	0
8	E	1	0	0	0	0
8	K	1	0	0	0	0
All	All	40296	0	40099	1249	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

The worst 5 of 1249 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:J:403:ASN:HB3	3:J:404:LYS:CA	1.63	1.28
3:A:145:ARG:HD2	3:B:310:GLN:NE2	1.58	1.18
3:E:14:ILE:HD13	3:E:15:GLU:H	1.10	1.16
2:W:6:DT:H2''	2:W:7:DT:H5''	1.29	1.14
3:J:403:ASN:CB	3:J:404:LYS:HA	1.75	1.14

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	A	413/454 (91%)	369 (89%)	31 (8%)	13 (3%)	4 26
3	B	424/454 (93%)	365 (86%)	38 (9%)	21 (5%)	2 16
3	C	432/454 (95%)	385 (89%)	29 (7%)	18 (4%)	3 20
3	D	425/454 (94%)	376 (88%)	37 (9%)	12 (3%)	5 29

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	E	412/454 (91%)	373 (90%)	27 (7%)	12 (3%)	4	28
3	F	415/454 (91%)	377 (91%)	27 (6%)	11 (3%)	5	30
3	G	415/454 (91%)	374 (90%)	37 (9%)	4 (1%)	15	54
3	H	419/454 (92%)	375 (90%)	39 (9%)	5 (1%)	13	49
3	I	412/454 (91%)	375 (91%)	34 (8%)	3 (1%)	22	61
3	J	410/454 (90%)	369 (90%)	37 (9%)	4 (1%)	15	54
3	K	415/454 (91%)	354 (85%)	51 (12%)	10 (2%)	6	34
3	L	413/454 (91%)	372 (90%)	36 (9%)	5 (1%)	13	49
All	All	5005/5448 (92%)	4464 (89%)	423 (8%)	118 (2%)	6	34

5 of 118 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	B	97	PRO
3	B	154	LYS
3	B	331	SER
3	B	409	LYS
3	B	410	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	353/386 (92%)	291 (82%)	62 (18%)	2	9
3	B	362/386 (94%)	293 (81%)	69 (19%)	1	8
3	C	361/386 (94%)	305 (84%)	56 (16%)	2	12
3	D	356/386 (92%)	303 (85%)	53 (15%)	3	14
3	E	348/386 (90%)	293 (84%)	55 (16%)	2	12
3	F	355/386 (92%)	298 (84%)	57 (16%)	2	11
3	G	351/386 (91%)	307 (88%)	44 (12%)	4	21
3	H	350/386 (91%)	308 (88%)	42 (12%)	5	22

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	I	350/386 (91%)	307 (88%)	43 (12%)	4	21
3	J	350/386 (91%)	303 (87%)	47 (13%)	4	18
3	K	352/386 (91%)	300 (85%)	52 (15%)	3	14
3	L	349/386 (90%)	307 (88%)	42 (12%)	5	22
All	All	4237/4632 (92%)	3615 (85%)	622 (15%)	3	14

5 of 622 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	I	285	ASN
3	K	340	SER
3	I	373	LYS
3	I	264	ARG
3	J	334	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 124 such sidechains are listed below:

Mol	Chain	Res	Type
3	F	334	ASN
3	K	226	ASN
3	H	49	GLN
3	K	225	GLN
3	L	201	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 38 ligands modelled in this entry, 14 are monoatomic - leaving 24 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	GDP	A	502	5	24,30,30	1.22	2 (8%)	30,47,47	1.73	9 (30%)
6	GDP	I	501	5	24,30,30	0.95	0	30,47,47	1.68	9 (30%)
7	ALF	I	503	-	0,4,4	-	-	-	-	-
6	GDP	K	501	-	24,30,30	0.94	1 (4%)	30,47,47	1.34	4 (13%)
7	ALF	A	504	-	0,4,4	-	-	-	-	-
6	GDP	G	501	-	24,30,30	0.97	1 (4%)	30,47,47	1.27	4 (13%)
4	MES	C	505	-	12,12,12	2.12	1 (8%)	14,16,16	2.66	7 (50%)
7	ALF	K	502	-	0,4,4	-	-	-	-	-
6	GDP	D	501	5	24,30,30	1.22	2 (8%)	30,47,47	1.42	4 (13%)
7	ALF	G	502	-	0,4,4	-	-	-	-	-
6	GDP	L	501	5	24,30,30	0.94	0	30,47,47	1.41	6 (20%)
7	ALF	D	503	-	0,4,4	-	-	-	-	-
7	ALF	C	503	-	0,4,4	-	-	-	-	-
7	ALF	C	504	-	0,4,4	-	-	-	-	-
6	GDP	E	501	5	24,30,30	0.96	1 (4%)	30,47,47	1.49	4 (13%)
4	MES	E	504	-	12,12,12	1.94	1 (8%)	14,16,16	7.30	8 (57%)
7	ALF	J	502	-	0,4,4	-	-	-	-	-
4	MES	I	504	-	12,12,12	1.99	1 (8%)	14,16,16	2.61	7 (50%)
4	MES	W	101	-	12,12,12	2.17	1 (8%)	14,16,16	2.43	7 (50%)
6	GDP	F	501	5	24,30,30	1.01	2 (8%)	30,47,47	1.38	4 (13%)
6	GDP	C	501	5	24,30,30	1.41	4 (16%)	30,47,47	1.57	7 (23%)
7	ALF	E	503	-	0,4,4	-	-	-	-	-
7	ALF	L	502	-	0,4,4	-	-	-	-	-
6	GDP	J	501	-	24,30,30	0.98	1 (4%)	30,47,47	1.34	5 (16%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MES	W	101	-	-	2/6/14/14	0/1/1/1
6	GDP	A	502	5	-	6/12/32/32	0/3/3/3
4	MES	C	505	-	-	5/6/14/14	0/1/1/1
6	GDP	G	501	-	-	3/12/32/32	0/3/3/3
6	GDP	F	501	5	-	5/12/32/32	0/3/3/3
6	GDP	I	501	5	-	3/12/32/32	0/3/3/3
6	GDP	D	501	5	-	4/12/32/32	0/3/3/3
6	GDP	C	501	5	-	6/12/32/32	0/3/3/3
6	GDP	L	501	5	-	2/12/32/32	0/3/3/3
6	GDP	J	501	-	-	4/12/32/32	0/3/3/3
4	MES	I	504	-	-	3/6/14/14	0/1/1/1
6	GDP	E	501	5	-	4/12/32/32	0/3/3/3
4	MES	E	504	-	-	3/6/14/14	0/1/1/1
6	GDP	K	501	-	-	3/12/32/32	0/3/3/3

The worst 5 of 18 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	W	101	MES	C8-S	-7.26	1.67	1.77
4	C	505	MES	C8-S	-7.05	1.67	1.77
4	I	504	MES	C8-S	-6.44	1.68	1.77
4	E	504	MES	C8-S	-6.13	1.68	1.77
6	C	501	GDP	O4'-C1'	4.35	1.47	1.41

The worst 5 of 85 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	E	504	MES	O2S-S-C8	-16.10	87.53	106.92
4	E	504	MES	O1S-S-C8	-15.52	88.23	106.92
4	E	504	MES	O3S-S-C8	-12.44	85.64	105.77
4	I	504	MES	C5-N4-C3	6.32	123.05	108.83
4	E	504	MES	C5-N4-C3	6.25	122.89	108.83

There are no chirality outliers.

5 of 53 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	W	101	MES	C8-C7-N4-C5
4	W	101	MES	N4-C7-C8-S

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
4	C	505	MES	N4-C7-C8-S
4	C	505	MES	C7-C8-S-O1S
4	C	505	MES	C7-C8-S-O3S


There are no ring outliers.


13 monomers are involved in 54 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	502	GDP	4	0
6	I	501	GDP	10	0
6	K	501	GDP	3	0
7	A	504	ALF	2	0
6	G	501	GDP	5	0
7	K	502	ALF	1	0
6	D	501	GDP	3	0
6	L	501	GDP	5	0
6	E	501	GDP	4	0
4	W	101	MES	4	0
6	F	501	GDP	4	0
6	C	501	GDP	4	0
6	J	501	GDP	6	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	V	14/14 (100%)	-0.19	1 (7%) 16 9	82, 91, 181, 202	0
2	W	13/13 (100%)	-0.55	0 100 100	92, 107, 167, 208	0
3	A	418/454 (92%)	-0.18	6 (1%) 75 63	69, 100, 162, 214	3 (0%)
3	B	430/454 (94%)	-0.06	9 (2%) 63 49	66, 111, 181, 236	0
3	C	434/454 (95%)	-0.16	5 (1%) 79 67	66, 102, 157, 213	0
3	D	431/454 (94%)	-0.12	3 (0%) 87 81	67, 109, 171, 229	0
3	E	418/454 (92%)	-0.08	5 (1%) 79 67	68, 108, 164, 221	0
3	F	421/454 (92%)	-0.01	12 (2%) 51 36	66, 108, 167, 212	0
3	G	421/454 (92%)	0.42	29 (6%) 16 9	100, 165, 252, 322	0
3	H	425/454 (93%)	0.39	30 (7%) 16 9	95, 149, 253, 284	0
3	I	418/454 (92%)	0.20	8 (1%) 66 53	92, 158, 248, 326	0
3	J	416/454 (91%)	0.38	23 (5%) 25 14	89, 177, 285, 314	1 (0%)
3	K	421/454 (92%)	0.42	42 (9%) 7 4	77, 169, 283, 310	0
3	L	419/454 (92%)	0.56	43 (10%) 6 4	112, 174, 246, 301	0
All	All	5099/5475 (93%)	0.14	216 (4%) 36 23	66, 134, 228, 326	4 (0%)

The worst 5 of 216 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	H	405	ASP	11.8
3	H	400	ASP	7.5
3	L	270	PRO	7.2
3	H	404	LYS	6.7
3	K	256	GLY	6.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	MES	C	505	12/12	0.64	0.85	232,233,234,235	0
4	MES	W	101	12/12	0.68	0.66	247,250,253,253	0
5	CA	G	503	1/1	0.80	0.48	141,141,141,141	0
5	CA	B	501	1/1	0.81	0.58	147,147,147,147	0
6	GDP	K	501	28/28	0.82	0.24	165,178,204,208	0
5	CA	A	507	1/1	0.84	0.27	134,134,134,134	0
5	CA	A	505	1/1	0.85	0.26	117,117,117,117	0
6	GDP	L	501	28/28	0.86	0.23	129,162,173,177	0
4	MES	I	504	12/12	0.87	0.16	111,121,151,151	0
6	GDP	J	501	28/28	0.89	0.22	140,157,182,186	0
6	GDP	E	501	28/28	0.90	0.23	67,89,110,118	0
5	CA	A	506	1/1	0.90	0.36	140,140,140,140	0
4	MES	E	504	12/12	0.91	0.16	122,127,144,148	0
6	GDP	G	501	28/28	0.92	0.23	118,157,181,186	0
5	CA	A	501	1/1	0.92	0.22	109,109,109,109	0
5	CA	D	504	1/1	0.92	0.16	131,131,131,131	0
6	GDP	F	501	28/28	0.92	0.24	71,90,105,111	0
7	ALF	G	502	5/5	0.92	0.40	120,127,139,141	0
6	GDP	I	501	28/28	0.93	0.23	71,84,103,109	0
7	ALF	J	502	5/5	0.93	0.39	119,120,129,130	0
6	GDP	A	502	28/28	0.95	0.20	67,75,87,90	0
6	GDP	C	501	28/28	0.95	0.24	66,69,83,87	0
6	GDP	D	501	28/28	0.95	0.18	65,81,90,97	0
7	ALF	A	504	5/5	0.96	0.21	84,89,97,97	0
7	ALF	K	502	5/5	0.96	0.31	121,126,137,144	0
7	ALF	E	503	5/5	0.97	0.36	88,88,100,101	0
5	CA	F	502	1/1	0.97	0.37	92,92,92,92	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
5	CA	D	502	1/1	0.98	0.26	90,90,90,90	0
7	ALF	I	503	5/5	0.98	0.28	89,90,95,97	0
7	ALF	D	503	5/5	0.98	0.37	80,88,97,109	0
5	CA	C	502	1/1	0.98	0.41	101,101,101,101	0
7	ALF	L	502	5/5	0.98	0.21	118,121,124,126	0
5	CA	A	503	1/1	0.99	0.28	103,103,103,103	0
5	CA	E	502	1/1	0.99	0.34	110,110,110,110	0
5	CA	I	502	1/1	0.99	0.32	117,117,117,117	0
7	ALF	C	503	5/5	0.99	0.31	66,70,86,88	0
7	ALF	C	504	5/5	0.99	0.32	67,74,92,94	0
5	CA	L	503	1/1	0.99	0.30	155,155,155,155	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around GDP L 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GDP J 501:**




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around GDP F 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around GDP I 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around GDP A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around GDP C 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.