

# Full wwPDB X-ray Structure Validation Report (i)

May 23, 2020 – 06:08 am BST

PDB ID : 5EXA

Title: Small-molecule stabilization of the 14-3-3/Gab2 PPI interface

Authors: Bier, D.; Ottmann, C.

Deposited on : 2015-11-23

Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

Mol Probity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.11

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac: 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.11

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} \text{Whole archive} \\ (\#\text{Entries}) \end{array}$	$egin{aligned}  ext{Similar resolution} \ (\# ext{Entries},  ext{resolution range}( ext{Å})) \end{aligned}$
$R_{free}$	130704	$2580 \ (1.96 - 1.96)$
Clashscore	141614	2705 (1.96-1.96)
Ramachandran outliers	138981	2678 (1.96-1.96)
Sidechain outliers	138945	2678 (1.96-1.96)
RSRZ outliers	127900	2539 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain						
1	A	230	2%	12%	•				
1	В	230	3%	16%	•				
2	С	11	18%	27%	18%	18%	_		
2	D	11	45% 55%		18%	27%			



# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 4107 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called 14-3-3 protein zeta/delta.

Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace
1	Λ	229	Total	С	N	О	S	0	1	0
1	1 A		1830	1148	306	367	9	0		
1	D	230	Total	С	N	О	S	0	0	0
1	I B	230	1820	1141	303	366	10	0		0

• Molecule 2 is a protein called GRB2-associated-binding protein 2.


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace		
2	2 C	0	Total	С	N	О	Р	S	0	0	0
		9	76	44	16	14	1	1			
9	D	11	Total	С	N	О	Р	S	0	0	0
	2   D	D   11		53	19	19	1	1	U	U	

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference	
С	397	GLN	ASN	conflict	UNP Q9UQC2	
D	397	GLN	ASN	conflict	UNP Q9UQC2	


• Molecule 3 is Fusicoccin A-THF derivative (three-letter code: 5SO) (formula: C<sub>30</sub>H<sub>48</sub>O<sub>8</sub>).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 38 30 8	0	0
3	В	1	Total C O 38 30 8	0	0

 $\bullet$  Molecule 4 is BENZOIC ACID (three-letter code: BEZ) (formula:  $\mathrm{C_7H_6O_2}).$ 



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 9 7 2	0	0
4	В	1	Total C O 9 7 2	0	0



• Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total Mg 1 1	0	0

• Molecule 6 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	88	Total O 88 88	0	0
6	В	92	Total O 92 92	0	0
6	С	5	Total O 5 5	0	0
6	D	8	Total O 8 8	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: 14-3-3 protein zeta/delta





# 4 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 21 21 21	Depositor	
Cell constants	72.00Å 83.09Å 111.15Å	Depositor	
a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.00° 90.00° 90.00°	Depositor	
Resolution (Å)	48.87 - 1.95	Depositor	
Resolution (A)	48.87 - 1.95	EDS	
% Data completeness	100.0 (48.87-1.95)	Depositor	
(in resolution range)	99.8 (48.87-1.95)	EDS	
$R_{merge}$	0.04	Depositor	
$R_{sym}$	(Not available)	Depositor	
$< I/\sigma(I) > 1$	4.01 (at 1.95Å)	Xtriage	
Refinement program	REFMAC 5.5.0102	Depositor	
P. P.	0.198 , $0.234$	Depositor	
$R, R_{free}$	0.199 , 0.236	DCC	
$R_{free}$ test set	2460 reflections $(5.00\%)$	wwPDB-VP	
Wilson B-factor (Å <sup>2</sup> )	29.8	Xtriage	
Anisotropy	0.268	Xtriage	
Bulk solvent $k_{sol}(e/Å^3)$ , $B_{sol}(Å^2)$	0.36, 39.0	EDS	
L-test for twinning <sup>2</sup>	$ < L >=0.46, < L^2>=0.29$	Xtriage	
Estimated twinning fraction	No twinning to report.	Xtriage	
$F_o, F_c$ correlation	0.95	EDS	
Total number of atoms	4107	wwPDB-VP	
Average B, all atoms (Å <sup>2</sup> )	31.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 16.05% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of <|L|>,  $< L^2>$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

### 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, BEZ, MG, 5SO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
MIOI		RMSZ	# Z >5	RMSZ	# Z  > 5	
1	A	1.15	0/1858	0.95	0/2501	
1	В	1.11	0/1845	0.92	0/2486	
2	С	0.72	0/65	1.61	$2/84 \ (2.4\%)$	
2	D	0.86	0/82	1.08	0/107	
All	All	1.12	0/3850	0.95	$2/5178 \ (0.0\%)$	

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$\mathbf{Ideal}(^o)$
2	С	392	LEU	CB-CG-CD1	-10.61	92.96	111.00
2	С	392	LEU	CA-CB-CG	6.67	130.65	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1830	0	1810	22	0
1	В	1820	0	1780	30	0
2	С	76	0	76	7	0
2	D	93	0	88	4	0
3	A	38	0	0	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Non-H	H(model)	$oxed{ H(added)   Clashes   Sy}$		Symm-Clashes
3	В	38	0	0	1	0
4	A	9	0	5	0	0
4	В	9	0	5 0		0
5	В	1	0	0	0	0
6	A	88	0	0	3	2
6	В	92	0	0	0	2
6	С	5	0	0	0	0
6	D	8	0	0	0	0
All	All	4107	0	3764	57	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (57) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

A.1 1	A	Interatomic	Clash
Atom-1	Atom-2	${ m distance} \; ({ m \AA})$	overlap(Å)
1:B:87:GLU:HG2	1:B:132:VAL:HG11	1.19	1.09
1:B:87:GLU:HG2	1:B:132:VAL:CG1	1.90	1.00
1:B:224:ASN:HD21	2:C:390:ASN:H	1.17	0.92
1:A:164:HIS:HD2	1:A:166:ILE:H	1.25	0.83
1:A:190:SER:O	1:A:194:THR:HG23	1.83	0.79
1:A:77:GLN:HE21	1:A:80:ARG:NH2	1.82	0.78
2:C:389:ARG:HG2	2:C:389:ARG:HH21	1.46	0.77
1:B:124:ASP:OD1	1:B:149:TYR:OH	2.01	0.77
1:B:49:LYS:HG2	2:C:393:PRO:HG2	1.70	0.73
1:B:112:ALA:CB	1:B:159:GLU:HG2	2.19	0.72
1:B:190:SER:O	1:B:194:THR:HG23	1.90	0.71
1:B:87:GLU:CG	1:B:132:VAL:CG1	2.69	0.70
1:B:49:LYS:HD3	1:B:49:LYS:O	1.93	0.69
1:A:222:ARG:O	1:A:226:THR:HG23	1.93	0.68
1:B:59:TRP:CD1	1:B:87:GLU:HG3	2.29	0.67
1:A:9:LYS:HE3	1:B:82:TYR:CD1	2.32	0.64
1:B:84:GLU:O	1:B:88:THR:HG23	2.01	0.61
1:B:49:LYS:HD3	1:B:49:LYS:C	2.19	0.61
2:C:389:ARG:HG2	2:C:389:ARG:NH2	2.16	0.61
1:B:132:VAL:O	1:B:132:VAL:HG12	2.04	0.58
1:B:87:GLU:CG	1:B:132:VAL:HG11	2.12	0.58
1:B:63:SER:O	1:B:67:GLN:HG2	2.05	0.56
1:A:160:MET:HE2	1:A:164:HIS:CG	2.41	0.55
1:A:164:HIS:CD2	1:A:166:ILE:H	2.15	0.54
1:B:112:ALA:HB2	1:B:159:GLU:HG2	1.90	0.53

Continued on next page...



 $Continued\ from\ previous\ page...$ 

A	_	Interatomic	Clash
Atom-1	Atom-2	${f distance} \; ({f \mathring{A}})$	overlap (Å)
1:B:30:THR:HG21	1:B:100:LEU:HG	1.92	0.52
1:A:217:ILE:HG12	2:D:392:LEU:HD23	1.93	0.51
1:A:160:MET:CE	1:A:164:HIS:CG	2.94	0.50
1:A:217:ILE:HG12	2:D:392:LEU:CD2	2.42	0.49
3:B:302:5SO:CBG	3:B:302:5SO:CAY	2.91	0.49
1:A:77:GLN:HE21	1:A:80:ARG:HH22	1.55	0.49
1:A:206:LEU:HG	1:A:211:TYR:HB2	1.95	0.48
2:D:396:ASP:O	2:D:397:GLN:HB2	2.14	0.48
1:A:12:LEU:HD11	1:B:79:ALA:HA	1.96	0.48
1:B:59:TRP:NE1	1:B:87:GLU:HG3	2.30	0.46
1:B:171:ALA:O	1:B:175:SER:HB3	2.16	0.46
2:C:389:ARG:NH2	2:C:389:ARG:CG	2.79	0.46
1:B:177:PHE:CE2	1:B:182:LEU:HD22	2.52	0.45
1:A:101:LEU:HA	1:A:105:LEU:HB2	1.99	0.45
1:B:87:GLU:CD	1:B:132:VAL:CG1	2.85	0.45
1:A:157:LYS:HG3	6:A:401:HOH:O	2.16	0.45
1:A:160:MET:CE	1:A:164:HIS:CD2	3.00	0.45
2:C:388:ARG:NH1	2:C:389:ARG:NH1	2.65	0.44
1:A:180:GLU:OE1	2:D:389:ARG:NE	2.50	0.44
1:B:4:ASN:O	1:B:8:GLN:HG2	2.18	0.44
1:B:103:LYS:HD3	1:B:104:PHE:CE1	2.53	0.44
1:B:124:ASP:HA	1:B:149:TYR:OH	2.17	0.44
1:A:156:SER:OG	6:A:401:HOH:O	2.09	0.44
1:A:160:MET:HE3	1:A:166:ILE:HB	2.00	0.44
1:B:200:ILE:O	1:B:203:LEU:HD22	2.18	0.43
1:A:77:GLN:NE2	1:A:80:ARG:NH2	2.59	0.43
2:C:388:ARG:HG2	2:C:389:ARG:HD3	2.00	0.43
1:A:103:LYS:NZ	6:A:404:HOH:O	2.31	0.43
1:B:91:ARG:HG2	1:B:129:LEU:CD2	2.49	0.42
1:B:1:MET:HG3	1:B:1:MET:O	2.20	0.41
1:B:59:TRP:HE1	1:B:87:GLU:CG	2.34	0.40
1:A:49:LYS:HD2	1:A:49:LYS:HA	1.85	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$egin{aligned}  ext{Interatomic} \  ext{distance} \ ( ext{Å}) \end{aligned}$	Clash overlap (Å)
6:A:423:HOH:O	6:B:446:HOH:O[4_545]	0.69	1.51
6:A:484:HOH:O	6:B:424:HOH:O[4_555]	1.86	0.34



### 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	3
1	A	$228/230 \ (99\%)$	227 (100%)	1 (0%)	0	100 100	
1	В	$228/230 \ (99\%)$	224~(98%)	4 (2%)	0	100 100	
2	$^{\mathrm{C}}$	6/11 (54%)	6 (100%)	0	0	100 100	
2	D	8/11 (73%)	8 (100%)	0	0	100 100	
All	All	470/482 (98%)	465 (99%)	5 (1%)	0	100 100	

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric Outliers		Percentiles		
1	A	197/200 (98%)	184 (93%)	13 (7%)	16 6		
1	В	193/200 (96%)	177 (92%)	16 (8%)	11 3		
2	С	7/9 (78%)	5 (71%)	2 (29%)	0 0		
2	D	9/9 (100%)	5 (56%)	4 (44%)	0 0		
All	All	406/418 (97%)	371 (91%)	35 (9%)	10 3		

All (35) residues with a non-rotameric sidechain are listed below:

$\mathbf{Mol}$	Chain	Res	Type
1	A	3	LYS
1	A	12	LEU

Continued on next page...



Continued from previous page...

Continued from previous page							
Mol	Chain	Res	Type				
1	A	91	ARG				
1	A	96	ASP				
1	A	115	LYS				
1	A	144	GLN				
1	A	182	LEU				
1	A	194	THR				
1	A	203	LEU				
1	A	204	ASP				
1	A	206	LEU				
1	A	208	GLU				
1	A	229	THR				
1	В	2	ASP				
1	В	49	LYS				
1	В	73	GLU				
1	В	88	THR				
1	В	100	LEU				
1	В	114	SER				
1	В	115	LYS				
1	В	143	ASP				
1	В	144	GLN				
1	В	147	GLN				
1	В	174	PHE				
1	В	182	LEU				
1	В	183	ASN				
1	В	194	THR				
1	В	203	LEU				
1	В	230	SER				
2	С	389	ARG				
2	С	392	LEU				
2	D	388	ARG				
2	D	389	ARG				
2	D	392	LEU				
2	D	396	ASP				

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	4	ASN
1	A	8	GLN
1	A	77	GLN
1	A	146	GLN
1	A	164	HIS

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type
1	В	50	ASN
1	В	146	GLN
1	В	147	GLN
1	В	224	ASN

### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Chain	Chain	Chain	Chain	Res	Link	$\mathbf{B}_{0}$	ond leng	${ m gths}$	В	ond ang	les
WIOI			nes	Lilik	Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2				
2	TPO	С	391	2	8,10,11	1.20	1 (12%)	10,14,16	1.59	2 (20%)				
2	TPO	D	391	2	8,10,11	1.13	1 (12%)	10,14,16	1.22	1 (10%)				

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	TPO	С	391	2	-	1/9/11/13	-
2	TPO	D	391	2	-	2/9/11/13	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	${ m Observed}({ m \AA})$	$\mathbf{Ideal}( exttt{A})$
2	С	391	TPO	P-OG1	2.27	1.63	1.59
2	D	391	TPO	P-OG1	2.09	1.63	1.59



All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	$\mathbf{Z}$	$\mathbf{Observed}(^o)$	$Ideal(^{o})$
2	С	391	TPO	O3P-P-O1P	3.25	123.42	110.68
2	С	391	TPO	CG2-CB-CA	-2.41	108.41	113.16
2	D	391	TPO	CG2-CB-CA	-2.15	108.92	113.16

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	391	TPO	O-C-CA-CB
2	D	391	TPO	CB-OG1-P-O3P
2	D	391	TPO	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

### 5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry (i)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	В	Bond lengths			Bond angles		
MIGI					Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2	
4	BEZ	В	303	-	7,9,9	0.51	0	8,11,11	0.90	0	
3	5SO	A	301	-	39,42,42	2.73	17 (43%)	44,66,66	3.46	20 (45%)	
3	5SO	В	302	-	39,42,42	2.63	14 (35%)	44,66,66	3.57	20 (45%)	
4	BEZ	A	302	-	7,9,9	0.81	0	8,11,11	0.69	0	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral



centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	${ m Res}$	Link	Chirals	Torsions	Rings
4	BEZ	В	303	-	-	0/0/4/4	0/1/1/1
3	5SO	A	301	-	-	5/11/92/92	0/5/5/5
3	5SO	В	302	-	-	2/11/92/92	0/5/5/5
4	BEZ	A	302	-	-	0/0/4/4	0/1/1/1

All (31) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}( ext{\AA})$	Ideal(Å)
3	В	302	5SO	O6-CBK	9.68	1.55	1.43
3	A	301	5SO	O6-CBK	6.79	1.51	1.43
3	В	302	5SO	O4-CBK	6.17	1.51	1.43
3	A	301	5SO	CAZ-CAX	5.75	1.58	1.51
3	В	302	5SO	CAK-CAW	5.59	1.40	1.33
3	A	301	5SO	CBB-CAW	-5.51	1.45	1.51
3	A	301	5SO	O6-C6	5.01	1.50	1.43
3	A	301	5SO	CAK-CAW	4.92	1.39	1.33
3	A	301	5SO	CAL-CAX	-4.57	1.43	1.51
3	A	301	5SO	O4-CBK	4.02	1.48	1.43
3	В	302	5SO	C4-C5	3.40	1.61	1.52
3	A	301	5SO	CBI-CAY	-3.26	1.45	1.50
3	В	302	5SO	CAO-CBL	-3.23	1.50	1.55
3	В	302	5SO	CAL-CAX	-3.14	1.46	1.51
3	A	301	5SO	CBG-CAW	-3.10	1.47	1.51
3	A	301	5SO	C4-C5	3.06	1.61	1.52
3	В	302	5SO	CBB-CAW	-3.04	1.48	1.51
3	A	301	5SO	CAX-CAY	2.71	1.37	1.33
3	A	301	5SO	O5-C5	2.63	1.50	1.44
3	В	302	5SO	CBG-CAW	-2.53	1.48	1.51
3	В	302	5SO	CBG-CBA	2.50	1.58	1.56
3	A	301	5SO	CAF-CBK	2.40	1.56	1.51
3	A	301	5SO	CBG-CBA	2.39	1.57	1.56
3	В	302	5SO	O6-C6	2.38	1.47	1.43
3	В	302	5SO	O3-C3	-2.38	1.37	1.43
3	В	302	5SO	CAZ-CAX	2.26	1.54	1.51
3	В	302	5SO	CAF-CBK	2.19	1.56	1.51
3	A	301	5SO	OAR-CAA	2.17	1.53	1.42
3	A	301	5SO	CAE-CBK	2.05	1.55	1.51
3	В	302	5SO	CAM-CBB	2.04	1.59	1.54
3	A	301	5SO	O1-C1	2.00	1.47	1.41



All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	301	5SO	C1-O5-C5	-12.84	88.49	113.69
3	В	302	5SO	C1-O5-C5	-12.36	89.42	113.69
3	В	302	5SO	C1-C2-C3	-7.49	94.41	110.00
3	A	301	5SO	O1-C1-O5	7.20	130.79	110.67
3	В	302	5SO	O1-C1-O5	7.06	130.40	110.67
3	A	301	5SO	C1-C2-C3	-6.81	95.81	110.00
3	В	302	5SO	CBG-CBA-CBC	-6.04	106.03	112.39
3	A	301	5SO	CAL-CAX-CAZ	-5.85	111.78	120.06
3	В	302	5SO	CBG-CAW-CAK	-5.59	124.63	130.72
3	В	302	5SO	CAN-CBG-CAW	5.46	111.90	101.69
3	A	301	5SO	CAD-CBA-CBG	5.24	119.13	114.85
3	В	302	5SO	O6-CBK-O4	5.02	114.96	110.02
3	В	302	5SO	CAL-CAX-CAY	5.00	115.50	110.63
3	A	301	5SO	CAN-CBG-CAW	4.91	110.87	101.69
3	A	301	5SO	CBG-CBA-CBC	-4.76	107.38	112.39
3	A	301	5SO	O5-C5-C6	4.64	115.94	108.78
3	В	302	5SO	CAD-CBA-CBG	4.56	118.57	114.85
3	A	301	5SO	CBG-CAW-CAK	-4.52	125.80	130.72
3	A	301	5SO	CAL-CAX-CAY	4.20	114.73	110.63
3	В	302	5SO	CAF-CBK-CAE	-4.12	107.92	112.92
3	A	301	5SO	CAM-CBB-CAW	3.91	109.28	101.98
3	В	302	5SO	CAM-CBB-CAW	3.67	108.84	101.98
3	В	302	5SO	CAZ-CAX-CAY	-3.58	124.88	129.59
3	В	302	5SO	O6-C6-C5	-3.58	103.78	110.87
3	В	302	5SO	CAN-CBG-CBA	-3.52	102.74	110.81
3	В	302	5SO	CBA-CBG-CAW	3.33	122.37	118.11
3	В	302	5SO	O5-C5-C6	3.20	113.72	108.78
3	В	302	5SO	CAC-CAZ-CAX	-3.04	106.72	111.35
3	В	302	5SO	O1-C1-C2	2.96	115.78	108.10
3	A	301	5SO	CAZ-CAX-CAY	2.85	133.35	129.59
3	A	301	5SO	O6-C6-C5	-2.80	105.33	110.87
3	A	301	5SO	O5-C5-C4	2.74	115.53	109.75
3	A	301	5SO	CAF-CBK-CAE	-2.65	109.70	112.92
3	A	301	5SO	CAN-CBG-CBA	-2.53	105.01	110.81
3	A	301	5SO	CAB-CAZ-CAX	2.36	114.95	111.35
3	A	301	5SO	C6-O6-CBK	2.35	116.82	114.37
3	A	301	5SO	O2-C2-C1	2.32	115.68	110.05
3	A	301	5SO	CAC-CAZ-CAX	2.23	114.74	111.35
3	В	302	5SO	O5-C5-C4	2.08	114.14	109.75
3	В	302	5SO	CAN-CAM-CBB	2.02	107.00	103.66

There are no chirality outliers.



All (7) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	301	5SO	CAL-CAX-CAZ-CAB
3	A	301	5SO	CAY-CAX-CAZ-CAC
3	A	301	5SO	CAL-CAX-CAZ-CAC
3	В	302	5SO	O5-C1-O1-CBI
3	A	301	5SO	O5-C1-O1-CBI
3	A	301	5SO	CAY-CAX-CAZ-CAB
3	В	302	5SO	OAR-CAP-CBB-CAW


There are no ring outliers.


1 monomer is involved in 1 short contact:

I	Mol	Chain	Res	Type	Clashes	Symm-Clashes
	3	В	302	5SO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

### 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	$\#\mathrm{RSRZ}{>}2$	$\mathbf{OWAB}(\mathrm{\AA}^2)$	Q < 0.9
1	A	$229/230 \ (99\%)$	0.22	5 (2%) 62 70	17, 27, 48, 59	1 (0%)
1	В	$230/230 \; (100\%)$	0.19	6 (2%) 56 65	18, 30, 49, 70	0
2	С	8/11 (72%)	2.00	2 (25%) 0 0	33, 37, 54, 56	3 (37%)
2	D	10/11 (90%)	2.17	5 (50%) 0 0	30, 41, 58, 60	4 (40%)
All	All	477/482 (98%)	0.28	18 (3%) 40 50	17, 29, 50, 70	8 (1%)

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	387	PRO	5.5
2	С	387	PRO	5.0
1	В	1	MET	4.8
1	A	208	GLU	3.8
2	D	396	ASP	3.5
1	В	136	ASP	3.5
2	С	392	LEU	3.3
2	D	395	MET	3.3
2	D	397	GLN	3.3
1	В	132	VAL	3.2
1	A	136	ASP	2.8
1	A	230	SER	2.8
1	A	135	GLY	2.4
1	В	70	GLU	2.4
1	В	230	SER	2.4
1	В	67	GLN	2.2
1	A	3	LYS	2.1
2	D	388	ARG	2.0



### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
2	TPO	С	391	11/12	0.97	0.09	26,30,32,32	0
2	TPO	D	391	11/12	0.98	0.12	23,27,32,33	0

### 6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-factors}}({f \AA}^2)$	Q < 0.9
3	5SO	Α	301	38/38	0.86	0.21	22,44,64,65	0
3	5SO	В	302	38/38	0.91	0.15	27,35,52,54	0
4	BEZ	В	303	9/9	0.98	0.10	21,22,25,26	9
4	BEZ	A	302	9/9	0.98	0.11	21,22,25,25	9
5	MG	В	301	1/1	0.99	0.22	47,47,47,47	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



# Electron density around 5SO A 301: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)

# Electron density around 5SO B 302: 2mF<sub>o</sub>-DF<sub>c</sub> (at 0.7 rmsd) in gray mF<sub>o</sub>-DF<sub>c</sub> (at 3 rmsd) in purple (negative) and green (positive)



# 6.5 Other polymers (i)

There are no such residues in this entry.

